Mireya Fernández-
Fournier, MD
Javier Díaz de Terán, MD
Antonio Tallón Barranco,
MD, PhD
Inmaculada Puertas, MD,
PhD

*Neurol Neuroimmunol
Neuroinflammation*
2014;1:e31; doi: 10.1212/
NXI.0000000000000031

EARLY CERVICAL MYELITIS AFTER HUMAN PAPILLOMA VIRUS VACCINATION

OPEN

In 2006, an adjuvant vaccine against 4 human papilloma-virus (HPV) types was licensed. By 2012, 45 countries had introduced HPV vaccination.¹ HPV encompasses a group of more than 100 viruses, of which at least 13 are high-risk types, responsible for cervical and other anogen-ital cancers.² There are currently 2 vaccines that prevent HPV 16 and 18 infection (known to cause 70% of cer-vical cancers) and development of subsequent cervical precancerous lesions.¹

Case report. A 14-year-old immunocompetent girl noticed her left hand felt weak and numb. These symptoms spread over 5 days to the rest of her arm and ipsilateral thoracic region, abdomen, and leg. Three days earlier she had received a first dose of quadrivalent HPV recombinant vaccine (qHPV, Gardasil). She had not had fever or previous symptoms suggestive of infection. On examination, she had bilateral diminished sensation to light touch, pain, and vibration below C5 level; muscle weakness of her left arm and leg, being able to move joints against resistance (grade 4 according to the Medical Research Council scale for muscle strength); and hyperreflexia bilaterally with nonsustained clonus and a left extensor plantar response. Neurologic examination was otherwise normal.

Brain and cervical spine MRI revealed an isolated C1-C2 hyperintense lesion in T2-weighted images (figure). CSF analysis showed normal glucose (0.49 mg/dL) and protein levels without cells, an increased myelin basic protein (6.4 ng/mL), an increased nonspecific immuno-globulin M ratio, and the presence of oligoclonal bands. Serology and PCR ruled out infectious etiologies, with negative testing for *Borrelia*, *Treponema*, *Listeria*, herpes-viruses 1, 2, and 6, varicella-zoster virus, Epstein-Barr virus, cytomegalovirus, and HIV. Laboratory screening showed normal angiotensin-converting enzyme, vitamin B₁₂, and folate levels and was negative for antinuclear and antitransglutaminase antibodies.

Cervical transverse myelitis was diagnosed. Treat-ment with 1 g IV methylprednisolone was given for 5 days and clinical improvement ensued. Six months later, only mild left hand hypoesthesia and lower

extremity hyperreflexia persisted. Follow-up brain and spinal MRI revealed an improvement of the cer-vical lesion without development of new lesions.

Discussion. HPV infection is transmitted mainly through sexual contact and it is a necessary step in the pathogenesis of cervical cancers. Cervical cancer is the second most common cancer in women and causes more than 270,000 deaths per year, mostly in low-and middle-income countries.¹ In the United States, there are an estimated 11,000 newly diagnosed cases of HPV-associated cervical cancer and 4,000 resulting deaths annually.³ The World Health Organization recommends vaccination for girls aged 9–13 years as “the most cost-effective public health measure against cervical cancer.”¹ However, there are safety concerns regarding HPV vaccines.

The relationship between HPV vaccination and subsequent CNS inflammation remains unclear. A recent review summarizes 9 published cases of CNS demyelination following HPV vaccination; clinical syn-dromes vary and include myelitis, optic neuritis, and encephalitis.⁴ In the genesis of CNS inflammatory dis-orders post-HPV vaccination, both molecular mimicry between vaccine antigen and myelin proteins and toxic materials in vaccine components are seen as potential causative factors.⁵ There are previous reports of CNS inflammatory syndromes following HPV vaccination describing a 10-day to 5-month time lapse from vacci-nation to symptom onset, with a minimum of a 21-day interval in cases developing myelitis. Recent reviews on this topic did not find an increased rate of subsequent autoimmune diseases; nevertheless, they acknowledge limitations regarding statistical power⁶ and data con-cerning specific populations.⁷

We present a case of transverse myelitis 3 days after HPV immunization. To our knowledge, this is the ear-liest case of CNS inflammation following HPV immuni-zation, with a less than 7-day interval from vaccination to symptom onset. Regarding etiology, the patient was immunocompetent and there were no systemic signs of infection. The absence of cells in CSF analysis and the presence of oligoclonal bands with an increased non-specific immunoglobulin M ratio suggested a back-ground of a CNS inflammatory condition; however, the mechanism of disease, whether viral-induced or immune-mediated, remains to be determined.

Brain MRI T2-weighted images: axial (A) and coronal (B) images showing absence of brain lesions. Cervical MRI T2-weighted images: axial (C) and sagittal (D) images showing an isolated C1-C2 hyperintense lesion.

In conclusion, the probable benefits of mass vaccination with HPV seem to outweigh the potential risks. However, we must continue to report and investigate cases of inflammatory disorders of the CNS associated with qHPV vaccination, both for safety reasons and because it may help comprehend the pathogenesis of subsequent CNS demyelination.

From the Multiple Sclerosis and Neuroimmunology Unit, University Hospital La Paz, Madrid, Spain.

Author contributions: Dr. Fernández-Fournier: conceptualization and manuscript drafting. Dr. Díaz de Terán: manuscript drafting. Dr. Tallón Barranco: manuscript revision. Dr. Puertas: manuscript revision.

Acknowledgement: The authors thank the Neurology Department at La Paz University Hospital, especially Dr. Olga Trabajos, MD, who provided careful patient care and assistance during the patient's hospitalization.

Study funding: No targeted funding reported.

Disclosure: The authors report no disclosures. Go to Neurology.org/nn for full disclosures. The Article Processing Charge was paid by Fundación Investigación Biomédica del Hospital La Paz.

This is an open access article distributed under the terms of the Creative Commons Attribution-Noncommercial No Derivative 3.0 License, which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received June 20, 2014. Accepted in final form August 6, 2014.

Correspondence to Dr. Fernández-Fournier: fernandezfournier@hotmail.com

1. WHO. Human papillomavirus (HPV) and cervical cancer. World Health Organization. Available at: <http://www.who.int/mediacentre/factsheets/fs380/en/>. Accessed July 21, 2014.
2. Muñoz N, Bosch FX, de Sanjosé S, et al. Epidemiologic classification of human papillomavirus types associated with cervical cancer. *N Engl J Med* 2003;348:518–527.
3. Balog JE. The moral justification for a compulsory human papillomavirus vaccination program. *Am J Public Health* 2009;99:616–622.
4. Karussis D, Petrou P. The spectrum of post-vaccination inflammatory CNS demyelinating syndromes. *Autoimmun Rev* 2014;13:215–224.
5. Sutton I, Lahoria R, Tan I, Clouston P, Barnett M. CNS demyelination and quadrivalent HPV vaccination. *Mult Scler* 2009;15:116–119.
6. Grimaldi-Bensouda L, Guillemot D, Godeau B, et al. Auto-immune disorders and quadrivalent human papillomavirus vaccination of young female subjects. *J Intern Med* 2014; 275:398–408.
7. Macartney KK, Chiu C, Georgousakis M, Brotherton JML. Safety of human papillomavirus vaccines: a review. *Drug Saf* 2013;36:393–412.

Neurology[®] Neuroimmunology & Neuroinflammation

Early cervical myelitis after human papilloma virus vaccination

Mireya Fernández-Fournier, Javier Díaz de Terán, Antonio Tallón Barranco, et al.

Neurol Neuroimmunol Neuroinflamm 2014;1;

DOI 10.1212/NXI.0000000000000031

This information is current as of September 11, 2014

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2014 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/1/3/e31.full.html
References	This article cites 6 articles, 0 of which you can access for free at: http://nn.neurology.org/content/1/3/e31.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Adolescence http://nn.neurology.org/cgi/collection/adolescence All Demyelinating disease (CNS) http://nn.neurology.org/cgi/collection/all_demyelinating_disease_cns All Spinal Cord http://nn.neurology.org/cgi/collection/all_spinal_cord Patient safety http://nn.neurology.org/cgi/collection/patient__safety Post-infectious http://nn.neurology.org/cgi/collection/postinfectious_
Errata	An erratum has been published regarding this article. Please see next page or: /content/2/1/e64.full.pdf
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2014 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.

CORRECTION

Early cervical myelitis after human papilloma virus vaccination

In the article “Early cervical myelitis after human papilloma virus vaccination” by M. Fernández-Fournier et al. (*Neurology® Neuroimmunology & Neuroinflammation* 2014;1:e31–e32), there is an error in the name of the vaccine used, which was in fact the bivalent HPV vaccine with a combined diphtheria/tetanus/pertussis vaccine rather than the quadrivalent HPV vaccine Gardasil, as reported in the article. Accordingly, the authors wish to reword several sentences in the article. Under “Case report,” the third sentence should read: “Three days earlier she had received a first dose of bivalent HPV recombinant vaccine together with a combined diphtheria/tetanus/pertussis (DTP) vaccine.” In the third paragraph of the Discussion, the first 2 sentences should read: “We present a case of transverse myelitis 3 days after HPV and DTP immunization. To our knowledge, this is the earliest case of CNS inflammation following either HPV or DTP immunization, with a less than 7-day interval from vaccination to symptom onset.” Sentence 4 should read: “DTP vaccination might have triggered demyelination; however, CNS demyelination is more frequently associated with HPV vaccination in the medical literature.”⁴ The last sentence of the article should read: “However, we must continue to report and investigate cases of inflammatory disorders of the CNS associated with vaccination, both for safety reasons and because it may help comprehend the pathogenesis of subsequent CNS demyelination.” The authors regret the errors.