Kathrin Doppler, MD
Barbara Schleyer
Christian Geis, MD
Benedikt Grünewald,

PhD
Evelyn Putz, MD
Carmen Villmann, PhD
Claudia Sommer, MD

*Neurol Neuroimmunol
Neuroinflamm*
2016;3:e186; doi: 10.1212/
NXL.0000000000000186

LOCKJAW IN STIFF-PERSON SYNDROME WITH AUTOANTIBODIES AGAINST GLYCINE RECEPTORS

OPEN

A 37-year-old man was admitted to our hospital in 1998 with severe contraction of both masseter muscles, which made him unable to open his mouth. He could only drink using a straw fitting through a tooth gap. Spasms of the facial and masseter muscles had started in 1993 and transiently improved after treatment with benzodiazepines. The patient's history included treatment of back and joint pain in 1990 and several admissions to psychiatry for "conversion neurosis." At one time, his wife was suspected of poisoning him with an insecticide because of residues of parathion in his serum.

Neurologic workup including brain and spinal cord MRI, lumbar puncture, extensive blood tests, and nerve conduction studies were normal. EMG showed profuse activity of the masseter muscle that increased when the mouth was passively opened. The masseter inhibitory reflex (MIR), performed according to Aramideh and Ongerboer de Visser,¹ revealed lack of the late silent period (SP2) even at high stimulus intensities, while the early silent period (SP1) was present (figure 1A).

With a tentative diagnosis of autoantibody-mediated stiff-person-like disorder, plasmapheresis was initiated. After the third treatment, the patient was able to open his mouth and to eat normally. EMG showed normal muscle activity. Both SP1 and SP2 of the MIR were now present (figure 1B).

However, after 2 months, lockjaw reappeared. Again, plasmapheresis led to complete recovery. For 1 year, plasmapheresis was repeated every 6 weeks, because this was the interval when lockjaw recurred. Corticosteroids (methylprednisolone 1,000 mg/d for up to 5 days), azathioprine (150 mg/d), and IV immunoglobulins (Igs) (5×30 g, equaling 2 g/kg body weight) were ineffective and unable to prolong the intervals. In 1999, spasms of the right body and sudden falls occurred. Again, plasmapheresis led to fast improvement. Consecutive treatment with mycophenolate mofetil, cyclophosphamide, and rituximab was not effective. Immunoadsorption using a protein A-coated column was regularly performed since then,

and a Cimino fistula was installed in the patient's left forearm for reliable vascular access. Still, lockjaw was the main symptom, remitting rapidly after immunoadsorption and recurring a few weeks later (video at Neurology.org/nn). If an interval of 4 weeks for immunoadsorption was maintained, lockjaw could be prevented. Whenever the interval was prolonged—like most recently because of a strike of train drivers in May 2015, which made it impossible for the patient to attend our department for immunoadsorption—lockjaw and falls reoccurred. Limb spasms and dystonic malposition of the right fingers 3 to 5 are present most of the time, but myoclonus, other brainstem symptoms, and hyperekplexia have not occurred so far. Assays for autoantibodies against anti-glutamate decarboxylase or anti-amphiphysin were negative, as was tumor screening. Patient serum did not bind to sections of human, rat, or mouse nervous tissue but did bind to freshly dissociated mouse hippocampal neurons.

In 2014, anti-glycine receptor (GlyR) α autoantibodies were detected in the patient's serum and purified IgG by binding assays using unfixed human embryonic kidney 293 cells transfected with GlyR $\alpha 1$, $\alpha 2$, and $\alpha 3$ (figure 2). Binding to $\alpha 1$, $\alpha 2$, and $\alpha 3$ subunits suggests that GlyR autoantibodies recognize an epitope common to all α subunits. Similar reactivity to all α subunits that, however, did not relate to the overall autoantibody titers has been described in a recent study.² In fact, the use of unfixed cells has been reported to be more sensitive compared to fixed cells in the detection of GlyR $\alpha 1$ autoantibodies.³ GlyR $\alpha 1$ autoantibodies have been described in patients with progressive encephalomyelitis with rigidity and myoclonus (PERM),^{4–6} a more severe variant of stiff-person syndrome, and also in 12% of patients with stiff-person syndrome.⁷ In a larger cohort of GlyR α antibody-positive patients, the majority showed the typical clinical picture of PERM; only 2 patients had stiff-person syndrome.² Trigeminal, facial, and bulbar symptoms, including trismus, were reported at the onset of disease in 47% and in 57% during the course of disease.² Trismus was also predominant in an adult in this study, resulting in an inability to eat.⁶ However, all patients described so far presented with additional signs and symptoms,

**Supplemental data
at Neurology.org/nn**

Figure 1 Masseter inhibitory reflex

The masseter inhibitory reflex revealed lack of the late silent period (SP2) even at high stimulus intensities, while the early silent period (SP1) was present (A). After treatment, both silent periods (SP1 and SP2) were present (B).

facilitating diagnosis of stiff-person syndrome or PERM. Several studies aimed to determine the clinical syndrome of anti-GlyR antibody-related disease,

but concluded that it comprises a wide range of symptoms, mostly summarized as PERM.^{2,6} Brainstem symptoms are often reported, and binding assays with

Figure 2 GlyR antibody binding

Antibodies bind to extracellular domain of glycine receptor subunits. HEK293 cells were cotransfected with GFP and with different glycine receptor subunits 1, 2, 3. Cells were incubated with purified patient IgG (I-L) and IgG from a healthy control (E-H) in a 2 mg/mL concentration. The GlyR 1-specific antibody MAb2B was used as a positive control, as well as the pan-antibody MAb4A to stain 2 and 3 (A-D). Of note, the surface staining of all GlyR subunits upon incubation with the patient's IgG is only present in live cells (A-C, E-G, I-K), not in fixed and permeabilized cells (D, H, L), while the control antibodies bind in both conditions. This shows that the antibodies in patient serum bind only to the glycine receptors in their native state. White bar = 10 μ m. GFP = green fluorescent protein; GlyR = glycine receptor; IgG = immunoglobulin G; MAb = monoclonal antibody.

patients' sera on brainstem showed distinctive binding,² which is in line with the predominant distribution of GlyR in the brainstem.⁸ GlyR mediate inhibitory neurotransmission and dysregulation may induce uncontrolled excitation resulting in muscle spasms.⁸ The lack of the MIR SP2 in contrast to the preserved SP1 underscores a deficit in inhibitory neurotransmission at the level of the pontomedullary junction.¹ Of note, in GAD antibody-positive stiff-person syndrome, the MIR can be elicited with high stimulus intensities, in contrast to the glycine receptor disorder familial hyperekplexia,⁹ which supports the glycine-related pathophysiology in our patient.

We conclude that stiff-person syndrome and especially anti-GlyR-mediated disease needs to be considered in cases of severe lockjaw even if there are no other relevant symptoms of stiff-person syndrome or PERM at the onset of disease as the clinical presentation is heterogeneous. Appropriate treatment needs to be initiated to prevent disease progression. In our case, plasmapheresis was started before the detection of autoantibodies. An unequivocal response to plasmapheresis can be of diagnostic value in suspected autoantibody-mediated disease and can significantly improve patients' quality of life.

From the Department of Neurology (K.D., C.G., B.G., C.S.), Institute for Clinical Neurobiology (B.S., C.V.), and Department of Transfusion Medicine (E.P.), University Hospital Würzburg; and Hans Berger Department of Neurology and Center for Sepsis Control and Care (C.G., B.G.), Jena University Hospital, Jena, Germany.

Author contributions: K.D.: treatment of patient, writing of the manuscript. B.S.: detection of autoantibodies. C.G.: treatment of patient, binding assays, writing of the manuscript. B.G.: binding assays. E.P.: treatment of patient. C.V.: detection of autoantibodies. C.S.: planning of study, treatment of patient, writing of the manuscript.

Study funding: Research by C.S. and C.G. on the stiff-person syndrome was funded by Deutsche Forschungsgemeinschaft, DFG, SFB 581. Research by C.V. on glycine receptors is funded by DFG VI586. The funding source had no influence on study design, writing, or decision to publish.

Disclosure: K. Doppler received speaker honoraria from Baxter Germany. B. Schleyer reports no disclosures. C. Geis received speaker honoraria from Teva, Allergan, was a guest editor for Frontiers in Neurology, received research support from German Research

Council, German Ministry of Education. B. Grünewald and E. Putz report no disclosures. C. Villmann is an associate editor for Frontiers in Molecular Neuroscience, Biological Chemistry. C. Sommer is on the scientific advisory board for Astellas Pharma Inc., Baxter International Inc., Eli Lilly and Company, Pfizer Inc., UCB, received travel funding and/or speaker honoraria from Astellas Pharma Inc., Allergan Baxter International Inc., CSL Behring, Eli Lilly and Company, Pfizer Inc., is an associate editor for Journal of the Peripheral Nervous System, PAIN, is an academic editor for PLOS ONE, received research support from German Research Foundation. Go to Neurology.org/nn for full disclosure forms. The Article Processing Charge was paid by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received August 16, 2015. Accepted in final form September 21, 2015.

Correspondence to Dr. Sommer: sommer@uni-wuerzburg.de

1. Aramideh M, Ongerboer de Visser BW. Brainstem reflexes: electrodiagnostic techniques, physiology, normative data, and clinical applications. *Muscle Nerve* 2002;26:14–30.
2. Carvajal-Gonzalez A, Leite MI, Waters P, et al. Glycine receptor antibodies in PERM and related syndromes: characteristics, clinical features and outcomes. *Brain* 2014;137:2178–2192.
3. Alexopoulos H, Akrivou S, Dalakas MC. Glycine receptor antibodies in stiff-person syndrome and other GAD-positive CNS disorders. *Neurology* 2013;81:1962–1964.
4. Hutchinson M, Waters P, McHugh J, et al. Progressive encephalomyelitis, rigidity, and myoclonus: a novel glycine receptor antibody. *Neurology* 2008;71:1291–1292.
5. Clerinx K, Breban T, Schrooten M, et al. Progressive encephalomyelitis with rigidity and myoclonus: resolution after thymectomy. *Neurology* 2011;76:303–304.
6. Mas N, Saiz A, Leite MI, et al. Antiglycine-receptor encephalomyelitis with rigidity. *J Neurol Neurosurg Psychiatry* 2011;82:1399–1401.
7. McKeon A, Martinez-Hernandez E, Lancaster E, et al. Glycine receptor autoimmune spectrum with stiff-man syndrome phenotype. *JAMA Neurol* 2013;70:44–50.
8. Lynch JW. Molecular structure and function of the glycine receptor chloride channel. *Physiol Rev* 2004;84:1051–1095.
9. Khasani S, Becker K, Meinck HM. Hyperekplexia and stiff-man syndrome: abnormal brainstem reflexes suggest a physiological relationship. *J Neurol Neurosurg Psychiatry* 2004;75:1265–1269.

Neurology® Neuroimmunology & Neuroinflammation

Lockjaw in stiff-person syndrome with autoantibodies against glycine receptors

Kathrin Doppler, Barbara Schleyer, Christian Geis, et al.

Neurol Neuroimmunol Neuroinflamm 2016;3;

DOI 10.1212/NXI.0000000000000186

This information is current as of December 10, 2015

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/3/1/e186.full.html
Supplementary Material	Supplementary material can be found at: http://nn.neurology.org/content/suppl/2015/12/10/3.1.e186.DC1
References	This article cites 9 articles, 2 of which you can access for free at: http://nn.neurology.org/content/3/1/e186.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://nn.neurology.org/cgi/collection/all_clinical_neurology All Immunology http://nn.neurology.org/cgi/collection/all_immunology Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases EMG http://nn.neurology.org/cgi/collection/emg Stiff person syndrome http://nn.neurology.org/cgi/collection/stiff_person_syndrome
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.