Camille Legouy, MD
Anne-Celine Zeghoudi,
MD
Irina Taifas, MD
Elise Le Stanc, MD
Jerome Honnorat, MD,
PhD
Fernando Pico, MD, PhD

*Neurol Neuroimmunol
Neuroinflamm*
2015;2:e168; doi: 10.1212/
NXL.0000000000000168

CEREBELLAR VERMIS HYPERMETABOLISM IN OPSOCLONUS MYOCLONUS WITHOUT ONCONEURAL ANTIBODIES

OPEN

A 41-year-old woman presented with dizziness and tremor of the right hand. She had myoclonus of the upper limbs and opsoclonus.¹ Brain MRI including enhanced gadolinium sequences was normal, in particular the cerebellum (figure 1, A and B). CSF analysis disclosed 24 cell/mm³ with 97% lymphocytes and 0.69 g/L of proteinorachia with oligoclonal bands. A workup for infection and neoplasia was negative except for the presence of atypical neuronal antibodies^{2,3} in the CSF detected by immunohistochemistry in a reference center (J.H.) (figure 2, A and B). All other common onconeural antibodies associated or not associated with OMS were excluded, particularly anti-Ri, anti-Hu, anti-Ma2, and anti-NMDA receptor.³ The PET performed 8 days after brain MRI depicted a hypermetabolism⁴ in the cerebellar vermis, which was no longer observed at 4-month follow-up (compare figure 1, C and D). The patient recovered almost completely by the time of the second PET. This PET pattern is rarely seen. Whether it is associated with opsoclonus myoclonus syndrome or specific immune reaction requires further studies.

From the Neurology Department and Stroke Center (C.L., A.-C.Z., I.T., F.P.), Versailles Mignot Hospital, Versailles; Versailles Saint Quentin en Yvelines University (C.L., A.-C.Z., I.T., F.P.); Nuclear Medicine Department (E.L.S.), Foch Hospital, Suresnes; French Reference Center on Paraneoplastic Neurological Syndrome (J.H.), Department of Neuro-oncology, Hospices Civils de Lyon, Hôpital Neurologique, Bron; Lyon Neuroscience Research Center INSERM U1028/CNRS UMR 5292 (J.H.), Lyon; and Université de Lyon—Université Claude Bernard Lyon 1 (J.H.), France.

Author contributions: Dr. Legouy: study concept and design. Dr. Taifas: acquisition of data. Dr. Zeghoudi: analysis and interpretation. Dr. Le Stanc and Prof. Honnorat: critical revision of the manuscript for important intellectual content. Prof. Pico: study supervision.

Study funding: No targeted funding.

Disclosure: The authors report no disclosures. Go to Neurology.org/nn for full disclosure forms. The Article Processing Charge was paid by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and

Figure 1 Brain MRI

Brain MRI performed 8 days before PET (A and B). Axial fluid-attenuated inversion recovery (A) and T1 enhanced gadolinium (B) depicted no abnormalities in the cerebellar vermis. Hypermetabolism in the cerebellar vermis (C, white arrow), which disappeared at 4-month follow-up examination (D).

sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received December 22, 2014. Accepted in final form September 14, 2015.

Correspondence to Dr. Legouy: camille.legouy@gmail.com

Figure 2 Detection of atypical anti-neuronal antibodies

Immunohistofluorescence assay on rat brain sagittal slices incubated with the patient's CSF and revealed by fluorescent anti-human immunoglobulin G. In the striatum, a staining of some neuron nucleus is observed (A, $\times 200$). In the cerebellum, the patient's CSF stained nucleus and weakly the cytoplasm of Purkinje cells (B, $\times 400$).

1. Klaas PJ, Ahlskog E, Pittock SJ, et al. Adult-onset opsoclonus-myoclonus syndrome. *Arch Neurol* 2012;69:1598–1607.
2. Baumgartner A, Rauer S, Mader I, Meyer PT. Cerebral FDG-PET and MRI findings in autoimmune limbic encephalitis: correlation with autoantibody types. *J Neurol* 2013;260:2744–2753.
3. Spatola S, Stojanova V, Prior JO, Dalmau J, Rossetti AO. Serial brain 18FDG-PET in anti-AMPA receptor limbic encephalitis. *J Neuroimmunol* 2014;271:53–55.
4. Ances BM, Vitaliani R, Taylor RA, et al. Treatment-responsive limbic encephalitis identified by neuropil antibodies: MRI and PET correlates. *Brain* 2005;128:1764–1777.

Neurology[®] Neuroimmunology & Neuroinflammation

Cerebellar vermis hypermetabolism in opsoclonus myoclonus without onconeural antibodies

Camille Legouy, Anne-Celine Zeghoudi, Irina Taifas, et al.
Neurol Neuroimmunol Neuroinflamm 2015;2;
DOI 10.1212/NXI.0000000000000168

This information is current as of October 29, 2015

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/2/6/e168.full.html
References	This article cites 4 articles, 0 of which you can access for free at: http://nn.neurology.org/content/2/6/e168.full.html#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases Clinical neurology history http://nn.neurology.org/cgi/collection/clinical_neurology_history Myoclonus http://nn.neurology.org/cgi/collection/myoclonus Ocular motility http://nn.neurology.org/cgi/collection/ocular_motility PET http://nn.neurology.org/cgi/collection/pet
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.