Cliff W. Hampton, MD
Dina Galperin, MD
Enrique Alvarez, MD
Matthew S. West, MD

*Neurol Neuroimmunol
Neuroinflamm*
2015;2:e82; doi: 10.1212/
NXL.0000000000000082

ENCEPHALOMYELONEUROPATHY WITH CRMP-5 ANTIBODIES IN A PATIENT WITH A PRIMARY MEDIASTINAL SERINOMA

OPEN

Case report. A 23-year-old nonsmoking man presented with difficulty walking, numbness, and weakness progressing over several months. By the time of presentation, the patient was not ambulatory. He was afebrile with normal vital signs and general examination. Neurologic examination demonstrated normal vision, mildly decreased strength, and severely diminished sensation to vibration and proprioception in both lower extremities. His temperature and pinprick sensation were mildly diminished in both legs and arms. Reflexes were diffusely diminished but not absent. He exhibited significant dysmetria in both upper and lower extremities. No tremor or other abnormal movements were noted.

MRI of the spinal cord demonstrated increased T2 signal extending from the craniocervical junction to the conus medullaris with gadolinium enhancement (figure, A–D). MRI of the brain demonstrated a heterogeneous region of contrast enhancement in the cerebellum (figure, E–F). CSF analysis showed a nucleated cell count of 187 (87% lymphocytes), no red blood cells, glucose 75 mg/dL, and protein elevated to 165 mg/dL. Oligoclonal bands were elevated at 2 (normal 0–1), albumin index was 24.2 (normal 0–9), and IgG index was 0.71 (normal 0.28–0.66). (Additional laboratory values are presented in the supplemental appendix at Neurology.org/nn.) Approximately 2 months prior to presentation, the patient underwent a nerve conduction study at another facility, which demonstrated a demyelinating polyneuropathy. Extensive rheumatologic and infectious workup was negative. Aquaporin-4 antibody was negative in serum and CSF. Serum paraneoplastic studies demonstrated a positive IgG antibody (titer 1:30,720) to collapsing response-mediator protein 5 (CRMP-5). While awaiting these results, the patient received steroids and plasma exchange. CT of the chest revealed a 3.4-cm anterior mediastinal soft tissue mass, with biopsy and pathology results consistent with seminoma (figure, G–H). There was no evidence of testicular neoplasm on ultrasound.

The patient demonstrated some improvement in strength and sensation after steroids and plasmapheresis.

He received 4 cycles of etoposide and cisplatin and continued to improve with treatment of the primary seminoma. Repeat imaging demonstrated resolution of the inflammation seen previously in the cerebellum and spinal cord, and repeat serum studies were negative for the CRMP-5 antibody. Follow-up PET did not reveal any fludeoxyglucose uptake in the mediastinum. Nerve conduction studies performed approximately 6 months later revealed a sensory ganglionopathy with absent sensory responses. He continues to have some difficulty with ambulation related primarily to severe proprioceptive deficits.

Discussion. Antibodies to CRMP-5 were first described clinically in a series of patients presenting with a wide variety of neurologic symptoms, including (from most to least common) peripheral neuropathy, autonomic neuropathy, cerebellar ataxia, subacute dementia, cranial neuropathy, and neuromuscular junction dysfunction.¹ Subsequent reports have shown an association of CRMP-5 with optic neuritis and retinitis,² chorea,³ and a neuromyelitis optica–like syndrome of transverse myelitis with optic neuritis.⁴ The patient described in this report had a combination of cerebellar ataxia, longitudinally extensive transverse myelitis involving primarily the dorsal columns, and demyelinating polyneuropathy.

In the reported cases of anti-CRMP-5 paraneoplastic disorders, the primary tumor was most commonly small cell lung cancer (SCLC). However, association with renal cell carcinoma, thymoma, thyroid papillary carcinoma, lymphoma, and prostate adenocarcinoma has also been described. We report the association of anti-CRMP-5 with a seminoma.

Seminomas have been associated rarely with anti-Ma2 antibodies affecting the limbic system, diencephalon, or brainstem.⁵ Primary mediastinal germ cell tumors have been recently reported as being associated with paraneoplastic disorders in 2 cases.^{6,7} The first presented with encephalitis associated with anti-Ma2, whereas the second had encephalitis, sensorimotor polyneuropathy, vasculomyositis, and cerebellar ataxia associated with anti-NMDA and anti-neuronal nuclear autoantibody type 1.

In this case, the patient responded clinically to immunosuppression followed by treatment of the underlying cancer with cisplatin and etoposide, although

Supplemental data
at Neurology.org/nn

Figure MRI demonstrating cerebellar encephalitis, longitudinally extensive transverse myelitis, and pathology of seminoma

(A) Parasagittal T1 postcontrast images of the cervical spine demonstrate homogeneous enhancement of the spinal cord from the craniocervical junction that extends through the mid-thoracic spine and also several ill-defined mass-like areas of enhancement in both cerebellar hemispheres involving both cortex and white matter. (B) Parasagittal T2 image demonstrating increased signal in these same areas. (C) Axial T1 postcontrast image demonstrating that the enhancement is primarily dorsal. (D) Axial T2 image with increased signal in the dorsal cord. (E, F) Axial T1-weighted postcontrast images demonstrating multifocal areas of patchy enhancement in the cerebellum. (G) Mediastinal soft tissue mass demonstrates positive immunohistochemistry for placental alkaline phosphatase and (H) CD 117. These are characteristic of seminoma.

his recovery was complicated by the chemotherapeutic agents' possible role in neuropathy/ganglionopathy. In cases of subacute onset syndromes of neurologic dysfunction involving multiple components of the nervous system in which initial evaluation for infection and primary autoimmunity does not reveal a cause, paraneoplastic studies should be considered, even in nonsmokers. Chest CT should be considered as first-tier testing for primary neoplasm, as many paraneoplastic syndromes are associated with SCLC.^{1,2,4,5} However, as paraneoplastic syndromes including CRMP-5 become increasingly recognized, it is important to expand the differential diagnosis of tumors associated with paraneoplastic syndromes to include seminomas and other primary neoplasms.

From the Department of Neurology, University of Colorado Denver, Aurora, CO.

Author contributions: Drs. Hampton, Galperin, Alvarez, and West participated in drafting and revising of the article for intellectual content.

Study funding: No targeted funding reported.

Disclosure: C.W. Hampton serves on the editorial board for the Resident and Fellow section of Neurology. D. Galperin reports no disclosures. E. Alvarez has received consulting honoraria from Teva Neuroscience, Biogen Idec, and Genzyme. M.S. West reports no disclosures. Go to Neurology.org/nm for full disclosure forms. The Article Processing Charge was paid by the University of Colorado Health Science Library Fund.

This is an open access article distributed under the terms of the Creative Commons Attribution-Noncommercial No Derivative 3.0 License, which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received November 17, 2014. Accepted in final form January 15, 2015.

Correspondence to Dr. West: matthew.west@ucdenver.edu

1. Yu Z, Kryzer TJ, Griesmann GE, et al. CRMP-5 neuronal autoantibody: marker of lung cancer and thymoma-related autoimmunity. *Ann Neurol* 2001;49:146–154.
2. Cross SA, Salomao DR, Parisi JE, et al. Paraneoplastic autoimmune optic neuritis with retinitis defined by CRMP-5-IgG. *Ann Neurol* 2003;54:38–50.
3. Vernino S, Tuite P, Adler CH, et al. Paraneoplastic chorea associated with CRMP-5 neuronal antibody and lung carcinoma. *Ann Neurol* 2002;51:625–630.
4. Jarius S, Wandinger KP, Borowski K, Stoecker W, Wildemann B. Antibodies to CV2/CRMP5 in neuromyelitis-optica-like disease: case report and review of the literature. *Clin Neurol Neurosurg* 2012;114:331–335.
5. Dalmau J, Graus F, Villarejo A, et al. Clinical analysis of anti-Ma2-associated encephalitis. *Brain* 2004;127:1831–1844.
6. Bosemani T, Huisman TA, Poretti A. Anti-Ma2-associated paraneoplastic encephalitis in a male adolescent with mediastinal seminoma. *Pediatr Neurol* 2014;50:433–444.
7. Pohley I, Roesler K, Wittstock M, et al. NMDA-receptor antibody and anti-Hu antibody positive paraneoplastic syndrome associated with a primary mediastinal seminoma. *Acta Neurol Belg* 2014;1–3.

Neurology® Neuroimmunology & Neuroinflammation

Encephalomyeloneuropathy with CRMP-5 antibodies in a patient with a primary mediastinal serinoma

Cliff W. Hampton, Dina Galperin, Enrique Alvarez, et al.
Neurol Neuroimmunol Neuroinflamm 2015;2;
DOI 10.1212/NXI.0000000000000082

This information is current as of March 12, 2015

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/2/3/e82.full.html
Supplementary Material	Supplementary material can be found at: http://nn.neurology.org/content/suppl/2015/03/12/2.3.e82.DC1
References	This article cites 6 articles, 0 of which you can access for free at: http://nn.neurology.org/content/2/3/e82.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Immunology http://nn.neurology.org/cgi/collection/all_immunology Paraneoplastic syndrome http://nn.neurology.org/cgi/collection/paraneoplastic_syndrome Transverse myelitis http://nn.neurology.org/cgi/collection/transverse_myelitis
Errata	An erratum has been published regarding this article. Please see next page or: /content/2/4/e133.full.pdf
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.

CORRECTION

Encephalomyeloneuropathy with CRMP-5 antibodies in a patient with a primary mediastinal serinoma

In the Clinical/Scientific Note “Encephalomyeloneuropathy with CRMP-5 antibodies in a patient with a primary mediastinal serinoma” by C.W. Hampton et al. (*Neurology® Neuroimmunology & Neuroinflammation* 2015;2:e82), “serinoma” was misspelled in the title. The publisher regrets the error.