Kathrin Schanda, MSc
Patrick Waters, PhD
Hannah Holzer, BSc
Fahmy Aboulenein-

Djamshidian, MD
M. Isabel Leite, DPhil
Jacqueline Palace, DM
Sandra Vukusic, MD
Romain Marignier, MD
Thomas Berger, MD
Markus Reindl, PhD

*Neurol Neuroimmunol
Neuroinflamm*
2015;2:e160; doi: 10.1212/
NXL.0000000000000160

ANTIBODIES TO AQUAPORIN-1 ARE NOT PRESENT IN NEUROMYELITIS OPTICA

OPEN

Although more than 70% of all patients with neuromyelitis optica (NMO) are seropositive for aquaporin-4 (AQP4) antibodies, a substantial proportion of patients fulfilling the clinical criteria for NMO or limited forms of the disease (NMO spectrum disorders [NMOSD]) are negative for these antibodies.¹ This raises the possibility of other autoantibodies in these patients. One putative target, the ubiquitously expressed water channel aquaporin-1 (AQP1), which is partially lost or internalized in certain NMO lesions,² was described in a subset of patients with NMOSD.^{3–5} However, AQP1 antibodies were also found in patients with multiple sclerosis (MS), thus raising concerns about the specificity of these findings. Therefore, we developed a recombinant live cell immunofluorescence assay (CBA) for AQP1 antibodies based on our AQP4 antibody assay.⁶ We analyzed 176 serum samples from Austrian patients with NMOSD (n = 67), Austrian patients with MS (n = 31), and controls (n = 78) for the presence of IgG AQP1 antibodies and AQP4 antibodies. Furthermore, both assays were validated in a blinded cohort of 58 patients with NMOSD (n = 36) or MS (n = 22) from Lyon (France) and Oxford (United Kingdom). Clinical and demographic data of patients and controls are shown in the table.

Methods. Analysis of AQP1 antibodies and AQP4 antibodies was performed using a live CBA described previously.⁶ Briefly, HEK293A cells were transiently transfected using the pcDNA6.2C-EmGFP-GW/TOPO plasmid (Invitrogen, Carlsbad, CA), expressing AQP4 (isoform M23) or AQP1 (isoform 1) fused C-terminally to emerald green fluorescence protein. Transfected cells were blocked with goat IgG in phosphate-buffered saline (PBS)/10% fetal calf serum (FCS) (Sigma-Aldrich, St. Louis, MO) followed by serum diluted 1:20 and 1:40 in PBS/FCS for 1 hour at 4°C. Serum preabsorption with liver powder was not performed because a previous report indicated loss of AQP1 antibodies after pretreatment.³ Bound antibodies were detected using Cy3[™]-conjugated goat anti-human IgG antibody (Jackson ImmunoResearch Laboratories,

West Grove, PA) for 30 minutes at room temperature. Bound antibodies were determined using a fluorescence microscope (Leica DMI 4000B). All samples were evaluated by 2 independent, clinically blinded investigators who agreed on all samples.

Standard protocol approvals, registrations, and patient consents. The present study was approved by the ethical committee of Medical University of Innsbruck (#AM3041a), the Oxfordshire Research Ethics Committee (#10/H0606/56), and the French data protection authority.

Results. Expression of AQP1 was verified by staining with a commercial AQP1-specific antibody recognizing full-length AQP1. However, the antibody only recognized the target after fixation of AQP1-transfected HEK293A cells with paraformaldehyde (figure e-1A at Neurology.org/nn). In order to prove surface expression and correct topology of AQP1, a myc-tag was inserted at position T120 (extracellular loop C) by site-directed mutagenesis. Staining with an anti-myc-tag monoclonal antibody clearly showed the surface expression of AQP1 in live HEK293A cells (figure e-1B).

As seen in the table, AQP4 antibodies were detected in 81 of 103 (79%) patients with NMOSD in the combined cohorts. AQP4 antibodies were absent in 53 patients with MS and 78 controls; therefore, the specificity of the AQP4 antibody assay was 100%. In contrast, AQP1 antibodies were absent in all 234 samples from patients with NMOSD and MS and controls. Although the AQP4 antibody CBA showed high sensitivity and specificity, a comparable AQP1 antibody CBA did not detect any antibodies in 234 serum samples (figure e-1C).

Discussion. This finding is in contrast to previous studies by 2 groups reporting the presence of AQP1 antibodies in 17%–74% of patients with NMOSD,^{3–5} but also in controls. In our opinion, these differences could be explained by methodologic differences between the studies (using radioimmunoprecipitation, ELISA with peptides, or fixed CBAs might have exposed intracellular epitopes, whereas our assay only detects antibodies to extracellular epitopes). Methodologic differences have proven to be a constant problem in this field for decades. For example, there is now clear evidence that antibodies to the myelin oligodendrocyte glycoprotein are specific

Supplemental data
at Neurology.org/nn

Table AQP4-IgG and AQP1-IgG antibodies in patients with NMO spectrum disorders and controls

	NMOSD				p Value
	NMO	LETM/ON	MS	CTRL	
N	45	58	53	78	
Females, n (%)	40 (89)	40 (69)	32 (63)	61 (78)	0.017 ^a
Age, y, median (range)	49 (18–83)	51 (13–78)	40 (19–66)	44 (18–84)	0.007 ^b
AQP4-IgG, n (%)	40 (89)	41 (71)	0 (0)	0 (0)	<0.001 ^a
AQP4-IgG titer, median (range)	640 (0–20,480)	320 (0–10,240)	0	0	<0.001 ^b
AQP1-IgG, n (%)	0 (0)	0 (0)	0 (0)	0 (0)	1.000 ^a
AQP1-IgG titer, median (range)	0	0	0	0	1.000 ^b

Abbreviations: AQP1 = aquaporin-1; AQP4 = aquaporin-4; CTRL = controls (23 patients with other neurologic diseases, 25 patients with systemic lupus erythematosus, and 30 healthy controls); LETM/ON = patients with longitudinally extensive transverse myelitis (n = 45) or optic neuritis (n = 13); MS = multiple sclerosis according to the 2005 revisions to the McDonald criteria (n = 53); NMO = clinically definite neuromyelitis optica according to the Wingerchuk 2006 criteria; NMOSD = NMO spectrum disorders (NMO + LETM/ON).

^a Groups were compared using the χ^2 test.

^b Groups were compared using the Kruskal-Wallis test.

for a subset of demyelinating diseases only if appropriate CBAs are used (these antibodies are detected at similar frequencies in patients and controls using ELISA).¹ Similarly, although numerous studies confirm the importance of AQP4 antibodies as diagnostic biomarkers for NMOSD, the seropositivity rates are influenced by the assays used, and some methods, such as ELISA or immunoblotting, also detect these antibodies in controls.¹ A possible limitation of our AQP1 assay is the use of HE-K293A cells, because astrocytic AQP1 might be expressed in a complex against which the immune system could react.

However, the absence of AQP1 antibodies in NMOSD and controls fits very well with AQP1's role in hematology. AQP1, also known as channel-forming integral protein, is well-known in transfusion medicine because it contains the Colton blood group antigen expressed on erythrocytes. Anti-Colton antibodies are very rare and lead to significant delayed or acute transfusion reactions or hemolytic disease,⁷ a clinical phenotype absent in NMOSD. To conclude, our study failed to confirm the presence of AQP1 antibodies in NMOSD.

From the Clinical Department of Neurology (K.S., H.H., T.B., M.R.), Medical University of Innsbruck, Austria; Nuffield Department of Clinical Neurosciences (P.W., M.I.L.), University of Oxford, United Kingdom; Department of Neurology (F.A.-D.), Karl Landsteiner Institute for Neuroimmunological and Neurodegenerative Disorders, Sozialmedizinisches Zentrum Ost Donauespital, Vienna, Austria; Department of Neurology (J.P.), Oxford University Hospitals Trust, Oxford, United Kingdom; and Department of Neurology (S.V., R.M.), Lyon University Hospital and Lyon Neurosciences Research Centre, Lyon, France.

Author contributions: K.S., P.W., and M.R. conceived and designed the experiments. K.S., H.H., and M.R. performed the experiments. K.S., P.W., T.B. and M.R. analyzed the data. P.W., F.A.-D., M.I.L., J.P., S.V., R.M., and M.R. contributed reagents/materials/analyses. K.S., P.W., and M.R. wrote the manuscript.

Acknowledgment: The authors are grateful to the Austrian NMO study group for contributing to this work.

Study funding: This study was funded by the Eugene Devic European Network (EDEN) project (ERA-Net ERARE 2: <http://www.erare.eu/financed-projects/eden>; Austrian Science Fund FWF project I916). P.W., M.I.L., and J.P. are funded by the NHS National Specialised Services for Neuromyelitis optica and the NIHR Oxford Biomedical Research Centre.

Disclosure: K. Schanda reports no disclosures. P. Waters received speaker honoraria from Biogen Idec Japan and Euroimmun AG, is a review editor for Frontiers in Molecular Innate Immunity, and holds a patent for and receives royalties from assays for the detection of antibodies to Igi1, Caspr2, Tag-1, and GABAAR. H. Holzer and F. Aboulenein-Djamschidian report no disclosures. M.I. Leite received travel grants and/or speaker honoraria from Biogen Idec and Novartis, is supported by NHS National Specialised Commissioning Group for Neuromyelitis Optica and NIHR Oxford Biomedical Research Centre, and is on the editorial board for Neuromuscular Disorders. J. Palace is on the scientific advisory boards for Merck Serono, Bayer Schering, Biogen Idec, Teva, Novartis, Sanofi-Aventis, and Alexion; received travel funding and/or speaker honoraria from Merck Serono, Novartis, Biogen Idec, Bayer Schering, and Medimmune; has consulted for Ono Pharmaceuticals, Chugai, CI Consulting, Biogen Idec, GlaxoSmithKline, and Alexion; is on the speakers' bureau for Teva and CMSC; received research support from Bayer Schering, Merck Serono, Novartis, Teva, Department of Health Risk Sharing Scheme, Department of Health, MS Society, Guthy Jackson Foundation, and ISIS Innovation Limited, a wholly owned subsidiary of the University of Oxford; and has filed a patent application to protect for the use of metabolomics as a method to diagnose and stage disease in multiple sclerosis. S. Vukusic is on the scientific advisory board for Biogen Idec, Genzyme, Merck Serono, Novartis Pharma, Sanofi-Aventis, and Teva; received travel funding and/or speaker honoraria from Biogen Idec, Genzyme, Merck Serono, Novartis, Sanofi, and Teva; has consulted for Merck Serono, Sanofi-Aventis, Teva, Biogen Idec, Genzyme, and Novartis; and received research support from Bayer Schering, Biogen Idec, Genzyme, Merck Serono, Novartis, Sanofi-Aventis, and Teva. R. Marignier served on the scientific advisory board for Medimmune. T. Berger reports no disclosures. M. Reindl is an academic editor for PLoS ONE; is on the editorial board for Current Medicinal Chemistry and Autoimmune Diseases; received research support from Eugene Devic European Network and Austrian Science Fund FWF; and The University Hospital and Medical University of Innsbruck (Austria, Markus Reindl) receive payments for antibody assays (NMDAR, AQP4, and other autoantibodies) and for AQP4

antibody validation experiments organized by Euroimmun. Go to Neurology.org/nn for full disclosure forms. The Article Processing Charge was paid by the Austrian Science Funds (project I916, Erare EDEN).

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received June 12, 2015. Accepted in final form August 7, 2015.

Correspondence to Dr. Reindl: markus.reindl@i-med.ac.at

1. Berger T, Reindl M. Antibody biomarkers in CNS demyelinating diseases—a long and winding road. *Eur J Neurol* 2015;22:1162–1168.
2. Misu T, Hoftberger R, Fujihara K, et al. Presence of six different lesion types suggests diverse mechanisms of tissue injury in neuromyelitis optica. *Acta Neuropathol* 2013;125:815–827.
3. Tzartos JS, Stergiou C, Kilidireas K, Zisimopoulou P, Thomaidis T, Tzartos SJ. Anti-aquaporin-1 autoantibodies in patients with neuromyelitis optica spectrum disorders. *PLoS One* 2013;8:e74773.
4. Long Y, Zheng Y, Shan F, et al. Development of a cell-based assay for the detection of anti-aquaporin 1 antibodies in neuromyelitis optica spectrum disorders. *J Neuroimmunol* 2014;273:103–110.
5. Tüzün E, Tzartos J, Ekizoğlu E, et al. Aquaporin-1 antibody in neuromyelitis optical patients. *Eur Neurol* 2014;72:271–272.
6. Mader S, Lutterotti A, Di Pauli F, et al. Patterns of antibody binding to aquaporin-4 isoforms in neuromyelitis optica. *PLoS One* 2010;5:e10455.
7. Halverson GR, Peyrard T. A review of the Colton blood group system. *Immunohematology* 2010;26:22–26.

Neurology[®] Neuroimmunology & Neuroinflammation

Antibodies to aquaporin-1 are not present in neuromyelitis optica

Kathrin Schanda, Patrick Waters, Hannah Holzer, et al.

Neurol Neuroimmunol Neuroinflamm 2015;2;

DOI 10.1212/NXI.0000000000000160

This information is current as of October 1, 2015

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/2/6/e160.full.html
Supplementary Material	Supplementary material can be found at: http://nn.neurology.org/content/suppl/2015/10/01/2.6.e160.DC1
References	This article cites 7 articles, 0 of which you can access for free at: http://nn.neurology.org/content/2/6/e160.full.html##ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://nn.neurology.org/content/2/6/e160.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Demyelinating disease (CNS) http://nn.neurology.org/cgi/collection/all_demyelinating_disease_cns Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases Devic's syndrome http://nn.neurology.org/cgi/collection/devics_syndrome
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2015 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.