Masato Kadoya, MD
Akiko Kadoya, MD
Hiroyuki Onoue, MD,
PhD
Katsunori Ikewaki, MD,
PhD
Jun Shimizu, MD, PhD
Kenichi Kaida, MD, PhD

*Neurol Neuroimmunol
Neuroinflamm*
2016;3:e220; doi: 10.1212/
NXL.0000000000000220

REPEATED TREATMENTS WITH RITUXIMAB IN A PATIENT WITH AMYLOID-LIKE IgM DEPOSITION NEUROPATHY

OPEN

Amyloid-like immunoglobulin M (IgM) deposition neuropathy associated with Waldenström macroglobulinemia is a rare phenotype of IgM-related neuropathy. The prominent clinical features are painful sensory-dominant neuropathy followed by distal motor weakness and atrophy over a long clinical course^{1–4} that are refractory to immunosuppressive treatment in many patients.^{1–6} Herein, we report a case of amyloid-like IgM deposition neuropathy that was successfully treated with rituximab (RTX).

Classification of evidence. This provides Class IV evidence that in patients with amyloid-like IgM deposition neuropathy associated with Waldenström macroglobulinemia, RTX therapy ameliorates motor neuropathy along with cutaneous macroglobulinosis and repeated RTX prevents their exacerbations. This was a single observational study without controls.

Case report. A 63-year-old man presented with weakness of his left fingers, which had started 3 days earlier. He had experienced tingling and numbness in distal portions of all extremities over the past 3 years and intermittent sharp pain in the lower limbs several times a day over the past 2 years. Physical examinations revealed left finger weakness without atrophy in the ulnar nerve region and symmetrical mild numbness with a glove and stocking distribution. He also had small reddish skin rashes on the distal legs, which changed into discrete blister-like papules within 2 to 3 weeks (figure, A). Laboratory examinations showed the presence of IgM- λ monoclonal gammopathy (IgM 1,231 mg/dL) and an increased number of lymphoplasmacytic cells in bone marrow, which confirmed the diagnosis of Waldenström macroglobulinemia. A nerve conduction study revealed a reduced compound muscle action potential (CMAP) (7.4 mV) in the left ulnar nerve (LUN) with normal distal latency and slightly decreased conduction velocity. Sensory nerve responses in the left median, ulnar, and bilateral sural nerves were absent.

Microscopic analysis of a sural nerve biopsy revealed amorphous amyloid-like aggregates in the

endoneurium and subperineurium that were stained by an anti-IgM antibody (figure, B and C). A semithin section showed severe fiber loss (figure, D), and the aggregates were found to be composed of a filamentous material in electron microscopy (figure, E). Biopsy of the duodenal mucosa and skin rashes showed the same aggregates (figure, F and G). Amyloid-like IgM deposition neuropathy with cutaneous macroglobulinosis was diagnosed.⁷

Initial treatment with IV immunoglobulin was ineffective. CMAP in LUN further worsened to 2.3 mV. RTX infusion therapy (375 mg/m², 4 weekly infusions) was introduced, and led to gradual improvement of weakness of the left fingers over 5 months, along with a significant decline of intermittent sharp pain in the lower limbs. The blister-like papules in the lower limbs also ameliorated and changed to pigmented macules. The serum IgM level was markedly decreased (522 mg/dL) 5 months after the start of RTX.

Nine months after the start of RTX, CMAP in LUN had increased (11.0 mV), but the left extensor pollicis longus muscle showed gradual worsening of weakness with relapse of skin rashes and an increase in serum IgM (1,045 mg/dL). The second RTX therapy improved the weakness and skin rashes, but mild sensory disturbances in the distal limbs were unchanged. The patient has undergone RTX therapy every 6 months for the following 2 years with no relapse of finger weakness and skin rashes. The serum IgM normalized 1 year after the reintroduction of RTX therapy and has subsequently remained within the normal range.

Discussion. Clinical courses and treatment have been reported for 7 cases of amyloid-like IgM deposition neuropathy.^{1–6} Sensory disturbances preceded development of weakness by 6 months to 20 years in 5 cases.^{1–4} Immune-mediated treatments, such as prednisolone,^{1,3} chlorambucil,^{2,6} and cyclophosphamide,^{2,4,5} plasma exchange,^{1,6} and RTX infusion,^{1,4,5} were introduced in an early stage of development of weakness but resulted in poor improvement of limb weakness in 6 cases.^{1–3,5,6} One patient with mononeuropathy multiplex after 17 years of sensory neuropathy showed almost complete recovery of power after oral administration of cyclophosphamide.⁴

(A) Small reddish rashes (arrows) and blister-like papules (arrowhead) in the distal legs. (B) Hematoxylin & eosin staining of a sural nerve biopsy specimen. Extensive deposits of eosinophilic aggregates were present in the endoneurium and subperineurial spaces, chiefly around endoneurial microvessels. (C) Immunohistochemical analysis revealed that the deposits were clearly stained by anti-immunoglobulin M antibody. (D) On a semithin section stained with toluidine blue, a severe reduction in density of large myelinated fibers was observed. The aggregates (white asterisks) appeared to compress residual nerve fibers. (E) In electron microscopy, the aggregates existed in island-shaped forms and were composed of a randomly arranged filamentous material (inset), which was morphologically different from spicular amyloid fibrils. (F, G) Hematoxylin & eosin staining of duodenum and skin biopsy specimens. Abundant amorphous eosinophilic material was diffusely deposited in interstitial areas in the duodenum (F) and dermis (G). Scale bars = 50 μ m in B, C, D, F, G and 2.5 μ m in E.

The pathophysiology of amyloid-like IgM deposition neuropathy is unclear and may be heterogeneous. In addition to a direct effect of deposited materials on nerve fibers, a vasculitic process is suggested by clinicopathologic features such as acute-onset multiple mononeuropathy^{4–6} and epineurial inflammatory changes.¹ In our patient, sural nerve biopsy findings indicated that sensory polyneuropathy is likely to result from progressive deposition of amyloid-like IgM and subsequent compression of nerve fibers. The acute-onset left ulnar neuropathy might emerge through an acute inflammatory process secondary to IgM deposition, given the time course. The favorable response of left ulnar neuropathy to RTX therapy may be induced by amelioration of secondary nerve inflammation as well as reduction of IgM production and deposition. Our patient is unique in that RTX therapy was effective for acute-onset motor weakness along with cutaneous macroglobulinosis and that repeated RTX infusions were beneficial for preventing exacerbations.

From the Department of Neurology and Anti-aging Medicine (M.K., A.K., H.O., K.I., K.K.), National Defense Medical College; and Department of Neurology (J.S.), The University of Tokyo, Japan.

Author contributions: Dr. M. Kadoya: study concept and design. Dr. A. Kadoya: acquisition of data. Dr. Onoue: analysis and interpretation. Dr. Ikewaki: study supervision. Dr. Shimizu: analysis and interpretation, revision of the manuscript. Dr. Kaida: critical revision of the manuscript for important intellectual content.

Study funding: No targeted funding.

Disclosure: The authors report no disclosures. Go to Neurology.org/nn for full disclosure forms. The Article Processing Charge was paid by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially.

Received October 13, 2015. Accepted in final form February 11, 2016.

Correspondence to Dr. Kaida: adiak901@ndmc.ac.jp

1. Figueroa JJ, Bosch EP, Dyck PJ, et al. Amyloid-like IgM deposition neuropathy: a distinct clinico-pathologic and proteomic profile disorder. *J Peripher Nerv Syst* 2012;17:182–190.
2. Iwashita H, Argyrakakis A, Lowitzsch K, et al. Polyneuropathy in Waldenström's macroglobulinaemia. *J Neurol Sci* 1974; 21:341–354.

3. Yee WC, Hahn AF, Hearn SA, et al. Neuropathy in IgM λ paraproteinemia. *Acta Neuropathol* 1989;78:57–64.
4. Leschziner GD, Roncaroli F, Moss J, et al. Nineteen-year follow-up of case of mononeuritis multiplex onset with rituximab therapy for Waldenström's-associated neuropathy and Bing-Neel syndrome. *Muscle Nerve* 2009;39:95–100.
5. Luigetti M, Frisullo G, Laurenti L, et al. Light chain deposition in peripheral nerve as a cause of mononeuritis multiplex in case of mononeuritis multiplex onset with rituximab therapy for Waldenström's macroglobulinemia. *J Neurol Sci* 2010;291:89–91.
6. Lamarca J, Casquero P, Pou A. Mononeuritis multiplex in Waldenström's macroglobulinemia. *Ann Neurol* 1987;22:268–272.
7. Gressier L, Hotz C, Lelièvre JD, et al. Cutaneous macroglobulinosis. *Arch Dermatol* 2010;146:165–169.

Neurology® Neuroimmunology & Neuroinflammation

Repeated treatments with rituximab in a patient with amyloid-like IgM deposition neuropathy

Masato Kadoya, Akiko Kadoya, Hiroyuki Onoue, et al.
Neurol Neuroimmunol Neuroinflamm 2016;3;
DOI 10.1212/NXI.0000000000000220

This information is current as of March 31, 2016

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2016 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/3/3/e220.full.html
References	This article cites 7 articles, 0 of which you can access for free at: http://nn.neurology.org/content/3/3/e220.full.html#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://nn.neurology.org/content/3/3/e220.full.html#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://nn.neurology.org/cgi/collection/all_clinical_neurology Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases Class IV http://nn.neurology.org/cgi/collection/class_iv Hematologic http://nn.neurology.org/cgi/collection/hematologic Peripheral neuropathy http://nn.neurology.org/cgi/collection/peripheral_neuropathy
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2016 American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.