Maraviroc in PML-IRIS

A separate ball game under HIV infection and natalizumab?

OPEN

Israel Steiner, MD
Felix Benninger, MD

Correspondence to
Dr. Steiner:
israels@ekmd.huji.ac.il

*Neurol Neuroimmunol
Neuroinflamm*
2017;4:e331; doi: 10.1212/
NXL.0000000000000331

Progressive multifocal leukoencephalopathy (PML) is a severe, often fatal, opportunistic infection of the CNS. First reported in 1958 as a white matter disorder in 3 patients with lymphoproliferative disorders,¹ subsequent studies revealed a polyomavirus, named John Cunningham (JC) virus from the initials of the patient from whose brain it was initially isolated, as the causative agent. Despite previous reports, there is no evidence that PML is caused by either SV40 or BK virus.²

The pathogen is a ubiquitous DNA virus that infects only humans. Subclinical infection often takes place within the first decade of life. Both the mechanism of infection and clinical expression of primary infection remain undefined. Initially, predominantly associated with lymphoproliferative disorders, with the AIDS pandemic, PML became strikingly prevalent, observed in 4%–5% of all patients with HIV prior to the availability of highly active antiretroviral therapy (HAART). The era of monoclonal antibodies for immune-mediated conditions such as natalizumab (Tysabri) for MS and Crohn disease and efalizumab (Raptiva) for psoriasis heralded another context for PML. As of November 30, 2016, there have been 698 reported cases of PML under natalizumab.³

While HAART has a significant beneficial impact on the prognosis of PML in patients with HIV, it remains one of the 4 most common CNS opportunistic infections affecting patients with AIDS.⁴ The institution of screening patients with MS for seropositivity for JC virus has decreased the prevalence of PML in natalizumab-treated patients with MS. The termination of this therapy and using plasma exchange to remove the monoclonal antibody from the circulation when PML is diagnosed has a beneficial effect on prognosis.

However, the introduction of HAART and the discontinuation of natalizumab led to the recognition of the immune reconstitution inflammatory syndrome (IRIS), an entity that is not unique to PML

and has been observed also with mycobacterial diseases, leprosy, fungal infections, and herpes viruses.⁵ IRIS is the consequence of rapid entry of immune cells into the brain at the time of immune restoration. As many as 23% of patients with AIDS with PML who recently started HAART will develop PML-IRIS.⁶ IRIS has occurred in the majority of patients with natalizumab-associated PML following withdrawal of natalizumab and plasma exchange. The diagnosis of CNS-IRIS in patients with MS with natalizumab-associated PML can be challenging because the deterioration might be attributed to PML, MS, or some other opportunistic infection.⁷

Thus, diagnosis of PML in an immune-compromised patient is associated with a 2-edge challenge and risk: clearing the JC virus from the brain, which requires normalization of the immune state, and reconstitution of immune surveillance while minimizing infiltration of the brain with activated T cells that attempt to control an underlying CNS opportunistic infection.

Treatment of CNS-IRIS is mainly empirical; as of today, there are no randomized controlled trials to support antiviral or immune-modulating therapy in this condition. The suggestion to treat CNS-IRIS with steroids remains at present a hypothesis.⁸

In 2009, a case report on the use of maraviroc, a CCR5 antagonist that has been approved for the treatment of HIV infection to inhibit CCR5-dependent viral docking in AIDS suggested that this agent might be effective against CNS-IRIS.⁹ The observation was not confirmed in a large study.¹⁰

Maraviroc is also associated with blocking of CCR5-mediated inflammation and was reported to be beneficial in graft-vs-host disease in patients undergoing allogeneic bone marrow transplantation.¹¹ It also decreases the amount of CD8 cells in the CNS of patients.¹² This led to an effort to extrapolate the use of maraviroc to PML under monoclonal antibodies therapy for MS.¹² This was associated with reduced trafficking of CCR5-expressing cells into

See e-323 and e-325

From the Department of Neurology, Rabin Medical Center, Beilinson Hospital, Petach Tikva, Israel.

Funding information and disclosures are provided at the end of the editorial. Go to Neurology.org/nn for full disclosure forms. The Article Processing Charge was paid by *Neurology® Neuroimmunology & Neuroinflammation*.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

the CSF and survival of the patient with relatively stable neurologic sequelae.

In this issue of *Neurology: Neuroimmunology & Neuroinflammation*, 2 independent reports are published on 3 natalizumab-treated patients with MS with CNS-IRIS handled with maraviroc.^{13,14} In all 3 patients, the syndrome seems to benefit, in part, from the therapy, but use of steroids (in 2), avoidance of plasma exchange (in all), and young age (in 2) might have contributed to the outcome. Maraviroc is not an innocent compound and is associated with hepatotoxicity, infections, cancer, and cardiovascular disorders.¹⁵

The conflicting results on the outcome of IRIS in maraviroc-treated HIV-infected patients and natalizumab-treated patients with MS may suggest a different pathogenesis and/or mechanism of action of the antiviral agent. However, currently available information is too less to draw any conclusions.

To summarize, the available information on the potential of maraviroc in monoclonal antibody-associated PML is too less to enable any recommendation. More anecdotal reports will eventually point to the possibility that a large phase II is warranted.

STUDY FUNDING

No targeted funding reported.

DISCLOSURE

I. Steiner served on the DSMB for Novartis and Actelion and on the editorial board for *Journal of Neurovirology*, *Journal of Neurological Sciences*, and *Neurology: Neuroimmunology & Neuroinflammation*; received research support from Israel Science Foundation, Israel Ministry of Health Chief Scientist, and Teva NNE; and is an expert witness in legal cases involving patients involved in car accidents. F. Benninger received research support from Israeli Science Foundation. Go to Neurology.org/nn for full disclosure forms.

REFERENCES

1. Astrom KE, Mancall EL, Richardson EP Jr. Progressive multifocal leuko-encephalopathy; a hitherto unrecognized complication of chronic lymphatic leukaemia and Hodgkin's disease. *Brain* 1958;81:93–111.
2. Steiner I, Berger JR. Update on progressive multifocal leukoencephalopathy. *Curr Neurol Neurosci Rep* 2012; 12:680–686.

3. chefarztfräulicher: beobachter. *Journal für Asymmetrische Polemic*. Available at: <https://chefarztfrau.de/>. Accessed December 31, 2016.
4. Collazos J. Opportunistic infections of the CNS in patients with AIDS: diagnosis and management. *CNS Drugs* 2003;17:869–887.
5. Lawn SD, Meintjes G. Pathogenesis and prevention of immune reconstitution disease during antiretroviral therapy. *Expert Rev Anti Infect Ther* 2011;9:415–430.
6. Falcó V, Olmo M, del Sá SV, et al. Influence of HAART on the clinical course of HIV-1-infected patients with progressive multifocal leukoencephalopathy: results of an observational multicenter study. *J Acquir Immune Defic Syndr* 2008;49:26–31.
7. Steiner I. PML: underdiagnosed in MS patients on natalizumab. *Lancet Neurol* 2010;9:564.
8. Antoniol C, Jilek S, Schluep M, et al. Impairment of JCV-specific T-cell response by corticotherapy: effect on PML-IRIS management? *Neurology* 2012;79:2258–2264.
9. Martin-Blondel G, Cuzin L, Delobel P, et al. Is maraviroc beneficial in paradoxical progressive multifocal leukoencephalopathy-immune reconstitution inflammatory syndrome management? *AIDS* 2009;23:2545–2546.
10. Sierra-Madero JG, Ellenberg SS, Rassool MS, et al. Effect of the CCR5 antagonist maraviroc on the occurrence of immune reconstitution inflammatory syndrome in HIV (CADIRIS): a double-blind, randomised, placebo-controlled trial. *Lancet HIV* 2014;1:e60–e67.
11. Reshef R, Luger SM, Hexner EO, et al. Blockade of lymphocyte chemotaxis in visceral graft-versus-host disease. *N Engl J Med* 2012;367:135–145.
12. Giacomini PS, Rozenberg A, Metz I, Araujo D, Arbour N, Bar-Or A; Maraviroc in Multiple Sclerosis–Associated PML–IRIS (MIMSAPI) Group. Maraviroc and JC virus-associated immune reconstitution inflammatory syndrome. *N Engl J Med* 2014;370:486–488.
13. Hodecker SC, Stuermer KH, Becker V, et al. Maraviroc as possible treatment for PML-IRIS in natalizumab-treated multiple sclerosis patients. *Neurol Neuroimmunol Neuroinflamm* 2017;4:e325. doi: 10.1212/NXI.0000000000000325.
14. Bsteh G, Auer M, Iglseder S, Walchhofer LM, Dietmar Langenscheidt D, Stefan Koppi S, Schauer-Maurer G, Stockhammer G, Berger T. Severe early natalizumab-associated PML in MS: effective control of PML-IRIS with maraviroc. *Neurol Neuroimmunol Neuroinflamm* 2017;4:e323. doi: 10.1212/NXI.0000000000000323.
15. Woollard SM, Kanmogne GD. Maraviroc: a review of its use in HIV infection and beyond. *Drug Des Devel Ther* 2015;9:5447–5468.

Neurology[®] Neuroimmunology & Neuroinflammation

Maraviroc in PML-IRIS: A separate ball game under HIV infection and natalizumab?

Israel Steiner and Felix Benninger

Neurol Neuroimmunol Neuroinflamm 2017;4;

DOI 10.1212/NXI.0000000000000331

This information is current as of February 8, 2017

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/4/2/e331.full.html
References	This article cites 14 articles, 0 of which you can access for free at: http://nn.neurology.org/content/4/2/e331.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Multiple sclerosis http://nn.neurology.org/cgi/collection/multiple_sclerosis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2017 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.