

A young man with numbness in arms and legs

From the National Multiple Sclerosis Society Case Conference Proceedings

Andrew R. Romeo, MD, Robert P. Lisak, MD,* Ethan Meltzer, MD,*† Edward J. Fox, MD, PhD,* Esther Melamed, MD, PhD,* Ashlea Lucas, PA-C,* Leorah Freeman, MD, PhD,* Teresa C. Frohman, MPAS, MSCS, PA-C,*‡ Kathleen Costello, MS, ANP-BC,*‡ Scott S. Zamvil, MD, PhD,*§ Elliot M. Frohman, MD, PhD,*§ and Jeffrey M. Gelfand, MD, MAS*

Correspondence

Dr. Gelfand
Jeffrey.Gelfand@ucsf.edu
or Dr. E.M. Frohman
Elliot.frohman@austin.utexas.edu

Neurol Neuroimmunol Neuroinflamm 2018;5:e509. doi:10.1212/NXI.0000000000000509

Case presentation

A gentleman in his late 40s developed a “pins and needles” sensation and numbness in his left hand that was exacerbated by leaning on his left arm or hyperextending his neck. The numbness developed over a few weeks. Over the next 2 months, the numbness spread proximally in the left arm, then to the right arm, and in patches over his bilateral upper chest and next to the bilateral posterior thighs. He denied pain, weakness, fatigue, cognitive changes, vision changes, dysarthria, or dysphagia or changes in bladder or bowel function. A review of systems was otherwise negative or normal. Past medical history was notable for hyperlipidemia, treated with atorvastatin 20 mg daily. There was no family history of neurologic or autoimmune disease. Vital signs and general physical examination were normal. Neurologic examination was notable for normal mental status and cranial nerve examinations. Gait, coordination, and the remainder of the motor examinations were normal. Sensation was mildly reduced to light touch circumferentially throughout both arms, in patches over the anterior chest, and over the posterior thighs, with preserved sensation to vibration, pinprick, and temperature. The Romberg sign was not present. Deep tendon reflexes were normal. The plantar response was flexor bilaterally.

Serum testing was negative for aquaporin-4 IgG, and targeted infectious, metabolic, and hematologic studies were unrevealing (table 1). The MRI showed a longitudinally extensive transverse myelitis (LETM, where the T2 hyperintensity extends ≥ 3 vertebral segments)¹ (figure 1). CSF examination showed no pleocytosis, normal glucose, elevated total protein (84 mg/dL), normal IgG index, and zero oligoclonal bands (table 2).

Differential diagnosis

This patient’s symptom onset and evolution were both *subacute*. The differential of a partial myelopathy includes structural (compressive), inflammatory, metabolic, toxic, infectious, paraneoplastic, vascular (especially spinal dural arteriovenous fistula), and malignant causes. Genetic/inherited causes usually present more insidiously.² In *acute* cases of myelopathy where inflammation of the spinal cord is demonstrated by imaging or in CSF, but no more specific

*National Multiple Sclerosis Society Case Conference Proceedings Faculty.

From the Department of Neurology (A.R.R., S.S.Z., J.M.G.) and Program in Immunology (S.S.Z.), University of California San Francisco; Department of Neurology (R.P.L.), Wayne State University, Detroit, MI; Partner’s Neurology Training Program (E.M.); MGH and Brigham and Women’s Hospitals, Harvard Medical School, Boston, MA; and E.M. is now with the Department of Neurology, Dell Medical School at the University of Texas at Austin; Central Texas Neurology Consultants (E.J.F.), and Department of Neurology, Dell Medical School at the University of Texas at Austin; Department of Neurology (E.M., A.L.), and Department of Neurology and Ophthalmology (T.C.F., E.F.), Dell Medical School at the University of Texas at Austin; Department of Neurology (L.F.), University of Texas at Houston; and the National Multiple Sclerosis Society, New York, NY.

Funding information and disclosures are provided at the end of the article. Full disclosure form information provided by the authors is available with the full text of this article at Neurology.org/NN.

†Guest Assistant Editor: Ethan Meltzer, MD.

‡Guest Managing Editors: Kathy Costello, MS, ANP-BC and Teresa C. Frohman, MPAS, MSCS, PA-C.

§Guest Editors-in-Chief: Elliot M. Frohman, MD, PhD, and Scott S. Zamvil, MD, PhD.

The Article Processing Charge was funded by the National Multiple Sclerosis Society.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Table 1 Laboratory results: blood

	Result	Reference range
Erythrocyte sedimentation rate	4 mm/h	2–28 mm/h
C-reactive protein	1.5 mg/L	<6.3 mg/L
Antinuclear antibodies	1:160, speckled	<1:40
Anti-proteinase 3	<10 CU	<20 CU
Anti-myeloperoxidase	<10 CU	<20 CU
Smith antibody	<10 CU	<20 CU
RNP antibody	<10 CU	<20 CU
Aquaporin-4 IgG	Not detected	
Vitamin B12, ng/L	388	211–911
Methylmalonic acid, $\mu\text{mol/L}$	0.19	<0.3
Copper, $\mu\text{g/dL}$	76	70–140
Alpha tocopherol, mg/L	15.7	5.7–19.9
Beta and gamma tocopherols, mg/L	1.7	<4.3
Lactate dehydrogenase, U/L	94	102–199
Serum protein electrophoresis and immunofixation	Normal pattern/Negative	
HIV antigen/antibody	Negative	
Nontreponemal screen (RPR)	Nonreactive	
Lyme disease antibody total (EIA)	Negative	
Coccidioides antibody immunodiffusion	Negative	
<i>Bartonella henselae</i> IgG and IgM	Negative	
<i>Bartonella quintana</i> IgG and IgM	Negative	
Quantiferon Gold	Negative	

Abbreviations: EIA = enzyme-linked immunosorbent assay; RNP = ribonucleoprotein; RPR = rapid plasma reagin.

etiology is found, the label *idiopathic acute transverse myelitis* may be applied, typically reaching nadir in 4–21 days.³ This case presented as a subacute partial myelitis.

Table 2 Laboratory results: CSF


	Result	Reference range
White blood cell count, cells/ μL	4	0–5
Red blood cell count, cells/ μL	3	0–10
Protein, total, mg/dL	84	15–45
Glucose, mg/dL	53	40–80
Oligoclonal bands	No bands identified	
IgG index	0.49	<0.66
VDRL	Nonreactive	

Abbreviation: VDRL = Venereal Disease Research Laboratory.

A key clinical feature of this patient's presentation is the relative lack of early major clinical deficits despite such an extensive longitudinal spinal cord lesion, which favors neurosarcoidosis, differing from neuromyelitis optica spectrum disorder (NMOSD) pathophysiology (which typically causes extensive tissue destruction all along the lesion, and profound early deficits). While LETM is exceptionally uncommon in MS, it is highly characteristic of NMOSD and can be associated with other inflammatory myelitides, particularly with neurosarcoidosis. Dorsal subpial post-gadolinium enhancement is characteristic of neurosarcoidosis myelitis when compared to NMOSD, whereas the ring configuration of enhancement is highly reminiscent of the inflammatory pattern most commonly affiliated with those diagnostic entities under the rubric of the NMOSD.⁴ The “trident sign” describes central canal enhancement with dorsal subpial enhancement in neurosarcoidosis myelitis.⁵ This pattern of enhancement can also be seen with CNS infection (including granulomatous infection) and lymphoma.

The circumferential, nondermatomal pattern of sensory loss favors a CNS lesion. The contiguous spread from arms to chest

Figure 1 MRI of inflammatory myelitis before and after treatment



(A and B) Sagittal and axial T2-weighted images that reveal hyperintensity from C4 to C7 involving the central gray and dorsal white matter bilaterally. T1 post-gadolinium images revealed partial, dorsal enhancement of the lesion with likely pial involvement (C and D) with some involvement of the leptomeninges (arrowhead). Repeat MRI after 2 months of oral prednisone (E–H) shows near resolution of the hyperintense lesion previously extending from C4 to C7, but persistent dorsal enhancement likely indicating a nidus of active granulomatous inflammation.

to legs, yet sparing the face, suggests an evolving or expanding cervical spinal cord process below the level of the spinal trigeminal nucleus, which carries pain and temperature as low as C2–C4.² The syndrome points towards a *partial* cervical myelopathy with only 1 of the 3 main spinal cord pathways affected (sensory, specifically dorsal column, but not motor or bowel/bladder), as opposed to a true transverse myelopathy.⁶

Given a strong clinical suspicion and high pretest probability for neurosarcoidosis causing his myelitis, a chest CT with IV contrast was performed. The chest CT revealed bilateral hilar and mediastinal calcified lymph nodes and perilymphatic pulmonary nodules, consistent with pulmonary sarcoidosis. Fine needle aspiration revealed rare nonnecrotizing granulomas consistent with sarcoidosis, with no evidence of infection or malignancy.

Final diagnosis

Probable neurosarcoidosis, manifesting as a partial longitudinally extensive transverse cervical myelitis, supported by biopsy-confirmed pulmonary sarcoidosis.

Discussion

Pathobiological mechanisms of noncaseating granulomatous inflammation

CNS involvement from sarcoidosis occurs in approximately 5%–15% of sarcoidosis patients and can manifest with variable

combinations of leptomeningitis, meningoencephalitis, pachymeningitis, optic neuropathy, other cranial neuropathies, hypothalamic/pituitary involvement, myelitis, or radiculitis.⁷ While sarcoidosis is classically and formally described as a multisystem disease, about 10%–20% of neurosarcoidosis cases have seemingly exclusive CNS involvement.^{7,8}

On MRI, neurosarcoidosis spinal cord parenchymal involvement can appear longitudinally extensive, smaller segmental or multifocal.^{7,9} In addition to the enhancement pattern seen in this case, there may also be enhancement involving the central canal, nerve roots, meninges, or other parts of the CNS if also affected by neurosarcoidosis.⁹ Neurosarcoidosis lesions can exhibit persistent T1 post-gadolinium enhancement for months or years at a time, even with treatment, whereas inflammatory-demyelinating lesions of MS and NMO typically remit within 1–2 months.^{4,8,10,11}


Our patient's chest CT revealed bilateral hilar and mediastinal calcified lymph nodes and perilymphatic pulmonary nodules, consistent with pulmonary sarcoidosis. If the CT is negative, a whole-body Fludeoxyglucose Positron Emission Topography (FDG-PET) can be diagnostically valuable to look for metabolically active but still normal sized lymph nodes that may be targets for biopsy. A skin examination looking for evidence of cutaneous sarcoidosis and eye examination (and sometimes conjunctival biopsy) may also be helpful in this context.

Angiotensin-converting enzyme (ACE) level is commonly considered in the diagnostic evaluation for sarcoidosis, but it is a nonspecific marker. While serum ACE tends to be higher on average in patients with sarcoidosis (and especially active pulmonary sarcoidosis) compared to those without sarcoidosis, sensitivity for sarcoidosis is low, 29%–60%, with specificity of about 89%.^{12–15} Similarly, in the CSF, sensitivity and specificity of ACE for neurosarcoidosis are 24%–55% and 94%, respectively.^{16,17} In summary, a normal ACE should not exclude neurosarcoidosis, and an elevated ACE can be

nonspecific and sometimes seen in association with other inflammatory, infectious, malignant, or metabolic processes or polymorphisms in the ACE gene.

The inflammation of sarcoidosis is characterized by well-formed, noncaseating (necrotizing) granulomas containing monocytes and macrophages, T lymphocytes, B lymphocytes, and fibroblasts, among other cell types (figure 2).^{18,19} Granulomas in the CNS tend to have a perivascular predilection. The granulomatous inflammation of sarcoidosis

Figure 2 Putative mechanisms of noncaseating granulomatous inflammation in neurosarcoidosis



This figure illustrates putative mechanisms the assembly and organization of the complex coordination of putative cellular and molecular mechanisms, which are thought to represent the pathobiological underpinnings for noncaseating granulomatous inflammation in neurosarcoidosis. Immune cells traffic into the “target tissue” via arterioles and can subsequently exhibit properties of antigen presentation. A collection of various immune cell types (e.g., B and T cells, macrophages, and plasma cells) acquire an affinity to become part of what we analogize as an “island of inflammatory cells,” delimited by a perimeter principally composed of hyaline collagen (shown on the figure). As opposed to granulomatous inflammation associated with tuberculosis and other processes, those compositional cellular elements in sarcoidosis usually do not undergo necrotic granulomatous transformation. The M1 designated macrophage is an important constituent of the sarcoid granuloma, and most particularly with respect to its ability to coordinate the inception and prolongation of “pro-inflammatory” cascades, thereby representing a key feature of the noncaseating granuloma of sarcoidosis. Alternatively, the M2 macrophage is characterized by its ability to provide reciprocal properties, in striking contradistinction, to the M1 macrophage, by exhibiting cardinal anti-inflammatory characteristics, including, but not limited to, the elaboration of a highly stereotyped set of anti-inflammatory cytokines and chemokines. Taken together, the repertoire and heterogeneity of intragranulomatous mononuclear cells serve to orchestrate the immune regulatory networks that provide for both the ignition and the complex coordination of the cellular and humoral factors, which have now become classic hallmarks of granulomatous inflammation. The noncaseating granuloma is equipped with counterbalancing mechanisms (i.e., the inflammatory “braking system”) capable of both high precision attenuation, as well as a corresponding ability to abolish those cascades that serve to provoke and perpetuate granulomatous inflammation in nearly every organ and tissue within the human body. TCR = T cell receptor.

is primarily T cell mediated, and classically considered Th1 driven, but emerging evidence promotes a Th17-driven process, at least in the lungs and mediastinal lymph nodes.^{20,21} Common cytokines involved in signaling in sarcoidosis include IFN γ , TNF α , and various interleukins and chemokines.^{18,19} Environmental and infectious exposures have been proposed as possible contributors to sarcoidosis susceptibility, but none have yet been convincingly demonstrated.¹⁹ Genetic susceptibility to sarcoidosis has been associated with specific human leukocyte antigen alleles, supporting an autoimmune etiology.¹⁹

Several proposed diagnostic approaches to neurosarcoidosis have been used in the literature over the years.^{22,23} Updated consensus diagnostic criteria for neurosarcoidosis were published in 2018.²⁴ Diagnosis of “definite” neurosarcoidosis is supported by a confirmatory biopsy from the nervous system consistent with sarcoidosis in the context of a consistent clinical phenotype and rigorous exclusion of other causes, particularly infection and malignancy. However, CNS biopsy is often not preferable or advisable due to risk of morbidity. A diagnosis of “probable” neurosarcoidosis, as in this case, can be made with a syndrome consistent with granulomatous inflammation of the CNS and a confirmatory biopsy of sarcoidosis from another organ system. Cases in which sarcoidosis is suspected but in which there is no biopsy confirmation are best designated as “possible” neurosarcoidosis.

There are no randomized controlled trials of treatment of CNS neurosarcoidosis. While glucocorticoids are effective for most patients with neurosarcoidosis, the doses needed to achieve or sustain remission can be prohibitive due to glucocorticoid toxicity.²² Common steroid-sparing therapies in clinical practice include methotrexate, azathioprine, mycophenolate mofetil, leflunomide, hydroxychloroquine, and, increasingly, infliximab.²⁵ Retrospective analyses suggest that mycophenolate mofetil may be less effective than methotrexate (at least for preventing “relapse”), but such studies risk confounding by indication.^{25,26} In retrospective analyses, infliximab, a TNF α inhibitor, is associated with favorable outcomes, including some cases refractory to other therapies.⁸

In the patient presented here, given concern about glucocorticoid adverse effects, the patient elected to initiate infliximab with weekly oral methotrexate and tapered off glucocorticoids completely within 4 months with gradual resolution of symptoms without functional limitation. Surveillance MRIs at 7 and 12 months showed complete remission.

Author contributions

A.R. Romeo: conception, drafting of manuscript, critical revision of manuscript. S.S. Zamvil and R.P. Lisak: conception, critical revision of manuscript for intellectual content. E. Meltzer, E.J. Fox, E. Melamed, A. Lucas, L. Freeman,

T.C. Frohman, and K. Costello: critical revision of manuscript for intellectual content. E.M. Frohman: conception, critical revision of manuscript for intellectual content, along with the design and development of figures 1 and 3 in collaboration with the medical illustrator Mr. Jason Ooi. J.M. Gelfand: conception and critical revision of manuscript for intellectual content.

Acknowledgment

The authors wish to express their gratitude to their medical illustrator, Mr. Jason Ooi, for his evidence-based rendition of the putative mechanisms underlying noncaseating granulomatous inflammation in neurosarcoidosis. The authors also wish to acknowledge the UCSF MS EPIC Study Team. The National Multiple Sclerosis Society Case Conference Proceedings is a peer review publishing initiative from the monthly; Diagnostic and Treatment Challenges in MS and Neuroimmunology Webinars; sponsored by the National Multiple Sclerosis Society Fellowship Training Program; and Directed by Elliot M Frohman, MD, PhD.

Study funding

Diagnostic and Treatment Challenges in MS and Neuroimmunology Webinars is sponsored by the National Multiple Sclerosis Fellowship Training Program.

Disclosure

A.R. Romeo received research support from National Multiple Sclerosis Society. R.P. Lisak served on the scientific advisory board from Syntimmune, Alexion, National Multiple Sclerosis Society, Myasthenia Gravis Foundation of America, Guillain-Barre Syndrome/CIDP Foundation; received travel funding and/or speaker honoraria from Teva; is section editor for *Neuroimmunology*, *Clinical Neuropharmacology*; is on the editorial board for *Immunology Research*; receives publishing royalties from Blackwell and Oxford; consulted for GLG Consulting, Insight Consulting, Alpha Sites Consulting, Slingshot Consulting, Informa Pharma Consulting; served on the speaker's bureau for Teva; is immediate past president and member of Consortium of Multiple Sclerosis Centers; is on the Executive Committee of the Board of Directors; is member of the Board of Governors of the Foundation of the Consortium of Multiple Sclerosis Centers Society; is on the Board of Governors for National Multiple Sclerosis Society; received research support from Teva, Alexion, Genentec, Novartis, Medimmune, Chigai, Ra Pharmaceuticals paid to Wayne State University. E. Meltzer reports no disclosures. E.J. Fox served on the scientific advisory board for Acorda, Bayer, Biogen, EMD Serono, Genentech Roche, Novartis, Sanofi Genzyme, Teva; received travel funding and speaker honoraria from Acorda, Biogen, EMD Serono, Genentech Roche, Novartis, Sanofi Genzyme, Teva; consulted for Bayer, Biogen, EMD Serono, Genentech Roche, Novartis, Sanofi Genzyme, Teva; served on the speaker's bureau for Acorda, Biogen, EMD Serono, Genentech Roche, Novartis, Sanofi Genzyme, Teva; received research support from Biogen, Celgene,

Chugai, EMD Serono, Genentech Roche, Mallinckrodt, MedDay, Novartis, Sanofi Genzyme, Teva, TG Therapeutics, National Multiple Sclerosis Society; provided expert testimony for Teva. E. Melamed served on the scientific advisory board for EMD, received research support from William & Ella Owens Medical Research Foundation. A. Lucas and L. Freeman report no disclosures. T.C. Frohman received travel funding and/or speaker honoraria from Novartis, Acorda, Genzyme, Biogen; is on the editorial board for NMSS Publication; receives publishing royalties for UpToDate; consulted for Acorda, Novartis, Genzyme; received research support from NIH and NMSS. K. Costello received travel funding from Advanced Studies in Medicine, Johns Hopkins Medicine. S.S. Zamvil served on data safety monitoring boards for Opexa, BioMS, Teva, and Eli Lilly; is a member of the clinical advisory board for Myelin Repair Foundation; is deputy editor for *Neurology: Neuroimmunology & Neuroinflammation*; holds a patent for Aquaporin-4 peptides; consulted for Biogen, Teva, EMD-Serono, Genzyme, Novartis, Roche; serves on the speaker's bureau for Advanced Health Media, Biogen; received research support from Biogen, Teva, NIH, NMSS, Alexander M. and June L. Maisin Foundation. E.M. Frohman received speaker honoraria from Genzyme, Novartis, Acorda; served on the editorial board for *Journal of Neurology, Journal of Neurology, Neurosurgery, and Psychiatry*; receives publishing royalties from UpToDate, Cambridge University Press; served on the speaker's bureau from Genzyme, Novartis, Acorda; received research support from National Multiple Sclerosis Society. J.M. Gelfand is on the editorial board for *Neurology: Neuroimmunology & Neuroinflammation*; consulted for Biogen; received research support to University of California, San Francisco from Genentech, MedDay, Quest Diagnostics; received research support from National Multiple Sclerosis Society, received compensation for medical-legal consulting; his spouse is an associate editor for *JAMA Neurology*, received honoraria from UpToDate, consulted for Zosano, Eli Lilly, Biohaven, and eNeura. Full disclosure form information provided by the authors is available with the full text of this article at Neurology.org/NN.

Received August 1, 2018. Accepted in final form August 15, 2018.

References

1. Frohman EM, Wingerchuk DM. Clinical practice: transverse myelitis. *N Engl J Med* 2010;363:564–572.
2. Blumenfeld H. *Neuroanatomy Through Clinical Cases*, 2nd ed. Sunderland: Sinauer Associates; 2010.
3. Transverse Myelitis Consortium Working Group. Proposed diagnostic criteria and nosology of acute transverse myelitis. *Neurology* 2002;59:499–505.
4. Flanagan EP, Kaufmann TJ, Krecke KN, et al. Discriminating long myelitis of neuromyelitis optica from sarcoidosis. *Ann Neurol* 2016;79:437–447.
5. Zalewski NL, Krecke KN, Weinschenker BG, et al. Central canal enhancement and the trident sign in spinal cord sarcoidosis. *Neurology* 2016;87:743–744.
6. Cree BA. Acute inflammatory myelopathies. *Handb Clin Neurol* 2014;122:613–667.
7. Krumholz A, Stern BJ. Neurologic manifestations of sarcoidosis. *Handb Clin Neurol* 2014;119:305–333.
8. Gelfand JM, Bradshaw MJ, Stern BJ, et al. Infliximab for the treatment of CNS sarcoidosis: a multi-institutional series. *Neurology* 2017;89:2092–2100.
9. MacLean HJ, Abdoli M. Neurosarcoidosis as an MS Mimic: the trials and tribulations of making a diagnosis. *Mult Scler Relat Disord* 2015;4:414–429.
10. Cotton F, Weiner HL, Jolesz FA, Guttmann CR. MRI contrast uptake in new lesions in relapsing-remitting MS followed at weekly intervals. *Neurology* 2003;60:640–646.
11. Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* 2015;85:177–189.
12. Thi Hong Nguyen C, Kambe N, Kishimoto I, Ueda-Hayakawa I, Okamoto H. Serum soluble interleukin-2 receptor level is more sensitive than angiotensin-converting enzyme or lysozyme for diagnosis of sarcoidosis and may be a marker of multiple organ involvement. *J Dermatol* 2017;44:789–797.
13. Grönhagen-Riska C, Selroos O, Wägar G, Fyhrquist F. Angiotensin-converting enzyme. II: serum activity in early and newly diagnosed sarcoidosis. *Scand J Respir Dis* 1979;60:94–101.
14. DeRemee RA, Rohrbach MS. Serum angiotensin-converting enzyme activity in evaluating the clinical course of sarcoidosis. *Ann Intern Med* 1980;92:361–365.
15. Chopra A, Kalkanis A, Judson MA. Biomarkers in sarcoidosis. *Expert Rev Clin Immunol* 2016;12:1191–1208.
16. Tahmouh AJ, Amir MS, Connor WW, et al. CSF-ACE activity in probable CNS neurosarcoidosis. *Sarcoidosis Vasc Diffuse Lung Dis* 2002;19:191–197.
17. Khoury J, Wellik KE, Demaerschalk BM, Wingerchuk DM. Cerebrospinal fluid angiotensin-converting enzyme for diagnosis of central nervous system sarcoidosis. *Neurologist* 2009;15:108–111.
18. Bagnato F, Stern BJ. Neurosarcoidosis: diagnosis, therapy and biomarkers. *Expert Rev Neurother* 2015;15:533–548.
19. Chen ES, Moller DR. Etiologies of sarcoidosis. *Clin Rev Allergy Immunol* 2015;49:6–18.
20. Ramstein J, Broos CE, Simpson LJ, et al. IFN-gamma-Producing T-helper 17.1 cells are increased in sarcoidosis and are more prevalent than T-helper type 1 cells. *Am J Respir Crit Care Med* 2016;193:1281–1291.
21. Broos CE, Koth LL, van Nimwegen M, et al. Increased T-helper 17.1 cells in sarcoidosis mediastinal lymph nodes. *Eur Respir J* 2018;51:1701124.
22. Zajicek JP, Scolding NJ, Foster O, et al. Central nervous system sarcoidosis—diagnosis and management. *QJM* 1999;92:103–117.
23. Marangoni S, Argentiero V, Tavalato B. Neurosarcoidosis: clinical description of 7 cases with a proposal for a new diagnostic strategy. *J Neurol* 2006;253:488–495.
24. Stern BJ, Royal W, 3rd, Gelfand JM, et al. Definition and Consensus Diagnostic Criteria for Neurosarcoidosis: From the Neurosarcoidosis Consortium Consensus Group. *JAMA Neurol Epub* 2018 Aug 27.
25. Bitoun S, Bouvry D, Borie R, et al. Treatment of neurosarcoidosis: a comparative study of methotrexate and mycophenolate mofetil. *Neurology* 2016;87:2517–2521.
26. Gelfand JM. Demystifying neurosarcoidosis and informing prognosis. *JAMA Neurol* 2017;74:1296–1298.

Neurology[®] Neuroimmunology & Neuroinflammation

A young man with numbness in arms and legs: From the National Multiple Sclerosis Society Case Conference Proceedings

Andrew R. Romeo, Robert P. Lisak, Ethan Meltzer, et al.
Neurol Neuroimmunol Neuroinflamm 2018;5;
DOI 10.1212/NXI.0000000000000509

This information is current as of October 23, 2018

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2018 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.



Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/5/6/e509.full.html
References	This article cites 24 articles, 1 of which you can access for free at: http://nn.neurology.org/content/5/6/e509.full.html##ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://nn.neurology.org/content/5/6/e509.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Immunology http://nn.neurology.org/cgi/collection/all_immunology All Spinal Cord http://nn.neurology.org/cgi/collection/all_spinal_cord Clinical neurology examination http://nn.neurology.org/cgi/collection/clinical_neurology_examination Transverse myelitis http://nn.neurology.org/cgi/collection/transverse_myelitis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2018 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

