

MuSK myasthenia gravis monoclonal antibodies

Valency dictates pathogenicity

Maartje G. Huijbers, PhD, Dana L. Vergoossen, MSc, Yvonne E. Fillié-Grijpma, Inge E. van Es, BASc, Marvyn T. Koning, BSc, Linda M. Slot, MSc, Hendrik Veelken, MD, PhD, Jaap J. Plomp, PhD, Silvère M. van der Maarel, PhD,* and Jan J. Verschuuren, MD, PhD*

Correspondence

Dr. Huijbers
M.G.M.Huijbers@lumc.nl

Neurol Neuroimmunol Neuroinflamm 2019;6:e547. doi:10.1212/NXI.0000000000000547

Abstract

Objective

To isolate and characterize muscle-specific kinase (MuSK) monoclonal antibodies from patients with MuSK myasthenia gravis (MG) on a genetic and functional level.

Methods

We generated recombinant MuSK antibodies from patient-derived clonal MuSK-specific B cells and produced monovalent Fab fragments from them. Both the antibodies and Fab fragments were tested for their effects on neural agrin-induced MuSK phosphorylation and acetylcholine receptor (AChR) clustering in myotube cultures.

Results

The isolated MuSK monoclonal antibody sequences included IgG1, IgG3, and IgG4 that had undergone high levels of affinity maturation, consistent with antigenic selection. We confirmed their specificity for the MuSK Ig-like 1 domain and binding to neuromuscular junctions. Monovalent MuSK Fab, mimicking functionally monovalent MuSK MG patient Fab-arm exchanged serum IgG4, abolished agrin-induced MuSK phosphorylation and AChR clustering. Surprisingly, bivalent monospecific MuSK antibodies instead activated MuSK phosphorylation and partially induced AChR clustering, independent of agrin.

Conclusions

Patient-derived MuSK antibodies can act either as MuSK agonist or MuSK antagonist, depending on the number of MuSK binding sites. Functional monovalency, induced by Fab-arm exchange in patient serum, makes MuSK IgG4 antibodies pathogenic.

*Shared senior authorship.

From the Department of Neurology (M.G.H., Y.E.F.-G., I.E.v.E., J.J.P., J.J.V.), Department of Human Genetics (M.G.H., D.L.V., Y.F.-G., I.E.v.E., S.M.v.d.M.), Department of Hematology (M.T.K., H.V.), and Department of Rheumatology (L.M.S.), Leiden University Medical Center, The Netherlands.

Funding information and disclosures are provided at the end of the article. Full disclosure form information provided by the authors is available with the full text of this article at [Neurology.org/NN](https://neurology.org/NN).

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

AChR = acetylcholine receptor; **Lrp4** = LDL receptor–related protein 4; **MG** = myasthenia gravis; **MIR** = main immunogenic region; **MuSK** = muscle-specific kinase; **NMJ** = neuromuscular junction; **PBMC** = peripheral blood mononuclear cell; **V_H** = heavy chain; **V_L** = light chain.

Myasthenia gravis (MG) is a debilitating autoimmune disease where autoantibodies against neuromuscular junction (NMJ) proteins impair neuromuscular transmission and cause fatigable skeletal muscle weakness. Approximately 5% of patients with MG has autoantibodies against muscle-specific kinase (MuSK).¹ These autoantibodies are predominantly of the IgG4 subclass.² IgG4 is considered an “anti-inflammatory” antibody being unable to bind complement or many Fc receptors on immune cells.³ IgG4 antibodies furthermore exchange Fab-arms with other IgG4 molecules, which renders them functionally bispecific and monovalent, preventing antigen crosslinking and internalization.⁴ Consequently, the pathogenicity of MuSK IgG4 autoantibodies was initially questioned.⁵ However, retrospective longitudinal epitope mapping with polyclonal serum showed that disease severity correlates with IgG4 reactivity against the N-terminal Ig-like 1 domain of MuSK.⁶ Furthermore, passive transfer of purified IgG4 from patients with MuSK MG dose dependently induced muscle weakness in mice.⁷ Last, in vitro studies showed that MuSK IgG4 autoantibodies block MuSK-LDL receptor–related protein 4 (Lrp4) interaction, thereby preventing acetylcholine receptor (AChR) clustering, which explains the impaired neuromuscular transmission in MG.^{8–10} It is now well established that MuSK IgG4 autoantibodies cause MG.

In some patients, low titers of IgG1 and IgG3 MuSK autoantibodies coincide with high levels of IgG4 MuSK autoantibodies. Whether this IgG1 and IgG3 can cause MuSK MG remains enigmatic.^{7,9,11}

To further understand the pathomechanism of MuSK MG and investigate whether the unique functional features of IgG4, like Fab-arm exchange, contribute to the pathogenesis, we isolated and functionally characterized monoclonal MuSK antibodies from patients with MuSK MG.

Methods

Patient selection and study approval

Patients with MuSK MG were recruited in our MG outpatient clinic at the Leiden University Medical Center and were selected based on the presence of a positive MuSK antibody test (RSR Ltd). Both patients were symptomatic and on immunosuppressive treatment, whereas 1 patient had been previously treated with rituximab. The study was conducted in accordance with the Declaration of Helsinki and was approved by the local medical ethics committee. Both patients signed informed consent.

Isolation of monoclonal autoantibodies from patients with MuSK MG

MuSK-binding memory B cells were isolated using a fluorescence-activated cell sorter (FACSaria; BD Biosciences, San Jose, CA) from cryopreserved peripheral blood mononuclear cells (using mouse anti-human monoclonals CD19-BV421 HIB19, CD20-AF700 2H7, CD27-APCHy7 M-T271 all from BD Biosciences; in 0.1 % BSA, 2 mM EDTA/Dulbecco's PBS). To remove dead cells and non-B cells, a dump channel was included (7-AAD,00-6993-50, CD3/FITC UCHT1 BD and CD14 (labeled with 7-AAD from Thermo Fisher, Waltham, MA; CD3/FITC UCHT1 and CD14/FITC MSE2 from BD Biosciences; and CD56/FITC HCD56 from BioLegend, San Diego, CA). Antigen-specific cells were isolated using recombinant MuSK produced in *Escherichia coli*⁶ labeled with R-PE (AS-72113, AnaSpec) and MuSK produced in yeast tagged with DyLight 650 (a kind gift of Konstantinos Lazaridis and Socrates Tzartos, Thermo Fisher). Single sorted cells were cultured on irradiated CD40L cells (a kind gift from Kees van Kooten) in a 96-well plate in complex RPMI medium.¹² After 2 weeks, the medium was tested in duplicate for MuSK antibody production using the MuSK ELISA as described previously.⁶

RNA isolation, cDNA production, and antibody sequence isolation

Single wells containing MuSK antibody–producing cells were lysed with 150 μ L Qiazol, and RNA was isolated using standard chloroform extraction and isopropanol/ethanol precipitation. RNA was rehydrated in 8 μ L H₂O and stored at -80° C until further use. cDNA was directly synthesized (without pre-amplification or purification) using Smartscribe Reverse Transcriptase (Takara Bio Europe) an Oligo-dT40VN primer (5'-AAGCAGTGGTATCAACGCAGAGTACT40VN-3') and a template switching oligo (5'-AAGCAGTGGTATCAA CGCAGAGTACATrGrG + G-3') for 10 rounds of amplification. Full-length V(D)J was obtained by ARTISAN PCR using gene-specific primers (general forward primer 5'-CTT AAGCAGTGGTATCAACGCAGAGTACATG-3', IgG HC 5'-GGAAGGTGTGCACGCCGCTGGTC-3', h κ LC 5'-CTGATGGGTGACTTCGCMG-3', and h λ LC 5'-CACACYAGTGTGGCCTTGTTGGCTTG-3').¹³

Recombinant antibody production, purification, and characterization

Heavy and light chain sequences were ordered at GeneArt (Thermo Fisher) in an IgG1 and IgG4 backbone pcDNA3.1 vector and transfected in suspension FreeStyle HEK293-F cells (R790-07, Thermo Fisher) using Fectin (12347-019,

Thermo Fisher) in FreeStyle™ 293 Expression Medium (12338, Thermo Fisher). To increase transfection and production efficiency, the cells were cotransfected with SV40 large T antigen, hp21, and hp27 (kindly provided by Theo Rispens). After 6 days, culture medium was collected, cell debris was removed by centrifugation, and immunoglobulin G (IgG) was purified using a HiTrap Protein A affinity column (17-0402-01, GE Healthcare) on an Akta Pure (GE Healthcare). Antibodies were dialyzed to PBS, filter sterilized, and stored at -20°C . Fab fragments were generated from these recombinant antibodies using papain according to the manufacturer's instructions (20341, Thermo Fisher).

To determine the binding characteristics of the recombinant antibodies, we performed an epitope mapping ELISA.⁶ Recombinant antibodies were used to immunostain fixed mouse levator auris longus muscle at $1\ \mu\text{g}/\text{mL}$ overnight at room temperature. Synaptic regions were labeled with $0.5\ \mu\text{g}/\text{mL}$ AlexaFluor488-conjugated α -bungarotoxin (B13422, Thermo Fisher), and bound recombinant antibodies were detected with $0.5\ \mu\text{g}/\text{mL}$ AlexaFluor594-conjugated goat anti-human IgG (A110414, Thermo Fisher) in PBS for 1 hour. Muscles were imaged on a Leica SP8 confocal laser scanning microscope and analyzed using LasX software.

To determine the ability of the antibodies to inhibit MuSK phosphorylation and AChR clustering, we exposed C2C12 myotube cultures (Cell Lines Service) to them as described previously.⁸ Differentiated myotubes were stimulated with $0.1\ \text{nM}$ agrin (550-AG-100, R&D systems) in the presence or absence of $100\ \text{ng}/\text{mL}$ recombinant antibodies or Fab fragments. For MuSK phosphorylation data, myotubes were lysed after 30 minutes of exposure, and MuSK was precipitated using $5\ \mu\text{L}/\text{sample}$ rabbit anti-MuSK polyclonal serum (ab94276 or ab94277, a kind gift of Steve Burden) during an overnight incubation at 4°C . Bound antigen-antibody complexes were precipitated using protein A agarose beads (11134515001, Roche), which were extensively washed. Samples were subsequently ran on sodium dodecyl sulfate polyacrylamide gel electrophoresis and transferred to polyvinylidene difluoride membrane. MuSK and phosphorylated MuSK were detected using goat anti-rat MuSK (AF562, R&D systems) and mouse anti-

phosphotyrosine clone 4G10 (05-321, Millipore) as primary antibodies and donkey anti-mouse-680RD (926-68072, Licor) and donkey anti-goat 800 CW (926-32214, Licor). To confirm equal loading, whole cell lysates, that formed the immunoprecipitation input samples, were analyzed for tubulin levels using a mouse anti- α -tubulin antibody (T6199, Sigma) and donkey anti-mouse-800 CW secondary antibody (926-32212, Licor). Bound antibodies were detected using the Odyssey CLx (Licor).

AChR clustering was studied after 16 hours of exposing myotubes to $100\ \text{ng}/\text{mL}$ recombinant antibodies or Fab fragments in the absence or presence of $0.1\ \text{nM}$ agrin. Subsequently, the cells were washed 3 times with differentiation medium (DMEM, 31966 Gibco, 2% heat-inactivated horse serum 26050-088, Gibco, 1% pen/strep and 1% L-glutamine) and incubated with $0.5\ \mu\text{g}/\text{mL}$ AlexaFluor488-conjugated α -bungarotoxin (B13422, Thermo Fisher) in differentiation medium for 30 minutes at 37°C . After staining, cells were fixed in 4% formalin solution for 5 minutes, washed with PBS, and mounted using Hardset Mounting Medium (H-1500, Vector Laboratories). Twenty fields divided over 2 coverslips per condition were randomly selected and imaged with the Leica DM5500 microscope. AChR cluster count and size were analyzed using ImageJ (1.48v). MuSK phosphorylation studies were performed in duplicate, and AChR clustering was performed in triplicate.

Statistics

All data are expressed as mean \pm standard error of the mean. Statistical significance of differences between treatment groups was tested with Student *t*-tests, with corrections for multiple testing wherever appropriate. Differences with *p* values <0.05 were considered statistically significant.

Data availability statement

All data related to this article will be made available on request by a qualified investigator.

Results

Patients with MuSK MG

To obtain MuSK-specific B cells, PBMCs were isolated from 2 patients with MuSK MG. The clinical characteristics of these patients are described in table 1.

Table 1 Clinical characteristics from study participants at the time of PBMC/blood donation

Patient ID	Age (y)	Sex	MuSK titer (nmol/L)	Disease severity (QMG)	Thymus status	Treatment
1	59	M	1.19	9/39	Thymectomy 1987; no thymoma	Prednisone, mycophenolate mofetil, pyridostigmine, plasmapheresis, and 2 courses of rituximab 4 and 2 y before
2	59	V	0.75	Not available	No thymectomy; no thymoma	Prednisone, azathioprine, and plasmapheresis

Abbreviation: MuSK = muscle-specific kinase.

Isolation and genetic characterization of patient-derived MuSK autoantibodies

Antigen-specific single-cell sorting yielded 8 MuSK-binding B cells from 2 patients with MuSK MG. The frequency of circulating MuSK clones was ~7 per 100 million PBMCs for patient 1 and 2.5 per 100 million PBMCs for patient 2. An overview of the isolated MuSK autoantibody characteristics is shown in table 2. From 6 of 7 MuSK autoantibody-producing clones from patient 1, we could derive the variable region sequences of the heavy chain (V_H) and light chain (V_L). For 1 clone, only the V_H region could be sequenced.

Surprisingly, the majority (5/7) of the antibodies isolated were of the IgG1 isotype. We furthermore isolated 1 IgG4 and 1 IgG3 clone. All IgG1 clones and the IgG3 clone used the same V_H and V_L genes (*IGHV3-21* and *IGKV1-39*), suggesting a clonal relationship.

Fab-arm exchange is an important feature of IgG4 and might affect the functional characteristics of MuSK autoantibodies.¹⁴ Previous work on polyclonal purified fractions suggested that MuSK MG IgG4 has the ability to undergo Fab-arm exchange.¹⁴ We sequenced the Fc part of our MuSK monoclonal antibodies and confirmed that the IgG4 clone indeed possessed the serine at position 228 and the arginine 409 required to undergo Fab-arm exchange (data not shown).

N-linked glycosylation of the Fab can be important for antigen binding, e.g., in rheumatoid arthritis ACPA autoantibodies.¹⁵ The NXS/T (where X cannot be a proline) N-linked glycosylation motif was found in the variable region of two of our monoclonal MuSK autoantibodies, suggesting that glycosylation of the Fab is not essential for all MuSK autoantibodies.

Functional characteristics of recombinant MuSK monoclonal autoantibodies

The Ig-like 1 domain of MuSK was previously recognized as the main immunogenic region (MIR) of MuSK in polyclonal patient IgG4 fractions and serum.⁶ For 5 of the patient-derived MuSK antibodies, the epitope mapped to the first Ig-like domain of MuSK (table 2). We have thus far identified no monoclonal autoantibodies against other domains. To establish the functional characteristics of the MuSK autoantibodies, recombinant antibodies were produced from an original IgG1 and IgG4 isolated clone. To assess the importance of autoantibody subclass in MuSK MG, each of these variable regions was subcloned in both an IgG1 and IgG4 backbone. We produced biotin antibodies as a negative control for all experiments.


To assess the ability of recombinant patient-derived IgG1 and IgG4 MuSK antibodies to bind to MuSK at the post-synaptic membrane of NMJs, we performed immunostaining on isolated mouse skeletal muscle. Both the IgG1 and IgG4 versions of the recombinant monoclonal MuSK

Table 2 MuSK autoantibody characteristics

Clone	Pt ID	Isotype	HC VDJ genes	VH mutations V _H	LC VJ genes	VL mutations V _L	Glycosylation motif present	Epitope
11-3D9	1	IgG1k	IGHV3-21*01F IGH3-3*01F IGHJ4*02F	16;6	IGKV1-39*01F IGKJ2*01F	16;4	1 in CDR1 of the HC	Ig-like 1
11-3F6	1	IgG1k	IGHV3-21*01F IGH3-19*01F IGHJ4*02F	19;6	IGKV1-39*01F IGKJ2*02F	30;1	1 in FR3 of the LC	Ig-like 1
11-8G4	1	IgG1k	IGHV3-21*01F IGH3-19*01F IGHJ4*02F	34;5	NA	NA	No	NA
11-7C5	1	NA	NA	NA	NA	NA	NA	NA
13-3B5	1	IgG4λ	IGHV1-2*02F IGH3-10*01F IGHJ5*02F	39;11	IGLV3-25*03F IGLJ2*01F	32;5	No	Ig-like 1
13-3D10	1	IgG1k	IGHV3-21*01F IGH3-16*01F IGHJ4*02F	45;7	IGKV1-39*01F IGKJ2*01F	26;2	No	Ig-like 1
13-4D3	1	IgG3k	IGHV3-21*02F IGH3-16*01F IGHJ4*02F	31;6	IGKV1-39*01F IGKJ2*01F	12;3	No	Ig-like 1
16-8B3	2	IgG1	NA	NA	NA	NA	NA	NA

Abbreviations: FR = frame work; HC = heavy chain; LC = light chain; NA = not available; VH = heavy chain variable region; VL = light chain variable region.

Figure 1 Patient-derived recombinant MuSK antibodies bind mouse NMJs




antibodies clearly bound to NMJs (figure 1, data shown for the IgG4 recombinant antibodies).

Successful neuromuscular transmission depends on properly clustered AChR, a process that is orchestrated through the agrin-Lrp4-MuSK signaling cascade.¹⁶ Agrin is released by the motor nerve terminal and binds Lrp4, which subsequently binds MuSK and thereby stimulates MuSK homodimerization and transphosphorylation. Activation of MuSK phosphorylation stimulates a variety of intracellular signaling cascades of which one culminates in AChR clustering. Purified polyclonal patient IgG4 MuSK autoantibodies inhibit MuSK-Lrp4 interaction, subsequent MuSK dimerization and phosphorylation, and thereby prevent agrin-induced AChR clustering in C2C12 myotube cultures and cause myasthenia in mice.⁷⁻⁹ Surprisingly, patient-derived recombinant MuSK monoclonal antibodies (both IgG1 and IgG4) activated rather than inhibited MuSK phosphorylation (figure 2A). This effect was observed in both the absence and presence of agrin. Activation of MuSK phosphorylation was concentration dependent (figure 2B) and differed slightly between the 2 clones. This shows that patient-derived bivalent monospecific MuSK

antibodies binding the Ig-like 1 domain facilitate dimerization and activation of MuSK in vitro.

Recombinant monoclonal IgG1 and IgG4, however, both engage in bivalent monospecific antibody-antigen interactions. To investigate the functional effects of the bispecificity and functional monovalency of Fab-arm exchanged IgG4 MuSK antibodies in patients, we generated monovalent Fab fragments from these recombinant antibodies by papain digestion. In vitro, these Fab fragments inhibited agrin-dependent MuSK phosphorylation (figure 2C) and AChR clustering similar to patient serum-derived MuSK IgG4 (figure 2D). Despite fully activating MuSK phosphorylation, bivalent monospecific MuSK antibodies partially inhibited agrin-dependent AChR clustering as compared to Fab fragments and a biotin control antibody. Importantly, that AChR clustering could be partially induced using bivalent monospecific antibodies independent from agrin (figure 2D). Thus, monovalent MuSK binding blocks MuSK signaling and the AChR clustering pathway, whereas bivalent monospecific MuSK antibodies fully stimulate MuSK phosphorylation and partially induce or restore AChR clustering in this tissue culture model.

Figure 2 Patient-derived recombinant MuSK antibodies can activate or inhibit MuSK phosphorylation and AChR clustering depending on the antibody valency




Bivalent monospecific recombinant MuSK antibodies (clones 11-3F6 and 13-3B5) activated MuSK phosphorylation in the presence and absence of agrin (A). Activation of MuSK phosphorylation was dose dependent (B). Clone 13-3B5 was slightly more potent compared with 11-3F6. A biotin control antibody did not affect (agrin dependent) MuSK phosphorylation. Monovalent Fab fragments generated from these recombinant MuSK monoclonals inhibited MuSK phosphorylation (C). Agrin-dependent AChR clustering was unaffected when exposed to a biotin control antibody or Fab fragments thereof. Bivalent monospecific recombinant IgG4 MuSK antibodies and Fab fragments inhibited the number of agrin-dependent AChR clustering significantly compared with the biotin control antibody (for IgG4: $p < 0.001$, for Fab fragments: $p < 0.001$, one-way analysis of variance (ANOVA) test, Dunnett corrected). Fab fragments reduced AChR clusters to the level of purified patient IgG4 and the “no agrin” condition. Bivalent monospecific antibodies significantly increased AChR clustering compared with Fab fragment-mediated inhibition that seems independent from agrin (D & E; 11-3F6: $p < 0.001$, 13-3B5: $p = 0.03$, biotin antibody: $p = 0.30$, unpaired t test, no agrin conditions 11-3F6: $p = 0.03$, 13-3B5: $p = 0.20$ one-way ANOVA Dunnett corrected). Data represent mean \pm SEM. Scale bar represents 50 μ m. AChR = acetylcholine receptor; MuSK = muscle-specific kinase.

Discussion

In this study, we provide new and surprising insight on the disease mechanism of MuSK MG by functional characterization of patient-derived monoclonal MuSK antibodies. The major findings of this study are graphically depicted in figure 3. Previous studies with polyclonal purified IgG4 from patients with MuSK MG showed that this IgG4 obstructs Lrp4-MuSK binding.^{8,9} This interaction is normally required for MuSK to become dimerized and phosphorylated. Polyclonal patient IgG4, by blocking Lrp4-MuSK interaction, prevents MuSK dimerization, phosphorylation, and subsequent signaling toward AChR clustering leading to muscle weakness. Experiments by others furthermore confirmed that polyclonal MuSK IgG4 antibodies in patient serum have undergone Fab-arm exchange¹⁴ and thus are functionally bispecific, binding to MuSK in a monovalent manner. We here

show that monovalent Fab fragments, generated from patient-derived monoclonal MuSK antibodies, mimicked monovalent Fab-arm exchanged serum IgG4 and recapitulated the pathomechanism of MuSK MG. Surprisingly, bivalent patient-derived monoclonal MuSK antibodies had the complete opposite effect. These antibodies activated, rather than inhibited MuSK phosphorylation and partially induced AChR clustering. As MuSK phosphorylation and signaling requires MuSK dimerization, our data suggest that bivalent monospecific MuSK antibodies can force this dimerization, thereby stimulating MuSK phosphorylation and AChR clustering. This bypasses the need for agrin-bound Lrp4 to interact with MuSK. Because Fab-arm exchange renders circulating IgG4 functionally monovalent, serum MuSK IgG4 is unable to force dimerization and blocks MuSK signaling. Our data suggest that IgG4 MuSK autoantibodies become pathogenic when, due to Fab-arm exchange, they have become more

Figure 3 Graphical depiction of the conclusions of this study



functionally monovalent. MuSK autoantibody valency and, consequently, MuSK autoantibody subclass are thus key determinants in the pathogenic mechanism underlying MuSK MG. Inhibition of Fab-arm exchange might therefore have therapeutic potential in MuSK MG and perhaps also in other identified IgG4-mediated autoimmune disorders.^{17,18} Our experiments do not exclude the possibility that the low levels of IgG1-3 MuSK antibodies might damage NMJs through activation of complement or immune cell-mediated cytotoxicity.

The agonistic (i.e., bivalent) patient-derived MuSK monoclonal antibodies did not induce AChR clustering in the C2C12 myotube assay to the same degree as the natural agonist agrin. This is likely not a dose-dependent potency difference (data not shown). Alternatively, binding of bivalent monospecific antibodies to the Ig-like 1 domain of MuSK might affect downstream aspects of the AChR clustering pathway differently from agrin-induced MuSK stimulation. These results show strong similarities to a study using IgG obtained from an active immunization model of MuSK MG in rabbits,¹⁹ to studies showing functionally bivalent antibodies binding the MuSK extracellular domain,^{20,21} and to *in vitro* studies with polyclonal purified patient IgG1-3 fractions.⁹ One explanation for the incomplete AChR clustering ability of these MuSK

antibodies may be reduced Dok-7 protein levels, a protein acting downstream of MuSK essential for transducing the AChR clustering signal.¹⁹ It is conceivable that binding of bivalent monospecific MuSK antibodies alters the interaction between MuSK and Dok-7 or might increase MuSK internalization and affect subsynaptic gene expression, ultimately lowering AChR cluster numbers.¹¹ It will furthermore be important to learn whether the level of AChR clustering that can be achieved with patient-derived bivalent monospecific MuSK antibodies is sufficient to maintain adequate neuromuscular transmission *in vivo*.

Recombinant MuSK agonistic antibodies have previously been described.^{1,21,22} These antibodies bound to the Ig-like 1 domain or the Fz-domain of MuSK and also activated MuSK phosphorylation and AChR clustering *in vitro* and *in vivo*.²³ This suggests that MuSK signaling activation by bivalent antibodies might be epitope independent and solely requires forced dimerization of 2 MuSK molecules. Of interest, forced activation of MuSK signaling holds therapeutic promise in neuromuscular disorders characterized by NMJ deficits.²³

MuSK autoantibodies can also inhibit MuSK-collagen Q interaction.²⁴ It will be interesting to study the effect of these

monoclonal antibodies on this interaction. Furthermore, although the Ig-like 1 domain of MuSK is considered the MIR, antibodies to other parts of the ectodomain of MuSK have been found.^{6,25,26} The here described methodology should enable isolation of MuSK monoclonal antibodies binding to different MuSK domains, which will facilitate more detailed analysis of their (pathogenic) effects.

The frequency of MuSK-binding memory B cells was low in the patients studied. This may be explained by the immunosuppressive medication that the patients received. Both patients had, despite this medication, rather high levels of MuSK autoantibodies. Lighaam et al.¹² showed that numbers of circulating IgG4 memory B cells tightly correlated with IgG4 titers. This might, however, not reflect individual antigen-specific responses. As plasma cells mostly reside in the bone marrow, the isolation of circulating memory B cells might not be a representative of the ongoing MuSK autoantibody production. We can also not exclude that the experimental procedures affected the number or isotype of the isolated MuSK-binding cells. Experiments testing the isolation and culture of IgG1 and IgG4 memory B cells using this method did show a slight (~10%) survival advantage for single IgG1 memory B cells (data not shown); however, this is not sufficient to explain the lack of IgG4 MuSK-binding memory B cells isolated in these experiments.

The degree of somatic hypermutation, reflecting affinity maturation, was high in all MuSK antibody clones for both the V_H and V_L, strongly suggesting antigen-driven selection. In 2 recent publications on MuSK monoclonals, B-cell populations, and repertoire analysis in patients with MuSK MG, a different VDJ gene usage was reported.^{27,28}

At least 12 other autoimmune diseases affecting different organ systems are hallmarked by pathogenic IgG4 autoantibodies.^{17,18,29} For several of them, patient-derived monoclonal antibodies have been isolated and characterized.^{30,31} The VDJ gene usage differs between all these diseases. It is not known why these diseases are caused by IgG4 autoantibodies. The role of antibody valency for these diseases has only partially been studied. In pemphigus vulgaris, where the antigen forms a heterodimer (in contrast to MuSK, which forms a homodimer), pathogenicity seems more dependent on autoantibody titer and epitope.¹⁸ It will be interesting to learn whether autoantibody valency and thus Fab-arm exchange is relevant in other IgG4-mediated autoimmune diseases.

In conclusion, the pathomechanism of MuSK MG is dependent on at least 3 factors: (1) autoantibody titer,^{6,32} (2) recognized epitope,^{6,8,22} and, emerging from this and other^{9,19} studies (3) antibody valency/autoantibody subclass. This study thereby presents a new pathomechanism in antibody-mediated autoimmunity, requiring Fab-arm exchange of IgG4 to render autoantibodies monovalent and pathogenic.

Author contributions

M.G. Huijbers designed and analyzed experiments and wrote the manuscript. D.L. Vergoossen and Y.E. Fillié-Grijpma performed and analyzed experiments. I.E. van Es performed experiments. M.T. Koning designed, analyzed, and advised on experiments. L.M. Slot designed and advised on experiments. H. Veelken provided material and wrote the manuscript. J.J. Plomp analyzed the data and wrote the manuscript. S.M. van der Maarel and J.J. Verschuuren designed experiments, analyzed the data, and wrote the manuscript.

Acknowledgment

The authors thank Rochelle Vergroesen, Lise Hafkenscheid, Joris Schonkeren, Prof. Jelle Goeman, and Prof. Rene Toes for their valuable advice on this project, Dr. Konstantinos Lazaridis and Prof. Socrates Tzartos for the MuSK protein, Prof. Steve Burden for the immunoprecipitation antibody, Edwin de Haas for assistance with FACS sorting, Dr. Theo Rispens for his valuable advice and expression vectors, and the Prinses Beatrix Spierfonds (W.OR17-13) and the LUMC for supporting our work financially.

Study funding

This work was supported by the Prinses Beatrix Spierfonds and the LUMC.

Disclosure

M.G. Huijbers is coinventor on several patent applications related to MuSK MG pathogenesis. D.L. Vergoossen, Y.E. Fillié-Grijpma, and I.E. van Es report no disclosures. M.T. Koning received research support from the Professor Steenhuis Fund. L.M. Slot and H. Veelken report no disclosures. J.J. Plomp has 2 patents pending on the pathomechanisms of MuSK MG antibodies. S.M. van der Maarel served on the scientific advisory boards of the Prinses Beatrix Spierfonds, CORT, MRC Centre for Neuromuscular Diseases, Dutch CAA Foundation, and Spierziekten Centrum Nederland; served on the editorial board of *Neuromuscular Disorders*; is coinventor on several patent applications related to FSHD and to MuSK MG; and is a nonexecutive board member of GenomeScan. J.J. Verschuuren served on the scientific advisory board of the Myasthenia Foundation of America; is on the editorial board of the *Journal of Neuromuscular Diseases*; holds a patent for MuSK myasthenia gravis; received research support from the Prinses Beatrix Spierfonds, LUMC receives payments from Alexion, argenx, Ra Pharma due to consultancies, LUMC received royalty payments for development of MuSK Diagnostic Antibody test for TECAN/IBL. Full disclosure form information provided by the authors is available with the full text of this article at Neurology.org/NN.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* December 3, 2018. Accepted in final form January 14, 2019.

References

1. Hoch W, McConville J, Helms S, Newsom-Davis J, Melms A, Vincent A. Autoantibodies to the receptor tyrosine kinase MuSK in patients with myasthenia gravis without acetylcholine receptor antibodies. *Nat Med* 2001;7:365–368.

2. McConville J, Farrugia ME, Beeson D, et al. Detection and characterization of MuSK antibodies in seronegative myasthenia gravis. *Ann Neurol* 2004;55:580–584.
3. Lighaam LC, Rispens T. The immunobiology of immunoglobulin G4. *Semin Liver Dis* 2016;36:200–215.
4. van der Neut Kolfshoten M, Schuurman J, Losen M, et al. Anti-inflammatory activity of human IgG4 antibodies by dynamic Fab arm exchange. *Science* 2007;317:1554–1557.
5. Shiraishi H, Motomura M, Yoshimura T, et al. Acetylcholine receptors loss and postsynaptic damage in MuSK antibody-positive myasthenia gravis. *Ann Neurol* 2005;57:289–293.
6. Huijbers MG, Vink AF, Niks EH, et al. Longitudinal epitope mapping in MuSK myasthenia gravis: implications for disease severity. *J Neuroimmunol* 2016;291:82–88.
7. Klooster R, Plomp JJ, Huijbers MG, et al. Muscle-specific kinase myasthenia gravis IgG4 autoantibodies cause severe neuromuscular junction dysfunction in mice. *Brain* 2012;135:1081–1101.
8. Huijbers MG, Zhang W, Klooster R, et al. MuSK IgG4 autoantibodies cause myasthenia gravis by inhibiting binding between MuSK and Lrp4. *Proc Natl Acad Sci USA* 2013;110:20783–20788.
9. Koneczny I, Cossins J, Waters P, Beeson D, Vincent A. MuSK myasthenia gravis IgG4 disrupts the interaction of LRP4 with MuSK but both IgG4 and IgG1-3 can disperse preformed agrin-independent AChR clusters. *PLoS One* 2013;8:e80695.
10. Otsuka K, Ito M, Ohkawara B, et al. Collagen Q and anti-MuSK autoantibody competitively suppress agrin/LRP4/MuSK signaling. *Sci Rep* 2015;5:13928.
11. Cole RN, Ghazanfari N, Ngo ST, Gervásio OL, Reddel SW, Phillips WD. Patient autoantibodies deplete postsynaptic muscle-specific kinase leading to disassembly of the ACh receptor scaffold and myasthenia gravis in mice. *J Physiol (Lond)* 2010;588:3217–3229.
12. Lighaam LC, Vermeulen E, Bleker T, et al. Phenotypic differences between IgG4+ and IgG1+ B cells point to distinct regulation of the IgG4 response. *J Allergy Clin Immunol* 2014;133:267–270.
13. Koning MT, Kielbasa SM, Boersma V, et al. ARTISAN PCR: rapid identification of full-length immunoglobulin rearrangements without primer binding bias. *Br J Haematol* 2017;178:983–986.
14. Koneczny I, Stevens JA, De RA, et al. IgG4 autoantibodies against muscle-specific kinase undergo Fab-arm exchange in myasthenia gravis patients. *J Autoimmun* 2017;77:104–115.
15. van de Bovenkamp FS, Derksen NIL, van Breemen MJ, et al. Variable domain N-linked glycans acquired during antigen-specific immune responses can contribute to immunoglobulin G antibody stability. *Front Immunol* 2018;9:740.
16. Burden SJ, Huijbers MG, Remedio L. Fundamental molecules and mechanisms for forming and maintaining neuromuscular synapses. *Int J Mol Sci* 2018;19:E490.
17. Huijbers MG, Querol LA, Niks EH, et al. The expanding field of IgG4-mediated neurological autoimmune disorders. *Eur J Neurol* 2015;22:1151–1161.
18. Huijbers MG, Plomp JJ, van der Maarel SM, Verschuuren JJ. IgG4-mediated autoimmune diseases: a niche of antibody-mediated disorders. *Ann N Y Acad Sci* 2018;1413:92–103.
19. Mori S, Yamada S, Kubo S, et al. Divalent and monovalent autoantibodies cause dysfunction of MuSK by distinct mechanisms in a rabbit model of myasthenia gravis. *J Neuroimmunol* 2012;244:1–7.
20. Hopf C, Hoch W. Dimerization of the muscle-specific kinase induces tyrosine phosphorylation of acetylcholine receptors and their aggregation on the surface of myotubes. *J Biol Chem* 1998;273:6467–6473.
21. Herbst R, Burden SJ. The juxtamembrane region of MuSK has a critical role in agrin-mediated signaling. *EMBO J* 2000;19:67–77.
22. Xie MH, Yuan J, Adams C, Gurney A. Direct demonstration of MuSK involvement in acetylcholine receptor clustering through identification of agonist ScFv. *Nat Biotechnol* 1997;15:768–771.
23. Cantor S, Zhang W, Delestrée N, Remédio L, Mentis GZ, Burden SJ. Preserving neuromuscular synapses in ALS by stimulating MuSK with a therapeutic agonist antibody. *Elife* 2018;7:e34375.
24. Kawakami Y, Ito M, Hirayama M, et al. Anti-MuSK autoantibodies block binding of collagen Q to MuSK. *Neurology* 2011;77:1819–1826.
25. Ohta K, Shigemoto K, Fujinami A, Maruyama N, Konishi T, Ohta M. Clinical and experimental features of MuSK antibody positive MG in Japan. *Eur J Neurol* 2007;14:1029–1034.
26. Takamori M, Nakamura T, Motomura M. Antibodies against Wnt receptor of muscle-specific tyrosine kinase in myasthenia gravis. *J Neuroimmunol* 2013;254:183–186.
27. Stathopoulos P, Kumar A, Nowak RJ, O'Connor KC. Autoantibody-producing plasmablasts after B cell depletion identified in muscle-specific kinase myasthenia gravis. *JCI Insight* 2017;2:94263.
28. Vander Heiden JA, Stathopoulos P, Zhou JQ, et al. Dysregulation of B Cell repertoire formation in myasthenia gravis patients revealed through deep sequencing. *J Immunol* 2017;198:1460–1473.
29. Koneczny IA. New classification system for IgG4 autoantibodies. *Front Immunol* 2018;9:97.
30. Di ZG, Di LG, Corti D, et al. Pemphigus autoantibodies generated through somatic mutations target the desmoglein-3 cis-interface. *J Clin Invest* 2012;122:3781–3790.
31. Payne AS, Ishii K, Kacir S, et al. Genetic and functional characterization of human pemphigus vulgaris monoclonal autoantibodies isolated by phage display. *J Clin Invest* 2005;115:888–899.
32. Bartoccioni E, Scuderi F, Minicuci GM, Marino M, Ciaraffa F, Evoli A. Anti-MuSK antibodies: correlation with myasthenia gravis severity. *Neurology* 2006;67:505–507.

Neurology[®] Neuroimmunology & Neuroinflammation

MuSK myasthenia gravis monoclonal antibodies: Valency dictates pathogenicity

Maartje G. Huijbers, Dana L. Vergoossen, Yvonne E. Fillié-Grijpma, et al.

Neurol Neuroimmunol Neuroinflamm 2019;6;

DOI 10.1212/NXI.0000000000000547

This information is current as of February 21, 2019

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/6/3/e547.full.html
References	This article cites 32 articles, 5 of which you can access for free at: http://nn.neurology.org/content/6/3/e547.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases Myasthenia http://nn.neurology.org/cgi/collection/myasthenia
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

