

Toll-like receptor 3 deficiency in autoimmune encephalitis post–herpes simplex encephalitis

Thaís Armangue, MD, PhD,* Benjamin J. Baucells, MD,* Alexandru Vlăgea, MD, PhD,* Mar Petit-Pedrol, PhD, Ana Esteve-Solé, PhD, Angela Deyà-Martínez, MD, PhD, Raquel Ruiz-García, MD, PhD, Manel Juan, MD, PhD, Rebeca Pérez de Diego, PhD, Josep Dalmau, MD, PhD, and Laia Alsina, MD, PhD

Neurol Neuroimmunol Neuroinflamm 2019;6:e611. doi:10.1212/NXI.0000000000000611

Correspondence

Dr. Armangue
tarmangue@
sjdhospitalbarcelona.org
or Dr. Alsina
lalsina@sjdhospitalbarcelona.org

The Toll-like receptor 3 (TLR3) pathway is a key component of the innate immunity that prevents replication of viruses in the CNS. Inborn errors of this pathway (TLR3-pathway deficiency), which includes defects in the genes *TLR3*, *UNC93B1*, *TRIF*, *TRAF3*, *TBK1* and *IRF3*, occur in 10% of patients with herpes simplex encephalitis (HSE),^{1,2} and about 66% of these patients develop relapses of HSE.¹ A recent study showed that 27% of patients with HSE develop autoimmune encephalitis (AE) in the weeks or months ensuing the infection.³ It is unknown whether TLR3-pathway deficient patients can also develop AE post-HSE. Here we report a patient with TLR3-pathway deficiency who developed HSE and a relapse of the viral infection followed by AE post-HSE, highlighting the fact that TLR3-pathway deficient patients should be carefully followed for both HSE relapses and AE.

Case

A 6-year-old healthy Caucasian girl, with a family history of maternal grandfather with recurrent herpetic keratitis, developed acute-onset headache, fever, decreased level of consciousness, aphasia, and right hemiparesis. Brain MRI showed bilateral hemorrhagic temporal lesions leading to suspect HSE. Signs of cranial hypertension precluded CSF studies; however, a positive herpes simplex virus 1 (HSV-1) PCR was obtained in the blood samples. IV acyclovir resulted in symptom improvement, and 3 weeks later the patient was discharged home with residual aphasia.

Five months later, she was readmitted with severe headache and decreased level of consciousness. Brain CT showed new hemorrhagic temporal lesions. She required urgent decompressive craniectomy, and a brain tissue sample was HSV-1 PCR positive. Genetic studies showed that the patient and the mother carried a heterozygous missense mutation p.Glu110Lys (c.328G>A) in exon 2 of the gene *TLR3* (NM_003265.2) (figure, A and B). Such mutation shows a very low allelic frequency (0.00082%, 1/121,316). Functional studies on patient's fibroblasts (figure, C) and monocyte-derived dendritic cells showed a decrease in TLR3-mediated activation (methods in appendix e-1, links.lww.com/NXI/A141). The patient received a new 21-day course of IV acyclovir followed by oral valganciclovir.

One year later, 17 months after the initial episode of HSE, she developed behavioral changes characterized by aggressivity and paranoid thoughts. CSF studies showed pleocytosis (15 cells/ μ L) and elevated protein concentration (100 mg/dL) but were HSV-1 PCR negative. CSF immunochemistry studies on rat brain tissue (figure, D, upper panel) and cultured live neurons showed

*Shared first authorship.


From the Neuroimmunology Program (T.A., M.P.-P., J.D.), Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Hospital Clínic, Universitat de Barcelona, Barcelona, Spain; Pediatric Neuroimmunology Unit (T.A.), Neurology Department, Sant Joan de Déu (SJD) Children's Hospital, University of Barcelona, Barcelona, Spain; Clinical Immunology and Primary Immunodeficiencies Unit (B.J.B., A.D.-M., L.A., A.E.-S.), Pediatric Allergy and Clinical Immunology Department, Hospital Sant Joan de Déu, Institut de Recerca Sant Joan de Déu, Universitat de Barcelona, Spain; Clinical Immunology Unit (A.V., A.E.-S., R.R.-G., M.J., L.A.), Hospital Sant Joan de Déu-Hospital Clínic Barcelona, Barcelona, Spain; Laboratory of Immunogenetics of Human Diseases (R.P.d.D.), IdiPAZ Institute for Health Research, La Paz Hospital, Madrid, Spain; Innate Immunity Group (R.P.d.D.), IdiPAZ Institute for Health Research, La Paz Hospital, Madrid, Spain; Interdepartmental Group of Immunodeficiencies (R.P.d.D.), Madrid, Spain; Neurology Department (J.D.), University of Pennsylvania, Philadelphia; and Catalan Institute for Research and Advanced Studies (ICREA) (J.D.), Barcelona, Spain.

Go to Neurology.org/NN for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by Mutua Madrileña Foundation.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Figure Genetic, functional, and immunologic studies in a 6-year-old child with TLR3-pathway deficiency who developed HSE, and a relapse of the viral infection followed by AE post-HSE



(A and B) Genetic studies identified a missense heterozygous mutation p.Glu110Lys (c.328G>A) in exon 2 of the TLR3 gene (NM_003265.2) in the patient and her mother (who was asymptomatic). The patient's grandfather had history of severe recurrent herpetic keratitis but at the time our patient was studied, he was deceased, and no genetic studies were available. (C) IFN- β production by fibroblasts of the patient (red squares) and a healthy control (black dots) in 2 independent experiments after 24 hours stimulation with 2 different TLR3 agonists: Poly(I:C) or Poly(I:C) LMW; stimulation with its own culture media was used as negative control (negative). IL-6 production by TLR3-agonists stimulated MDDCs was also decreased in the patient compared with a healthy control in 2 independent experiments (data not shown). (D) Rat brain immunostaining with CSF of the patient (upper panel) indicating the presence of antibodies against neuronal surface antigens, compared with that of a healthy control individual (lower panel). Scale bar = 2,000 μ m. AE = autoimmune encephalitis; HSE = herpes simplex encephalitis; IFN- β = interferon-beta; LMW = low molecular weight; MDDC = monocyte-derived dendritic cell; Poly(I:C) = polyinosinic-polycytidylic acid; TLR3 = Toll-like receptor 3.

strong reactivity revealing the presence of neuronal antibodies. Autoantibodies against N-methyl-D-aspartate, γ -aminobutyric acid A, and other known receptors and cell surface proteins were all negative (methods in appendix e-1, links.lww.com/NXI/A141). The indicated patient's antibodies were absent in samples of CSF obtained during HSE (data not shown). With the diagnosis of AE post-HSE, she was started on high-dose IV steroids and immunoglobulins without a clear improvement. Subsequently, rituximab (2 doses 500 mg/m² 2 weeks apart) resulted in neurologic improvement. A CSF sample obtained 4 months later was no longer reactive with brain.

Discussion

We report a young girl with recurrent HSE and a new TLR3 mutation associated with absent interferon- β responses to TLR3 agonist who several months after the first HSE episode

developed recurrent HSE followed by AE. TLR3-pathway deficient patients, especially TLR3-deficient, are prone to develop HSE relapses, putting them at risk of developing AE. Indeed, a recent series of patients with HSE showed that 27% subsequently developed AE; none of them were investigated for TLR3-pathway deficiency but the current case indicates that patients with this deficiency are also at risk of AE. The symptoms of our patient (predominant behavioral abnormalities) are typical of AE post-HSE in patients older than 4 years^{3,4}; yet the long interval between HSE and AE (17 months since the onset of HSE and 12 months since the relapse of HSE) is somewhat atypical (median 26 days in children aged 4 years or younger and 43 days in patients older than 4 years),^{3,5} but similar prolonged intervals occurred in 7% of patients in a recently reported series.³ The exact mechanisms underlying this severe immune-mediated complication are unknown. It has been postulated that the neuronal destruction caused by the virus leads to a release of antigens, which in the context of severe inflammation results in neuronal autoimmunity, such as that shown in our patient.³ In

patients with TLR3-pathway deficiency, the increased number of episodes of HSE theoretically increases the risk of this autoimmune complication. Our case suggests that patients with recurrent HSE or family history of HSE or HSV keratitis should be assessed for TLR3-pathway deficiency.^{1,6,7} If the function of this pathway is impaired, patients should be carefully followed for potential relapses of HSE or AE post-HSE.

Acknowledgment

The authors thank Maria Rodés, Marta Muñoz, Mercè Alba, Eva Caballero, and Esther Aguilar for excellent administrative and technical support.

Study funding

This study was supported in part by Mutua Madrileña Foundation Award (AP162572016, T.A.); Plan Nacional de I+D+i and cofinanced by the ISCIII—Subdirección General de Evaluación y Formento de la Investigación Sanitaria—and the Fondo Europeo de Desarrollo Regional (ISCIII-FEDER; PI18-00486 to T.A.; 17/00234 to J.D., PI15/01094, PFIS0200 (AC16/00025), and PI18/00223 to L.A.; PI14/00616 and PI17/00543 to R.P.d.D.); Pla estratègic de recerca i innovació en salut (PERIS), Departament de Salut, Generalitat de Catalunya (SLT006/17/00362 to T.A.; SLT006/17/00199 to L.A.); a 2017 Leonardo Grant for Researchers and Cultural Creators, BBVA Foundation to L.A.; Ramon Areces Foundation, through a grant from the XVII Concurso Nacional de Ayudas a la Investigación to L.A.; and Fundació CELLEX to J.D.

Disclosure

J. Dalmau holds patents for the use of Ma2, NMDAR, GABAbR, GABAaR, DPPX, and IgLON5 as autoantibody tests and receives royalties related to autoantibody tests from Athena Diagnostics and Euroimmun, Inc., and is editor of *Neurology*[®]: *Neuroimmunology & Neuroinflammation*. The other authors declare no conflict of interest related to this manuscript. Go to Neurology.org/NN for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* June 21, 2019. Accepted in final form July 26, 2019.

Appendix Authors

Name	Location	Role	Contribution
Thaís Armangué, MD, PhD	Sant Joan de Déu (SJD) Barcelona Children's Hospital, IDIBAPS and Hospital Sant Joan de Déu-Hospital Clínic	Author	Design and conceptualized study; performed the neuroimmunologic studies; analyzed the data; follows the patient, drafted the manuscript; developed the figures
Benjamin J. Baucells, MD	Sant Joan de Déu (SJD) Barcelona Children's Hospital	Author	Analyzed the data; drafted the manuscript

Appendix (continued)

Name	Location	Role	Contribution
Alexandru Vlăgea, MD, PhD	Hospital Sant Joan de Déu-Hospital Clínic Barcelona	Author	Performed the TLR3-pathway studies on dendritic cells and genetics, analyzed the data; drafted the manuscript, developed the figures
Mar Petit-Pedrol, PhD	IDIBAPS	Author	Performed the neuroimmunologic studies; analyzed the data; developed the figures, critical review of the manuscript
Ana Esteve-Solé, PhD	Sant Joan de Déu Research Institute and Hospital Sant Joan de Déu-Hospital Clínic Barcelona	Author	Performed the TLR3-pathway studies on dendritic cells, analyzed the data; critical review of the manuscript
Angela Deyà-Martínez, MD, PhD	Sant Joan de Déu (SJD) Barcelona Children's Hospital	Author	Follows the patient, analyzed the data; critical review of the manuscript
Raquel Ruiz-García, MD, PhD	Hospital Sant Joan de Déu-Hospital Clínic Barcelona	Author	Analyzed the data; critical review of the manuscript
Manel Juan, MD, PhD	Hospital Sant Joan de Déu-Hospital Clínic Barcelona	Author	Performed the TLR3-pathway studies on dendritic cells, analyzed the data; critical review of the manuscript
Rebeca Pérez de Diego, PhD	IdiPAZ Institute for Health research	Author	Performed the TLR3-pathway studies on fibroblasts, analyzed the data; developed the figures, critical review of the manuscript
Josep Dalmau, MD, PhD	IDIBAPS, Hospital Clínic and University of Pennsylvania, Philadelphia	Author	Analyzed the data; critical review of the manuscript
Laia Alsina, MD, PhD	Sant Joan de Déu (SJD) Barcelona Children's Hospital and Hospital Sant Joan de Déu-Hospital Clínic	Author	Design and conceptualized study; analyzed the data; follow the patient, drafted the manuscript; developed the figures

References

- Lim HK, Seppänen M, Hautala T, et al. TLR3 deficiency in herpes simplex encephalitis: high allelic heterogeneity and recurrence risk. *Neurology* 2014;83:1888–1897.
- Lafaille FG, Pessach IM, Zhang SY, et al. Impaired intrinsic immunity to HSV-1 in human iPSC-derived TLR3-deficient CNS cells. *Nature* 2012;491:769–773.
- Armangué T, Spatola M, Vlăgea A, et al. Frequency, syndromes, risk factors, and outcome of autoimmune encephalitis following herpes simplex encephalitis: a prospective observational study and a retrospective analysis of cases. *Lancet Neurol* 2018;17:760–772.
- Armangué T, Moris G, Cantarín-Extremera V, et al. Autoimmune post-herpes simplex encephalitis of adults and teenagers. *Neurology* 2015;85:1736–1743.
- Nosadini M, Mohammad S, Corazza F, et al. Herpes simplex virus-induced anti-N-methyl-D-aspartate receptor encephalitis: a systematic literature review with analysis of 43 cases. *Dev Med Child Neurol* 2017;59:796–805.
- Routes J, Abinun M, Al-Herz W, et al. ICON: the early diagnosis of congenital immunodeficiencies. *J Clin Immunol* 2014;34:398–424.
- Abel L, Plancoulaine S, Jouanguy E, et al. Age-dependent Mendelian predisposition to herpes simplex virus type 1 encephalitis in childhood. *J Pediatr* 2010;157:623–629.e1.

Neurology® Neuroimmunology & Neuroinflammation

Toll-like receptor 3 deficiency in autoimmune encephalitis post–herpes simplex encephalitis

Thaís Armangue, Benjamin J. Baucells, Alexandru Vlaga, et al.
Neurol Neuroimmunol Neuroinflamm 2019;6;
DOI 10.1212/NXI.0000000000000611

This information is current as of September 5, 2019

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.



Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/6/6/e611.full.html
References	This article cites 7 articles, 0 of which you can access for free at: http://nn.neurology.org/content/6/6/e611.full.html##ref-list-1
Citations	This article has been cited by 3 HighWire-hosted articles: http://nn.neurology.org/content/6/6/e611.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases Encephalitis http://nn.neurology.org/cgi/collection/encephalitis Post-infectious http://nn.neurology.org/cgi/collection/postinfectious_ Psychosis http://nn.neurology.org/cgi/collection/psychosis Viral infections http://nn.neurology.org/cgi/collection/viral_infections
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

