Expanding the MOG phenotype

Brainstem encephalitis with punctate and curvilinear enhancement

Susan Matesanz, MD, Chelsea Kotch, MD, Christopher Perrone, MD, Angela J. Waanders, MD, Brook Hill, MD, and Sona Narula, MD

Correspondence

Dr. Narula
narulas@email.chop.edu

Neurol Neuroimmunol Neuroinflamm 2019;6:e619. doi:10.1212/NXI.0000000000000619

Myelin oligodendrocyte glycoprotein antibody (MOG-Ab)-associated demyelination has been associated with a range of clinical phenotypes, including acute disseminated encephalomyelitis (ADEM), ADEM followed by recurrent optic neuritis, aquaporin 4-negative neuromyelitis optica spectrum disorder, and less commonly, brainstem encephalitis.¹ In contrast, chronic lymphocytic inflammation with pontine perivascular enhancement responsive to steroids (CLIPPERS) is a clinical, radiologic, and pathologic diagnosis without an associated antibody.^{2,3} There have been few reported cases of pediatric CLIPPERS,⁴ and most have not clearly met the strict case definition of CLIPPERS as proposed in 2017.³ Here, we present a pediatric case, ultimately diagnosed as MOG-Ab-associated demyelination, that was initially concerning for CLIPPERS due to subacute brainstem encephalitis with MRI showing curvilinear and punctate contrast enhancement in the pons.

Case presentation

A 17-year-old previously healthy boy developed headache and transient horizontal diplopia. Over the next 3 days, he developed paresthesias in his left hand, arm, and face. The symptoms rapidly progressed to include dysarthria, gait imbalance, and urinary retention. MRI of the brain demonstrated T2 prolongation in the pons with extension into the cerebral peduncles, with multiple areas of curvilinear and punctate contrast enhancement (figure). MRI of the spine revealed multifocal, patchy T2 signal abnormality throughout the thoracic cord and conus (figure). Although a diagnosis of diffuse midline glioma was initially considered, this was ultimately thought to be less likely due to the subacute onset of symptoms, the noted punctate and curvilinear enhancement, and the presence of spinal cord lesions. CSF analysis was notable for 64/ μ L white blood cells and a protein of 65 mg/dL. Flow cytometric analysis of the CSF showed that 58% of cells were mature lymphocytes. Of the lymphocytes, 90% were T-lymphocytes, with an elevated CD4:CD8 ratio of 8:1. Serum MOG-IgG1 testing, analyzed via a live cell-based flow cytometry assay at Mayo Clinic laboratories, was then sent, following a single dose of dexamethasone.

Because of clinical concern for CLIPPERS, IV methylprednisolone was started on hospital day 2. The patient's symptoms rapidly improved, and his neurologic examination at discharge was normal. Rituximab was started on hospital day 6 as a steroid-sparing agent, and a steroid taper was initiated. Following discharge, the MOG-IgG1 testing sent during admission returned positive with a titer of 1:1,000. A repeat brain and spine MRI performed 3 weeks later demonstrated near-complete resolution of previously seen abnormalities. Repeat MOG-IgG1 antibody testing sent 3 months after presentation remained positive, though with a decreased titer of 1:100. He remains symptom-free 6 months after presentation with no further lesion accrual.

From the Division of Neurology (S.M., S.N.), Division of Oncology (C.K.), and Division of Radiology (B.H.), Children's Hospital of Philadelphia; Department of Neurology (C.P., S.N.) and Department of Pediatrics (S.M., C.K., B.H., S.N.), The Perelman School of Medicine at the University of Pennsylvania, Philadelphia; Division of Hematology, Oncology, and Stem Cell Transplant (A.J.W.), Ann & Robert H. Lurie Children's Hospital of Chicago; and Department of Pediatrics (A.J.W.), Feinberg School of Medicine Northwestern University, Chicago, IL.

Go to Neurology.org/NN for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Figure Sagittal (A) and coronal (B) 3D T1-weighted postcontrast images showing punctate and curvilinear enhancement in the pons

Follow-up sagittal (C) and coronal (D) 3D T1-weighted postcontrast images approximately 3 weeks later showing resolution of enhancement after treatment with IV methylprednisolone and rituximab. Axial T2-weighted images of the thoracic spinal cord (E and F) and conus (G) at the time of presentation showing patchy T2 hyperintense lesions involving gray and white matter.

Discussion

Given the rarity of CLIPPERS in the pediatric population, recognition and evaluation of possible alternative diagnoses is key. The differential for CLIPPERS is broad and includes infectious, inflammatory, and neoplastic processes. Although the case presented above was characterized by a subacute presentation of brainstem symptoms, dramatic response to steroids, and curvilinear enhancement predominating in the pons and cerebellum, it did not meet the strict criteria for CLIPPERS proposed by Tobin et al., as the T2 signal abnormality exceeded the area of contrast enhancement and the MOG-Ab positivity provided an alternative diagnosis. Clinically, as with CLIPPERS, MOG-Ab-associated demyelination is often steroid responsive, and relapses can occur when steroids are weaned.¹ Radiologically, specific features can be suggestive of MOG-Ab-associated disease in the setting of a brainstem encephalitis such as lesions in the posterior fossa greater than 2 cm or lesions with ill-defined margins. As many patients with MOG-Ab-associated disease will also have supratentorial lesions involving the gray and white matter, optic pathway lesions, or spinal cord lesions, comprehensive imaging of the neuroaxis should be performed.⁵

Two previous case reports have reported positive MOG antibodies at the time of relapse in patients previously thought to have CLIPPERS.^{6,7} Although the pathologic

role of MOG antibodies produced by B cells requires further study, the underlying biology of relapses may be different in MOG-Ab-associated demyelination than in CLIPPERS, which is characterized by a predominantly T-cell infiltrate. These differences may have implications for the safety, use, and duration of long-term steroid-sparing therapies.

Early identification of brainstem predominant MOG-Ab-associated disease will lead to better understanding of the clinical phenotype, prognosis, and treatment response. We recommend consideration of MOG-Ab testing in pediatric patients where there is clinical concern for CLIPPERS due to subacute brainstem encephalitis with punctate and curvilinear contrast enhancement, as the prognosis and treatment of MOG-Ab-associated disease and CLIPPERS may ultimately differ.

Study funding

No targeted funding reported.

Disclosure

The authors report no relevant disclosures. Disclosures available: Neurology.org/NN.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* June 5, 2019. Accepted in final form August 15, 2019.

Appendix Authors

Name	Location	Role	Contribution
Susan Matesanz, MD	Children's Hospital of Philadelphia	Author	Cowrote the first draft of the manuscript and critically revised the manuscript
Chelsea Kotch, MD	Children's Hospital of Philadelphia	Author	Cowrote the first draft of the manuscript
Christopher Perrone, MD	Perelman School of Medicine	Author	Critically reviewed and revised the manuscript
Angela J. Waanders, MD	Ann & Robert H. Lurie Children's Hospital of Chicago	Author	Critically reviewed and revised the manuscript
Brook Hill, MD	Children's Hospital of Philadelphia	Author	Compiled figure and reviewed the manuscript
Sona Narula, MD	Children's Hospital of Philadelphia	Author	Conceptualized the study and critically reviewed and revised the manuscript

References

1. Jarius S, Paul F, Aktas O, et al. MOG encephalomyelitis: international recommendations on diagnosis and antibody testing. *J Neuroinflammation* 2018;15:134.
2. Pittock SJ, Debruyne J, Krecke KN, et al. Chronic lymphocytic inflammation with pontine perivascular enhancement responsive to steroids (CLIPPERS). *Brain* 2010;133:2626–2634.
3. Tobin WO, Guo Y, Krecke KN, et al. Diagnostic criteria for chronic lymphocytic inflammation with pontine perivascular enhancement responsive to steroids (CLIPPERS). *Brain* 2017;140:2415–2425.
4. Sa M, Green L, Abdel-Mannan O, et al. Is chronic lymphocytic inflammation with pontine perivascular enhancement responsive to steroids (CLIPPERS) in children the same condition as in adults? *Dev Med Child Neurol* 2019;61:490–496.
5. Jurynczyk M, Gerales R, Probert F, et al. Distinct brain imaging characteristics of autoantibody-mediated CNS conditions and multiple sclerosis. *Brain* 2017;140:617–627.
6. Berzero G, Taieb G, Marignier R, et al. CLIPPERS mimickers: relapsing brainstem encephalitis associated with anti-MOG antibodies. *Eur J Neurol* 2018;25:e16–e17.
7. Symmonds M, Waters PJ, Küker W, Leite MI, Schulz UG. Anti-MOG antibodies with longitudinally extensive transverse myelitis preceded by CLIPPERS. *Neurology* 2015;84:1177–1179.

Neurology[®] Neuroimmunology & Neuroinflammation

Expanding the MOG phenotype: Brainstem encephalitis with punctate and curvilinear enhancement

Susan Matesanz, Chelsea Kotch, Christopher Perrone, et al.
Neurol Neuroimmunol Neuroinflamm 2019;6;
DOI 10.1212/NXI.0000000000000619

This information is current as of September 13, 2019

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/6/6/e619.full.html
References	This article cites 7 articles, 0 of which you can access for free at: http://nn.neurology.org/content/6/6/e619.full.html#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://nn.neurology.org/content/6/6/e619.full.html#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Demyelinating disease (CNS) http://nn.neurology.org/cgi/collection/all_demyelinating_disease_cns All Pediatric http://nn.neurology.org/cgi/collection/all_pediatric Encephalitis http://nn.neurology.org/cgi/collection/encephalitis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.