Vitamin D enhances responses to interferon- β in MS

Xuan Feng, PhD,* Zhe Wang, MD, PhD,* Quentin Howlett-Prieto, BS, Nathan Einhorn, BA, Suad Causevic, BA, and Anthony T. Reder, MD

Neurol Neuroimmunol Neuroinflamm 2019;6:e622. doi:10.1212/NXI.0000000000000622

Correspondence

Dr. Reder
areder@
neurology.bsd.uchicago.edu

Abstract

Objective

To determine the effect of vitamin D3 on interferon- β (IFN- β) response and immune regulation in MS mononuclear cells (MNCs).

Methods

MNCs from 126 subjects, including therapy-naïve patients with different forms of MS, plus patients with MS receiving IFN- β or glatiramer treatment, plus healthy controls were incubated in vitro with IFN- β -1b $\pm$ vitamin D3 (calcitriol). Activation of the IFN- β -induced transcription factor, p-Y-STAT1, and antiviral myxovirus A (MxA) protein was measured with flow cytometry and Western blots; serum proteins were measured with a customized 31-protein multiplex assay.

Results

Vitamin D enhanced in vitro IFN responses, as measured by induction of p-Y-STAT1 and MxA in MNCs, T cells, and monocytes. Vitamin D augmentation of IFN responses was seen in untreated and in IFN- β -1b-treated MS. The combination of vitamin D plus IFN- β reduced Th1 and Th17 cytokines, and increased Th2 responses, reversing the effect of IFN- β alone. Exacerbations and progression in untreated patients reduced the vitamin D enhancement of IFN responses. Vitamin D had less effect on IFN response in clinically stable glatiramer-treated than in IFN- β -treated patients.

Conclusion

Vitamin D enhances IFN- β induction of multiple proteins and also reverses the Th1/Th2 bias in MS seen with IFN- β alone. The combination of vitamin D and IFN- β has potential benefit in ameliorating MS.

*These authors contributed equally to this work.

From the Department of Neurology (X.F., Q.H.-P., N.E., S.C., A.T.R.), University of Chicago Medicine, IL; and Department of Neurology (Z.W.), The First Affiliated Hospital of Dalian Medical University, China.

Go to Neurology.org/NN for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

BDNF = brain-derived neurotrophic factor; **ConA** = concanavalin A; **EAE** = experimental autoimmune/allergic encephalomyelitis; **GA** = glatiramer acetate; **HC** = healthy control; **HGF** = hepatocyte growth factor; **IFN- β** = interferon- β ; **LIF** = leukemia inhibitory factor; **MFI** = median fluorescence intensity; **MNC** = mononuclear cell; **MxA** = myxovirus A; **NGF** = nerve growth factor; **PMA** = phorbol myristate acetate; **PMS** = progressive MS; **PPMS** = primary progressive MS; **RRMS** = relapsing/remitting MS; **RRMS-a** = active RRMS; **RRMS-s** = stable RRMS; **SPMS** = secondary progressive MS; **TPO** = thymopoietin; **VDR** = vitamin D receptor; **VDSI** = Vitamin D Stimulation Index.

Vitamin D in serum controls the onset and activity of MS, and vitamin D affects clinical responses to interferon- β (IFN- β) therapy. The mechanism is unknown.

Vitamin D serum levels correlate with onset of MS and with clinical responses to IFN- β .¹ The chance of developing MS in far Northern and far Southern latitudes and in cloudy regions of France is linked to lack of sunlight and to low serum vitamin D levels.² In established MS, exacerbations and disease progression are linked to low serum vitamin D levels.³ In the BENEFIT study of clinically isolated demyelinating syndrome, high vitamin D levels correlated with fewer second attacks, fewer active MRI lesions, less T2 lesion load, and less brain atrophy.⁴

When exogenous vitamin D is added to IFN- β -1b therapy, there is MRI and clinical benefit. For instance, in a randomized, double-blind trial, 20,000 U/wk of oral vitamin D3 added to IFN- β -1b therapy led to fewer active Gd+ MRI lesions, faster tandem walking speed than at baseline, and a trend for less disability.⁵ In the subgroup with prestudy activity during IFN- β therapy, those supplemented with vitamin D had fewer Gd-enhancing and new T2 lesions and less T2 lesion burden.⁶ In a mix of IFN- β -treated and untreated Tasmanian patients, only those on IFN- β therapy showed clinical benefit from high serum vitamin D levels,⁷ suggesting that there are complex dose/response relationships in vivo. For instance, blacks have lower serum vitamin D levels than whites,⁸ and there was less benefit from 2 forms of IFN- β -1a in blacks than whites.⁹

We hypothesized that combination of vitamin D with IFN- β would enhance IFN-regulated responses. We examined the mechanism of in vitro vitamin D effects on intracellular IFN signaling, antiviral responses, inflammatory cytokines, and neuroprotective proteins in immune cells from untreated and IFN-treated patients with MS and healthy controls (HCs). We demonstrate that vitamin D enhances IFN responses in activated MS immune cells and also provokes a marked shift from Th1 to Th2 responses.

Methods

Subjects

One hundred twenty-six subjects, 111 patients with MS, and 15 HCs were studied. Seventy-two therapy-naïve patients with MS included 33 stable relapsing/remitting MS (RRMS-s), 12 active

RRMS (RRMS-a), 17 secondary progressive MS (SPMS), and 10 primary progressive MS (PPMS). Thirty-nine treated patients with MS were receiving IFN- β (14 RRMS-s, 5 RRMS-a, and 5 SPMS) or glatiramer acetate (GA; 11 RRMS-s and 4 SPMS) (table).

Data availability

All data from this study are included in the article, but anonymized data will be shared by request from any qualified investigator. It is not a clinical trial.

Standard protocol approvals, registrations, and patient consents

Written consent, approved by the University of Chicago institutional review board, was obtained from all subjects.

In vitro IFN response

Mononuclear cells (MNCs) were isolated on lympholyte-H density gradients (Cedarlane Laboratories, Burlington, Canada). In preliminary experiments from 33 sets of MNCs from HCs and patients with MS, IFN- β alone and vitamin D alone had minimal effects on protein secretion by resting cells, but induced robust production by in vitro-activated cells. The dose effect and kinetics of vitamin D (10–400 nM, over 0–48 hours) in these cells were tested and optimized with vitamin D preincubation using replicates at 0, 1, 2, 3, 6, 8, 12, 16, 18, 20, and 24 hours for activation of the p-Y-STAT1 transcription factor and also 0, 24, and 48 hours for induction of the antiviral myxovirus A (MxA) protein. U937 monocytoid and Jurkat T-cell lines exhibited similar responses.

Vitamin D elevated IFN-induced p-Y-STAT1 and MxA at all points, but did not prolong the amplitude after the peak. IFN and IFN plus Vit D both caused maximal rise at 30 minutes, and then, p-Y-STAT1 levels began to decline. Although the IFN + VitD levels were higher at all points, the shape of the kinetic curve was the same. HC and MS MNCs exhibited the same kinetics, with and without vitamin D. MxA protein was induced at 24 and increased at 48 hours before declining. We chose 30 minutes and 48 hours to compare differences between combination therapy and IFN alone.

We also compared media, 5 μ g/mL concanavalin A (ConA) mitogen, or the combination of 5 ng/mL phorbol myristate acetate (PMA) and 1 μ M ionomycin (Sigma-Aldrich). Cytokine induction was minimal with media alone. Cytokine levels were more variable with PMA/ionomycin than with

Table Demographics

	Treatment	n (Total 126)	Age (mean ± SEM)	Sex (male/female)	Race (W/B/O)	Vitamin D (mean ± SEM)	Vitamin D (<30/≥30 ng/mL)
HCs	Ø ^a	15	44.4 ± 4.2	9/6	13W/2O	32.1 ± 3.4	6/9
RRMS-s	Ø	33	46.3 ± 1.9	5/28	24W/7B/2O	37.2 ± 2.8	10/23
RRMS-s	IFN-β	14	48.9 ± 2.8	2/12	12W/2B	39.2 ± 4.0	3/11
RRMS-s	Glatiramer	11	46.6 ± 2.8	1/10	11W	37.4 ± 2.9	1/10
RRMS-a	Ø	12	48.4 ± 2.9	2/10	11W/1B	32.8 ± 4.9	5/7
RRMS-a	IFN-β	5	45.2 ± 6.7	0/5	4W/1B	37.4 ± 11.4	2/3
SPMS	Ø	17	57.7 ± 2.4	7/10	10W/5B/1O	39.5 ± 4.0	5/12
SPMS	IFN-β	5	56.8 ± 5.3	1/4	3W/1B/1O	49.6 ± 6.7	0/5
SPMS	Glatiramer	4	58.7 ± 3.8	1/3	2W/2B	43.5 ± 6.0	1/3
PPMS	Ø	10	54.2 ± 3.2	4/6	8W/1B/1O	34.5 ± 5.0	4/6

Abbreviations: HC = healthy control; PPMS = primary progressive MS; RRMS-a = active RRMS; RRMS-s = stable RRMS; SPMS = secondary progressive MS.
^a Untreated (Ø), Caucasian (W), African American (B), Other (O), and serum 25-OH vitamin D (Vit D).

ConA and exhibited less distinction between Th1, Th2, and other cytokine families than ConA. There were also no differences in kinetic response curves between IFN-β-1b (Bayer) and IFN-β-1a (Serono) at 160 U/mL for 0–2 hours (p-Y-STAT1) or 0–48 hours (antiviral MxA).

For all subsequent experiments, 4×10^6 MNCs/mL in 250 μL AIM-V medium (Life Technologies) were preincubated with 200 nM activated vitamin D3 (1,25 [OH]₂ vitamin D3/calcitriol, Enzo Life Sciences) or media at 37°C in 5% CO₂ for 12 hours and then activated for 2 days with ConA as a model of inflammation. We measured the peak p-Y-STAT1 activation at 30 minutes and MxA at 48 hours after ConA activation, using flow cytometry and Western blots.

Flow cytometry

Stimulated cells were fixed, permeabilized, and stained with monoclonal rabbit antibodies to phospho-STAT1 (Tyr701) (clone 58D6, Alexa Fluor 488 conjugate, Cell Signaling Technology, Danvers, MA) or monoclonal primary rat anti-MxA (Biogen, Cambridge, MA) followed by goat anti-rat immunoglobulin G (IgG) FITC (BioLegend), according to the manufacturer's protocols. To correct for fluorescence intensity variation between experiments, we used multiple internal controls. These included unstained background control, isotype control, medium control, and untreated sample control. Isotype control (Cell Signaling Technology) of rabbit IgG Alexa Fluor 488 was performed to correct for nonspecific binding of p-Y-STAT1 staining. We observed very minimal signal on isotype and unstained controls. Flow cytometry was performed on an LSRFortessa cell analyzer using FACS-Diva software (Becton Dickinson, BD). Numbers of MNCs, lymphocytes, and monocytes, based on forward and side scatter, and median fluorescence intensity (MFI) of protein expression¹⁰ were analyzed with FlowJo software (BD).

After FlowJo software analysis (BD Biosciences), MFI raw values from each condition were exported to an Excel spreadsheet for calculating fold of change. We first normalized all samples by subtracting isotype controls. We then used the Vitamin D Stimulation Index (VDSI) to determine fold change in MFI of protein expression with a 4-component formula, [(IFN+VitD) – VitD alone]/[(IFN alone – media)]. This index measures the additive effect of vitamin D plus IFN-β compared with IFN alone and corrects for differences in induction by IFN alone and for the minor effects of vitamin D alone or media alone.

Western blots

MNCs were lysed in Laemmli buffer plus protease inhibitors and 1 mM Na orthovanadate.¹¹ Cell lysates were run on 8% separating/5% stacking gels and transferred to Immobilon-P polyvinylidene difluoride membranes for enhanced chemiluminescence analysis with luminol/enhancer solution and peroxide solution (Bio-Rad). Western blots were quantitated on a CHEMI Genius Bio Imaging System using GeneSnap Image Acquisition Software and analyzed on Gene tools (Syngene, Frederick, MD). Antibodies were primary rat anti-MxA (Biogen), p-STAT1 (Y701) rabbit MAb (Cell Signaling Technology), actin (I-19) goat polyclonal IgG, donkey anti-goat IgG-horseradish peroxidase (HRP), chicken anti-rat IgG-HRP, and goat anti-rabbit IgG-HRP (Santa Cruz Biotechnology). The relative fold of change between IFN alone and vitamin D added to IFN was calculated with the VDSI, below.

Gel densitometry (ChemiDoc XRS + system imager, Bio-Rad) is based on charge-coupled device high-resolution, high-sensitivity detection, followed by Gene Tools analysis (Syngene). This software automatically adjusts densitometry readouts and maintains the same ratio of MxA/actin regardless of the changing background or sharpness on image. Representative figures capture the intensity at one brightness

level, but the raw data are imaged at a dynamic range of >4.0 orders of magnitude with a 65,535 pixel density (gray levels). During chemiluminescent exposure, care was taken to generate images that were not overexposed to prevent signal plateau/saturation and loss of dynamic range.

We found that Western blot experiments were actually more reproducible than other assays for this intracellular protein. We and other researchers find that Western blots of both p-Y-STAT1 and MxA are reliable markers of IFN responses and also show subtle differences in molecular weight due to posttranslational modifications that are missed with other techniques.¹²

Cytokine profiles

Supernatants from in vitro-activated MNCs with medium alone, vitamin D alone, IFN- β alone, and vitamin D added to IFN- β were measured with 31-cytokine customized multiplex (Procarta Immunoassay Kit, Affymetrix) in 40 patients. Targets were based on the MS literature and our preliminary experiments extracted from 280 candidates, including cytokines, chemokines, and neuroendocrine and neurotrophic proteins. Many of these induced proteins are detectable in vitro, unlike in serum, where most cytokines are at very low levels.¹³ Analytes with significant changes are listed in figure 6. Patient subtypes and HCs were randomized between plates. Assays were run according to the manufacturer's guidelines.

Serum vitamin D levels

Total 25-OH D, 25-OH D3, and 25-OH D2 were quantitated with liquid chromatography/mass spectroscopy (Quest Diagnostics, Schaumburg, IL). Total serum 25-OH D levels less than 20 ng/mL indicate vitamin D deficiency; 20–30 ng/mL suggest insufficiency; 30–100 ng/mL is considered optimal for healthy normal subjects.

Statistical analysis

Values between subject groups were compared with unpaired student *t* tests. Baseline vs later drug-induced values were analyzed with paired *t* tests in all conditions. Figures were generated using R package ggplot2. To compare protein expression after induction with vitamin D added to IFN vs IFN alone, we used the VDSI. It determines fold change in MFI with a 4-component formula, [(IFN+VitD) – VitD alone]/(IFN alone – media). This index measures the additive effect of vitamin D plus IFN- β compared with IFN alone, and corrects for differences in induction by IFN alone, and minor effects of vitamin D or media alone. Statistical significance was delineated as **p* < 0.05 and ***p* < 0.01.

Results

Enhancing effects of vitamin D on intracellular IFN- β responses in MNCs

Untreated MS

IFN- β responses in vitro were first examined by measuring activation of the p-Y-STAT1 transcription factor and

induction of the antiviral MxA protein in ConA-activated MNCs from 72 untreated patients with MS. IFN- β alone induced rapid phosphorylation of p-Y-STAT1, typically a 3-fold increase in p-Y-STAT1 levels compared with media alone (figure 1A). Twelve-hour pretreatment with vitamin D alone had only minimal effects on activation of p-Y-STAT1 (figure 1B). IFN- β plus vitamin D displayed dramatic induction, with a 10-fold increase in p-Y-STAT1 levels compared with IFN- β alone (*p* = 0.00002, paired *t* test).

The effect of vitamin D on intracellular IFN- β response was compared between subgroups of untreated patients with different forms of MS. HCs were the best responders to in vitro vitamin D plus IFN- β and had the highest median elevation in p-Y-STAT1 (*p* < 0.01) (figure 2A). All patient groups except SPMS showed significant vitamin D enhancement of STAT1 activation. There was less induction in SPMS than in HC.

For MxA protein, IFN- β alone showed greater induction than media or vitamin D alone in cells from therapy-naïve patients with MS (figure 1C). IFN- β plus vitamin D induced much more MxA compared with IFN- β alone. Thus, vitamin D activates STAT1 and modifies the downstream response to IFN- β .

Vitamin D enhanced IFN-induced MxA production in all MS groups and in HCs (figure 2B). Therapy-naïve patients with PPMS, with the highest median fold change in (*p* < 0.05, MxA), and patients with SPMS were the best responders to in vitro MxA induction by IFN- β plus vitamin D. The trend for enhanced responses in PPMS compared with HC was not significant. This is because the VDSI corrects for baseline MxA levels when it measures the incremental change from IFN- β alone vs the combination of IFN- β plus vitamin D. IFN- β alone induced far more MxA in HC (14.7-fold induction) than in all forms of MS (2- to 6-fold); RRMS-s had the highest level in untreated patients (described below; figure 6A). This low level of MxA parallels the subnormal response to type I IFN during active MS.¹² Vitamin D corrected the inability of IFN- β alone to induce a robust response. The combination of vitamin D and IFN- β induced MxA more during relapses and progression than in HC.

Treated MS

Vitamin D enhanced IFN- β induction of p-Y-STAT1 levels in MNCs from IFN-treated RRMS-s and RRMS-a, but had no effect in IFN-treated SPMS cells (figure 3). Responses in IFN- β -treated RRMS-s were greater than RRMS-a, and in turn greater than SPMS, paralleling our demonstration of IFN resistance in active and progressive MS.¹² With glatiramer treatment, there was more activation of p-Y-STAT1 in RRMS-s and a trend for higher p-Y-STAT1 in a small number of SPMS (nonsignificant).

There was an age effect in IFN-treated patients with RRMS-s, but not in other groups. Younger patients with RRMS-s on IFN- β therapy (<45 years old) had better in vitro response to IFN- β plus vitamin D (3.5-fold induction) compared with older

Figure 1 Vitamin D enhances IFN- β -induced expression of p-Y-STAT1 and MxA in MNCs

(A) In vitro vitamin D enhances IFN- β induction of p-Y-STAT1 in ConA-activated MNCs from 72 untreated patients with MS ($p = 0.00002$). (B) p-Y-STAT1 expression on flow cytometry histograms from a representative patient with RRMS-s. The media, IFN, VitD, and IFN+VitD curves are all stained with Alexa Fluor 488-labeled Mab to p-Y-STAT1. (C) MxA protein on Western blot of a representative patient with RRMS-s. ConA-activated MNCs were incubated in vitro with 160 U/mL IFN- β -1b for 30 minutes to 48 hours $\pm$ preincubation with 200 nM Vit D3 (calcitriol) for 12 hours. Gray-scale densities of each band are listed above figure. IFN- β = interferon- β ; MxA = myxovirus protein.

patients (1.5-fold induction) ($p < 0.05$). There were no sex or race effects on this response. MxA induction was similar to that seen with p-Y-STAT1 in treated patients (data not shown).

Thus, vitamin D improves in vitro responses to IFN- β in both untreated and in IFN- β -1b and GA-treated patients with MS. There is a lower response to IFN in MNCs from relapsing and progressive patients than in stable patients. This variability could be from the inflammatory milieu in MS or from differences in vitamin D serum levels.

Effect of low vs high serum levels of vitamin D

Subjects were divided into low (<30 ng/mL) and normal/high serum vitamin D (≥ 30 ng/mL) cohorts to compare their responses to in vitro IFN- β -1b plus vitamin D. In both low and high serum vitamin D groups, in vitro vitamin D enhanced activation of STAT1 by IFN- β (figure 4). HCs had good responses, regardless of serum vitamin D levels. In contrast, patients with RRMS-s, SPMS, and PPMS with high vitamin D were better responders than those with low vitamin D levels. This suggests that even on a background of high serum vitamin D levels, additional vitamin D enhances responses to IFN- β . Conversely, during exacerbations, untreated patients with RRMS-a with low serum vitamin D levels had even better in vitro induction of p-Y-STAT1 than those with high levels. This suggests that immune dysregulation and the characteristic low IFN- β responses during exacerbations would benefit from additional vitamin D.

Vitamin D effects on responses to IFN- β in immune cell subsets

Vitamin D affected p-Y-STAT1 induction in lymphocytes and monocytes. Vitamin D plus IFN- β (figure 5, blue line) in most therapy-naïve groups had a greater effect than IFN- β alone (yellow line) on activation of STAT1. However, there is little effect of vitamin D stimulation on lymphocytes from SPMS and RRMS-a (figure 5A), or on monocytes from SPMS (figure 5B), groups with known resistance to IFN- β stimulation.¹² During IFN- β therapy compared with untreated MS, in vitro vitamin D forces induction of additional p-Y-STAT1 in RRMS-s and RRMS-a. This suggests that IFN- β therapy has reduced negative effects of inflammation or has primed immune cells for greater responses to IFN.¹⁴

There were high correlations in expression of p-Y-STAT1 between monocytes and lymphocytes in all groups, after induction by IFN alone ($r = 0.82$, $p = 3.4 \times 10^{-12}$) and by IFN + VitD ($r = 0.91$, $p = 2.4 \times 10^{-17}$), and also in the fold of change ($r = 0.68$, $p = 2.6 \times 10^{-3}$).

Multiplex analysis of vitamin D effects on immune and neuroprotective factors in MS MNCs

Th1, Th17, and Th2 cytokines

A 31-analyte multiplex assay was used to investigate how the vitamin D plus IFN- β combination modulates immune and

Figure 2 Vitamin D enhances intracellular IFN- β responses in therapy-naive MS and HCs

(A) p-Y-STAT1, measured with flow cytometry, is generated in all groups, but there is less induction in SPMS than in HC. (B) MxA protein, measured with Western blots. The trend for enhanced responses in PPMS vs HC was not significant ($p < 0.13$, unpaired t test). ConA-activated MNCs were incubated in vitro with 160 U/mL IFN- β 1b for 30 minutes to 48 hours $\pm$ preincubation with 200 nM vitamin D3 (calcitriol) for 12 hours. Fold change of vitamin D plus IFN- β compared with IFN- β alone, determined using VDSI: $[(\text{IFN} + \text{VitD}) - (\text{VitD})]/[(\text{IFN}) - (\text{no IFN})]$; average with p values above SEM bar: * $p < 0.05$, ** $p < 0.01$. Comparisons between groups use unpaired t tests, brackets. PPMS = primary progressive MS; RRMS-a = active relapsing/remitting MS; RRMS-s = stable RRMS; SPMS = secondary progressive MS.

nonimmune responses. We targeted inflammatory and non-inflammatory cytokines, IFN-stimulated proteins, neurotrophic factors, and cell proliferation and adhesion molecules in 40 subjects (figure 6).

IFN- β alone tended to induce Th1 cytokines. Ratios of IFN/media are in figure 6A, and corresponding difference values in pg/mL are shown in figure e-1A (links.lww.com/NXI/A151). IL-2 was elevated in vitro in untreated and IFN-treated MS MNCs, especially during exacerbations, but not in cells from glatiramer-treated patients. IFN- γ was slightly increased in MNCs from untreated RRMS-a and progressive MS, but not in glatiramer-treated RRMS-s or HC. IFN- β alone inhibited Th2 cytokines (IL-4 in all groups except RRMS-a and HC; IL-5 in all groups except RRMS-a). IL-10

was an exception, with induction by IFN- β in all groups except RRMS-s on IFN therapy (figure 6A). This may be due to differential effects on IL-10 induction among immune subsets. We and others find that IFN- β induces IL-10 in T cells, but inhibits production in monocytes.^{15,16} During attacks, in untreated and IFN- β -treated MS, in vitro IFN- β elevated many Th1 and some Th2 cytokines. Thus, the Th1/Th2 ratio remained elevated during the “cytokine storm” that characterizes exacerbations.

Vitamin D alone, in contrast to IFN- β alone, markedly reduced Th1 and Th17 cytokines for all untreated groups and for IFN- β - and glatiramer-treated MS groups and HC. These cytokines included IL-2, IFN- γ , TNF- α , and TNFRFII, which is secreted by activated T cells, and IL-17F.

Figure 3 Vitamin D enhances IFN- β -induced p-Y-STAT1 expression in MNCs from patients with MS receiving MS therapies

Figure 4 Effect of high vs low serum vitamin D levels from therapy-naive patients with MS on IFN response in MNCs in vitro

Vitamin D, again in contrast to IFN- β , increased Th2 cytokine expression, including IL-4, IL-5, and IL-10 (figure 6B and figure e-1B, [links.lww.com/NXI/A151](https://www.lww.com/NXI/A151)). GA-treated patients were the exception, with no induction of Th2 cytokines compared with the media control, but Th2 cytokine levels were higher than levels seen with IFN- β alone.

Vitamin D added to IFN- β maintained the shift from Th1 to Th2 caused by vitamin D alone, and dramatically reversed the Th1-inducing effect of IFN- β alone, in therapy-naive and IFN-treated MS (figure 6C and figure e-1C, [links.lww.com/NXI/A151](https://www.lww.com/NXI/A151)). Again, the exception was in GA treated, where there was minimal induction. Overall, the change was close to that seen with vitamin D alone. IFN- β plus vitamin D markedly

Figure 5 Vitamin D and IFN- β effects on p-Y-STAT1 induction in lymphocytes (A) and monocytes (B)

Vitamin D plus IFN- β (blue line) in most therapy-naive groups had a greater effect than IFN- β alone (yellow line) on activation of STAT1 in ConA-activated lymphocytes and monocytes. Values in radar plot represent median fold change in log₂ scale expression of p-Y-STAT1 measured by flow cytometry. Blue line: (IFN+ VitD) - (Vit D); yellow: (IFN)-(media alone). IFN- β = interferon- β ; RRMS-a = active relapsing/remitting MS; RRMS-s = stable RRMS; SPMS = secondary progressive MS.

Figure 6 Multiplex of fold change of protein secretion by activated MNCs from HCs and different forms of MS

(A) IFN- β alone vs media, expressed as ratio of IFN- β /media. (B) Vitamin D alone vs media. (C) IFN- β plus vitamin D vs media alone. Selected targets are grouped into Th1, Th17, and Th2 immune pathways and clusters of IFN-stimulated proteins (ISGs), adhesion molecules, and cytokines controlling homeostatic proliferation, neurotrophic factors, and IL-12 family. Values are fold change in the stimulation condition/media. Intensity of shading shows upregulation (red) or downregulation (blue). Target proteins include BDNF = brain-derived neurotrophic factor; HGF = hepatocyte growth factor; IFN- γ = interferon- γ ; IL-2 = interleukin-2; TNF- α = tumor necrosis factor- α ; TNFR1 = TNF receptor type I; IL-17F, IL-4, IL-5, IL-10, IP-10 = IFN-induced protein-10 (CXCL10); MCP1 = macrophage chemotactic protein-1 (CCL2); I-TAC = IFN-inducible T-cell alpha chemoattractant (CXCL11); LIF = leukemia inhibitory factor; p-Y-STAT1 = phosphotyrosine-STAT1 transcription factor; MxA = myxovirus A; NGF = nerve growth factor; sICAM-1 = soluble intercellular adhesion molecule-1; TPO = thymopoietin; VCAM-1 = vascular cell adhesion molecule; VEGF- α = vascular endothelial growth factor- α , IL-15, IL-7, and IL-12 p40 and IL-12 p70 components.

reduced IL-17F in MNCs from HCs and all patients with MS, including those on glatiramer therapy.

Prototypical IFN-stimulated proteins

Induction of p-Y-STAT1 was enhanced 3-fold by vitamin D added to IFN- β in MNCs from HC and groups of therapy-naive and treated MS (figure 6C). MxA was induced in all groups, but much more in HCs than in any MS group (figure 6C). This reflects the low IFN-induced MxA protein levels in MS compared with HC reported by Feng et al.¹² IFN- β and vitamin D, alone and in combination, had only modest effects on chemokine levels in this system.

Adhesion and neurotrophic factors

Vitamin D and IFN- β together tended to stimulate secretion of adhesion molecules intercellular adhesion molecule-1 and vascular cell adhesion molecule-1, as well as vascular endothelial growth factor- α (figure 6C and figure e-1C, links.lww.com/NXI/A151). Although IFN- β alone caused slight induction of neurotrophic factors (leukemia inhibitory factor, brain-derived

neurotrophic factor, nerve growth factor [NGF], hepatocyte growth factor, and thymopoietin), vitamin D plus IFN- β had minimal or inhibitory effects. These neurotrophic factors are predominantly products of the nervous system or other tissues, and serum levels may differ from the modest ability of cultured MNCs to produce these factors. For instance, IFN- β has direct neuroprotective effects by inducing NGF in neurons and astrocytes,¹⁷ but we did not see induction in MNCs.

Multifunctional cytokines

Vitamin D plus IFN- β induced secretion of IL-12(p70), but inhibited secretion of IL-12(p40). IL-12, a monocyte product, induces Th1 cells, predominantly through the actions of the IL-12 p70 heterodimer that consists of p35 and p40 subunits. The IL-12p40 homodimer, however, blocks IL-12 p70 binding to its receptor. These data would suggest that the marked fall in IL-12 p70 and rise in IL-12 p40 would be detrimental in MS. However, there is no therapeutic benefit in MS of anti-IL-12p40 monoclonal antibody (binding IL-12 p40 homodimer, IL-12 p70 heterodimer, and the p40 subunit in IL-12/23).

The lack of efficacy with targeted therapy against IL-12 may arise from the effects on shared subunits of multiple IL-12 family cytokines.

Vitamin D plus IFN- β slightly reduced cytokines that augment homeostatic proliferation (IL-7, IL-15) in most HCs and patients with MS. Reduction of IL-7 may be beneficial in MS. IL-7 induces Th1 cells, and antibodies to IL-7 inhibit experimental autoimmune/allergic encephalomyelitis.¹⁸ IL-15 RNA expression is highly dysregulated in untreated MS.¹⁹ IL-15 induces proliferation of cytolytic CD8 cells,²⁰ which can sever axons in vitro,²¹ yet IL-15 also blunts the function of CD8 cytolytic T-cell.¹⁹ Reduction of IL-15 by IFN- β plus vitamin D potentially counterbalances cytolytic CD8 T-cell function and promotes in vivo and in vitro IFN- β -driven enhancement of regulatory CD8 T-cell function in MS seen in vitro and in vivo.²²

Discussion

Vitamin D enhances responses to IFN- β by increasing activation of p-Y-STAT1. It also reverses or enhances downstream effects of IFN- β alone. Augmentation of signaling by vitamin D occurs in cells from untreated MS (figures 1 and 2) and from patients on IFN- β and glatiramer therapy (figure 3).

Vitamin D alone downregulates proinflammatory Th1 and Th17 cytokines and increases anti-inflammatory Th2 cytokines (figure 6B and figure e-1b, links.lww.com/NXI/A151). IFN- β alone, conversely, clearly increases Th1 and Th17 production and decreases IL-4 and IL-5 secretion. The immune-enhancing effect of IFN- β is not surprising, as type I IFNs are potent antiviral agents, activate cytolytic T cells, and induce Th1 cells in vitro and in vivo.^{23,24}

Addition of vitamin D to IFN- β reversed some of the effects of IFN- β alone. The combination downregulated inflammatory Th1 and Th17 cytokines, but upregulated serum levels of anti-inflammatory Th2 cytokines (IL-4, IL-5, and IL-10), and also induced the MxA antiviral protein and adhesion molecules (figure 6C and figure e-1C, links.lww.com/NXI/A151). Vitamin D modifies IFN responses in both lymphocytes and in monocytes (figure 5). Thus, vitamin D, alone or combined with IFN- β therapy, shifts immunity in MS from a proclivity for Th1 and Th17 cytokine production to a Th2 profile, in addition to increasing regulatory CD4 T-cell numbers.²⁵

Relevant to the high monocyte:T-cell ratio in MS plaques,²⁶ IFN- β alone induces IL-10 production by activated T cells, but reduces IL-10 in monocytes.¹⁵ Vitamin D, as well as prostaglandins and cyclic adenosine monophosphate agonists, induces IL-10 in monocytes¹⁵ and is likely to be complementary to IFN therapy. High levels of serum vitamin D predicted stronger in vitro vitamin D plus IFN- β responses (figure 4). Combining vitamin D with IFNs to reverse the Th1/Th2 effect could have therapeutic benefit in MS and other autoimmune diseases.

Our in vitro data showing that vitamin D enhances IFN- β effects in most forms of MS (figure 4) provide an explanation for clinical observations. High serum vitamin D is linked to preventing the onset of MS, and enhancing the benefit of IFN- β therapy on MS relapses, accumulation of disability, and MRI lesions.⁴ Oral vitamin D supplementation in a prospective trial prevented a second episode of demyelination after optic neuritis.²⁷ Vitamin D supplementation during IFN- β therapy had MRI and clinical benefits.^{5–7}

Serum 25-OH vitamin D levels in MS were much lower before widespread vitamin D supplementation. Vitamin D levels averaged 12 ng/mL in Finnish women in 1983, but rose to 28 by 2009.²⁸ Serum vitamin D levels were 17–22 ng/mL in nonsupplementing Eastern Europe, but 26–30 in North America in 2004 in the BEYOND study of double-dose IFN- β -1b.²⁹ Vitamin D levels in our patients with MS are now above 30 ng/mL (table). The effect of enhanced IFN- β signaling by vitamin D in untreated and IFN-treated patients is stronger in RRMS-s than in SPMS, a form of MS with intrinsic low IFN responses.^{11,12} Low serum vitamin D levels and cohorts with more clinically advanced MS may have reduced the efficacy of IFN- β in early pivotal trials.

During exacerbations in untreated MS, vitamin D enhanced IFN- β effects to a degree intermediate between relapsing and progressive forms of MS (figure 3). Enhancement was greater when serum vitamin D levels were low (figure 4). This augmentation suggests that adequate serum vitamin D would help overcome the transient IFN resistance seen during exacerbations.^{11,12} Variability in response to vitamin D plus IFN- β is highest in untreated stable MS, suggesting that some patients with “stable” MS are in a danger zone with greater likelihood of having attacks.

During glatiramer treatment, in vitro vitamin D plus IFN- β enhanced p-Y-STAT1 activation in RRMS-s and SPMS and in HC (figure 3). The combination reduced IL-17 and caused a shift from a Th1 to a Th2 profile, more than in IFN- β -treated patients. Nonetheless, higher serum vitamin D levels have less clinical benefit during glatiramer treatment than during IFN- β therapy.²⁹ Vitamin D and glatiramer both bias immunity toward Th2 responses,^{30–32} so adding vitamin D to glatiramer may not have much additional effect. The Th2 shift may also provide a basis for the benefits of high serum vitamin D levels on natalizumab therapy.³³

Vitamin D has an opposite effect on Th1 vs Th2 family cytokines compared with IFN- β . This indicates that there are molecular effects in addition to activation of p-Y-STAT1. Vitamin D affects inflammatory responses, oxidative stress, cell adhesion, cellular growth, and DNA repair in multiple cell types. Three percent of all genes are controlled in some way by vitamin D.³⁴ Vitamin D regulates 229 genes in immortalized B cells and additional genes in other tissues.³⁵ The vitamin D receptor binds 2,776 genomic sites; these are overrepresented for active transcription and enhancement of

gene expression. Some of these are genes identified in MS genome-wide association studies, and many are related to IFN signaling (JAK1, Tyk2, STAT3, IRF8, and RGS1).³⁶ In the BENEFIT study of IFN- β -1b in clinically isolated syndromes, 63 vitamin D-regulated genes affected Gd+ MRI lesions; 62 of these were also regulated by IFN- β .³⁷ The molecular basis for these interactions is under investigation.

There were several limitations of this study. There were only modest numbers of patients with relapses. Nonetheless, responses to IFN- β in these patients were intermediate between stable RRMS and progressive MS, paralleling our studies of serum IFN levels and IFN signaling.¹¹

Second, cells were activated in vitro because resting MS MNCs generated only low levels of cytokines. Dysregulated immunity is a hallmark of MS and reflects CNS inflammation. Activation of a small number of cells can provoke significant autoimmune disease, and shifts in levels of cytokines can have significant effects on immunity.^{38,39} Specific brain antigens (myelin basic protein, myelin-oligodendrocyte glycoprotein, and proteolipid protein), tetanus, and viral antigens have been used to activate MS cells. They are not the causative agents in MS, but do induce weak cytokine responses in MS and in some HCs. As a model of acute and chronic low-level inflammation in MS, ConA was used to activate MNCs. This mannose-binding lectin broadly activates monocytes and CD4 and CD8 T cells, including the CD8⁺ CD28⁻ regulatory population that is subnormal in MS.^{40,41} This subset is not stimulated by commonly used anti-CD3 plus anti-CD28 antibodies, and ConA is thus a more appropriate activator for MS immune cells.

IFN- β is infrequently used in PPMS because it only minimally slows the worsening ambulatory dysfunction. Nonetheless, patients with PPMS had strong responses to IFN- β -1b plus vitamin D induction relative to IFN- β -1b alone (figure 2). It is possible that vitamin D could potentiate benefits of IFN- β in PMS, such as reducing breakthrough relapses and maintaining cognition.⁴²

In conclusion, vitamin D induces a shift from Th1 to Th2 immunity in activated MNCs. When combined with IFN- β , it enhances activation of STAT1, a transcription factor that controls IFN signaling, and also reverses most of the Th1-inducing effects seen with IFN- β alone. This provides an explanation for benefits of vitamin D in preventing and ameliorating different forms of MS before and during IFN- β therapy. Vitamin D is an inexpensive and safe addition to amplify the therapeutic benefit of IFN- β .

Author contributions

X. Feng: data acquisition, drafting/revising the manuscript, study concept or design, analysis or interpretation of data, statistical analysis, study supervision, and approval of data. Z. Wang: data acquisition, drafting/revising the manuscript, analysis or interpretation of data, and statistical analysis. Q.

Howlett-Prieto: data acquisition and statistical analysis. N. Einhorn: data acquisition and drafting/revising the manuscript. S. Causevic: data acquisition, study concept or design, analysis or interpretation of data, statistical analysis, and study supervision. A. Reder: data acquisition, drafting/revising the manuscript, study concept or design, analysis or interpretation of data, statistical analysis, study supervision, and obtaining funding.

Study funding

This study was funded by unrestricted grant support from Bayer Healthcare Pharmaceuticals and the US National Multiple Sclerosis Society (NMSS), RG4509A.

Disclosure

X. Feng has received unrestricted grant support from Bayer, Biogen, Roche-Genentech, Merck-Serono, Novartis, and Teva. Z. Wang, Q. Howlett-Prieto, N. Einhorn, and S. Causevic report no disclosures. A.T. Reder has received unrestricted grant support from Bayer, Biogen, Roche-Genentech, Merck-Serono, Novartis, and Teva and is a consultant for Bayer, Biogen, Roche-Genentech, Merck-Serono, and Novartis. Go to Neurology.org/NN for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* April 8, 2019. Accepted in final form August 13, 2019.

References

1. Sintzel MB, Rameeta M, Reder AT. Vitamin D and multiple sclerosis: a comprehensive review. *Neurol Ther* 2018;7:59–85.
2. Orton SM, Wald L, Confavreux C, et al. Association of UV radiation with multiple sclerosis prevalence and sex ratio in France. *Neurology* 2011;76:425–431.
3. Munger KL, Levin LI, Hollis BW, Howard NS, Ascherio A. Serum 25-hydroxyvitamin D levels and risk of multiple sclerosis. *JAMA* 2006;296:2832–2838.
4. Ascherio A, Munger KL, White R, et al. Vitamin D as an early predictor of multiple sclerosis activity and progression. *JAMA Neurol* 2014;71:306–314.
5. Soilu-Hänninen M, Aivo J, Lindström BM, et al. A randomised, double blind, placebo controlled trial with vitamin D3 as an add on treatment to interferon β -1b in patients with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2012;83:565–571.
6. Aivo J, Lindström BM, Soilu-Hänninen M. A randomised, double-blind, placebo-controlled trial with vitamin D3 in MS: subgroup analysis of patients with baseline disease activity despite interferon treatment. *Mult Scler Int* 2012;2012:802796.
7. Stewart N, Simpson S, van der Mei I, et al. Interferon- β and serum 25-hydroxyvitamin D interact to modulate relapse risk in MS. *Neurology* 2012;79:254–260.
8. Nwosu BU, Maranda L, Berry R, et al. The vitamin D status of prison inmates. *PLoS One* 2014;9:e90623.
9. Cree BAC, Al-Sabbagh A, Bennett R, Goodin D. Response to interferon beta-1a treatment in African American multiple sclerosis patients. *Arch Neurol* 2005;62:1681–1683.
10. Reder AT, Antel JP, Oger J, McFarland TA, Rosenkoetter M, Arnason BGW. Low T8 antigen density on lymphocytes in active multiple sclerosis. *Ann Neurol* 1984;16:242–249.
11. Feng X, Han D, Kilaru BK, Franek BS, Niewold TB, Reder AT. Inhibition of interferon-beta responses in multiple sclerosis immune cells associated with high-dose statins. *Arch Neurol* 2012;69:1303–1309.
12. Feng X, Petraglia AL, Chen M, Byskosh PV, Boos MD, Reder AT. Low expression of interferon-stimulated genes in active multiple sclerosis is linked to subnormal phosphorylation of STAT1. *J Neuroimmunol* 2002;129:205–215.
13. Pachner AR, DiSano K, Royce DB. Clinical utility of a molecular signature in inflammatory demyelinating disease. *Neurol Neuroimmunol Neuroinflamm* 2018;6:e520.
14. Feng X, Wang Z, Causevic S, et al. Vitamin D enhances interferon-beta response in multiple sclerosis. *Neurology* 2016;86:P3.103.
15. Feng X, Yau D, Holbrook C, Reder AT. Type I interferons inhibit interleukin-10 production in activated human monocytes and stimulate IL-10 in T cells: implications for Th1-mediated diseases. *J Interferon Cytokine Res* 2002;22:311–319.
16. Rudick RA, Ransohoff RM. Cytokine secretion by multiple sclerosis monocytes. Relationship to disease activity. *Arch Neurol* 1992;49:265–270.
17. Boutros T, Croze E, Yong VW. Interferon- β is a potent promoter of nerve growth factor production by astrocytes. *J Neurochem* 1997;69:939–946.

18. Lee LF, Axtell R, Tu GH, et al. IL-7 promotes T(H)1 development and serum IL-7 predicts clinical response to interferon- β in multiple sclerosis. *Sci Transl Med* 2011;3:93ra68.
19. Yamaguchi KD, Ruderman DL, Croze E, et al. IFN-beta-regulated genes show abnormal expression in therapy-naïve relapsing-remitting MS mononuclear cells: gene expression analysis employing all reported protein-protein interactions. *J Neuroimmunol* 2008;195:116–120.
20. Correale J, Villa A. Isolation and characterization of CD8+ regulatory T cells in multiple sclerosis. *J Neuroimmunol* 2008;195:121–134.
21. Denic A, Woodla B, Rodriguez M. CD8(+) T cells in multiple sclerosis. *Expert Opin Ther Targets* 2013;17:1053–1066.
22. Noronha A, Toscas A, Jensen MA. Contrasting effects of alpha, beta, and gamma interferons on nonspecific suppressor function in multiple sclerosis. *Ann Neurol* 1992;31:103–106.
23. Byskosh PV, Reder AT. Interferon beta-1b effects on cytokine mRNA in peripheral mononuclear cells in multiple sclerosis. *Mult Scler* 1996;1:262–269.
24. Tough DF. Modulation of T-cell function by type I interferon. *Immunol Cell Biol* 2012;90:492–497.
25. Royal W, Mia Y, Li H, Naunton K. Peripheral blood regulatory T cell measurements correlate with serum vitamin D levels in patients with multiple sclerosis. *J Neuroimmunol* 2009;213:135–141.
26. Lassmann H, Suchanek G, Ozawa K. Histopathology and the blood-cerebrospinal fluid barrier in multiple sclerosis. *Ann Neurol* 1994;36:S42–S46.
27. Derakhshandi H, Etemadifar M, Feizi A, et al. Preventive effect of vitamin D3 supplementation on conversion of optic neuritis to clinically definite multiple sclerosis: a double blind, randomized, placebo-controlled pilot clinical trial. *Acta Neurol Belg* 2013;113:257–263.
28. Munger KL, Hongell K, Åivo J, Soilu-Hänninen M, Surcel H-M, Ascherio A. 25-hydroxyvitamin D deficiency and risk of MS among women in the Finnish Maternity Cohort. *Neurology* 2017;89:1578–1583.
29. Fitzgerald KC, Munger KL, Köchert K, et al. Association of vitamin D levels with multiple sclerosis activity and progression in patients receiving interferon beta-1b. *JAMA Neurol* 2015;72:1458–1465.
30. Gröber U, Spitz J, Reichrath J, Kisters K, Holick MF. Vitamin D: update 2013: from rickets prophylaxis to general preventive healthcare. *Dermatoendocrinol* 2013;5:331–347.
31. Dhib-Jalbut S, Chen M, Said A, Zhan M, Johnson KP, Martin R. Glatiramer acetate-reactive peripheral blood mononuclear cells respond to multiple myelin antigens with a Th2-biased phenotype. *J Neuroimmunol* 2003;140:163–171.
32. Valenzuela RM, Kaufman M, Balashov KE, Ito K, Buyske S, Dhib-Jalbut S. Predictive cytokine biomarkers of clinical response to glatiramer acetate therapy in multiple sclerosis. *J Neuroimmunol* 2016;300:59–65.
33. Scott TF, Hackett CT, Dworek DC, Schramke CJ. Low vitamin D level is associated with higher relapse rate in natalizumab treated MS patients. *J Neurol Sci* 2013;330:27–31.
34. Bouillon R, Carmeliet G, Verlinden L, et al. Vitamin D and human health: lessons from vitamin D receptor null mice. *Endocr Rev* 2008;29:726–776.
35. Ramagopalan SV, Hanwell HE, Giovannoni G, et al. Vitamin D-dependent rickets, HLA-DRB1, and the risk of multiple sclerosis. *Arch Neurol* 2010;67:1034–1035.
36. Berlanga-Taylor AJ, Disanto G, Ebers GC, Ramagopalan SV. Vitamin D-gene interactions in multiple sclerosis. *J Neurol Sci* 2011;311:32–36.
37. Munger KL, Köchert K, Simon KC, et al. Molecular mechanism underlying the impact of vitamin D on disease activity of MS. *Ann Clin Transl Neurol* 2014;1:605–617.
38. Genç K, Dona DL, Reder AT. Increased CD80(+) B cells in active multiple sclerosis and reversal by interferon beta-1b therapy. *J Clin Invest* 1997;99:2664–2671.
39. Germain RN. The art of the probable: system control in the adaptive immune system. *Science* 2001;293:240–245.
40. Antel JP, Arnason BG, Medof ME. Suppressor cell function in multiple sclerosis: correlation with clinical disease activity. *Ann Neurol* 1979;5:338–342.
41. Karaszewski JW, Reder AT, Anlar B, Kim WC, Arnason BG. Increased lymphocyte beta-adrenergic receptor density in progressive multiple sclerosis is specific for the CD8+, CD28– suppressor cell. *Ann Neurol* 1991;30:42–47.
42. Lacy M, Hauser M, Pliskin N, Assuras S, Valentine MO, Reder A. The effects of long-term interferon-beta-1b treatment on cognitive functioning in multiple sclerosis: a 16-year longitudinal study. *Mult Scler* 2013;19:1765–1772.

Neurology[®] Neuroimmunology & Neuroinflammation

Vitamin D enhances responses to interferon- β in MS

Xuan Feng, Zhe Wang, Quentin Howlett-Prieto, et al.

Neurol Neuroimmunol Neuroinflamm 2019;6:e622

DOI 10.1212/NXI.0000000000000622

This information is current as of October 3, 2019

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/6/6/e622.full.html
References	This article cites 42 articles, 3 of which you can access for free at: http://nn.neurology.org/content/6/6/e622.full.html##ref-list-1
Citations	This article has been cited by 3 HighWire-hosted articles: http://nn.neurology.org/content/6/6/e622.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Demyelinating disease (CNS) http://nn.neurology.org/cgi/collection/all_demyelinating_disease_cns Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases Endocrine http://nn.neurology.org/cgi/collection/endocrine Multiple sclerosis http://nn.neurology.org/cgi/collection/multiple_sclerosis Nutritional http://nn.neurology.org/cgi/collection/nutritional
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.