Fatal PCR-negative herpes simplex virus-1 encephalitis with GABA_A receptor antibodies

Simon Schuster, MD, Ligia Abrante, Jakob Matschke, MD, Marc Lütgehetmann, MD, Brigitte Holst, MD, Mathias Gelderblom, MD, Hanna Braass, MD, Frank Leypoldt, MD,* and Tim Magnus, MD*

Correspondence
Dr. Schuster
s.schuster@uke.de

Neurol Neuroimmunol Neuroinflamm 2019;6:e624. doi:10.1212/NXI.0000000000000624

Herpes simplex virus-1 encephalitis (HSVE) is the most common cause of fatal sporadic encephalitis worldwide. Recently, post-HSVE relapses due to autoimmune encephalitis with NMDA receptor antibodies and other synaptic autoantibodies have been reported in up to 27% of patients within 1–2 months after HSVE.^{1,2} We report an unusual case of a patient with refractory status epilepticus due to HSVE with antibodies against GABA_A receptors (GABA_AR).

Case report

A 47-year-old man was admitted with a new-onset generalized seizure. Two days before admission, behavioral changes were observed. No psychiatric symptoms, headaches, or feverish infections had occurred in the previous weeks. He had a B-cell non-Hodgkin lymphoma 5 years ago and a Hodgkin lymphoma 16 years ago, both in complete remission after radio-chemotherapy. Eventually, focal status epilepticus developed provoking intensified anticonvulsive treatment.

Initial brain MRI showed hyperintense fluid-attenuated inversion recovery (FLAIR) signal of the left prefrontal gyrus without diffusion restriction (figure 1, A and D) or contrast enhancement. CSF analysis on the day of admission revealed lymphocytic pleocytosis (13 leukocytes/ μ L), elevated protein (574 mg/L), and lactate (2.74 mmol/L). IV acyclovir 750 mg TID was started. CSF herpes simplex virus-1 (HSV-1) quantitative polymerase chain reaction (qPCR) was negative. A second lumbar puncture (day 5 of admission) showed 29 lymphocytes/ μ L, normal protein, and slightly elevated lactate (2.27 mmol/L). HSV-1 DNA remained undetectable. Immunoglobulin G-HSV antibody index was unremarkable (<1.3); further analysis revealed no other infectious causes. However, GABA_AR antibodies were detected in serum (1:1,600) and CSF (1:32) of the second lumbar puncture using cell-based assays, tissue-based assays, and life embryonal hippocampal neuron cultures (figure 1, G–I)³; no other neuronal antibodies were identified.

Acyclovir was stopped, yet IV methylprednisolone did not induce clinical improvement. Follow-up MRI showed expansion of the left frontal hyperintense FLAIR lesion with accompanied diffusion restriction and new bilateral opercular diffusion restrictions (figure 1, B, C, E, F). Refractory status epilepticus continued (EEG, figure e-1, [links.lww.com/NXI/A145](https://www.lww.com/NXI/A145)). The patient died of bowel ischemia due to thrombosis of the mesenteric artery. Postmortem revealed extensive HSVE with necrosis, inflammation, positive HSV antigen, and tissue PCR (figure 1J). No evidence of lymphoma was found.

*These authors contributed equally to this manuscript.

From the Department of Neurology (S.S., M.G., H.B., T.M.), University Hospital Hamburg-Eppendorf; Neuroimmunology (L.A., F.L.), Institute of Clinical Chemistry, University Hospital Schleswig-Holstein, Kiel/Lübeck; Institute of Neuropathology (J.M.), University Hospital Hamburg-Eppendorf; Department of Medical Microbiology, Virology and Hygiene (M.L.), University Hospital Hamburg-Eppendorf; Department of Neuroradiology (B.H.), University Hospital Hamburg-Eppendorf; and Department of Neurology (F.L.), University Hospital Schleswig-Holstein, Kiel.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

MRI 1st day of admission (A and D) and 14 days after admission (B, C, E, and F), showing progression of the left frontal hyperintense lesion and new diffusion restriction left frontally and bilaterally in the operculum on day 14. (A–C) Axial fluid-attenuated inversion recovery-weighted images with hyperintense lesion of the left prefrontal gyrus (A and B) and the operculum bilaterally (C). (D–F) Diffusion-weighted images and apparent diffusion coefficient images (small insets) without diffusion restriction (D) and marked diffusion restriction in the left prefrontal gyrus (E) and the operculum bilaterally (F). (G) Immunolabeling of sagittal rat brain sections with the patient's CSF antibodies showing a characteristic pattern. Patient and control CSF 1:4. Anti-human IgG (H + L). Human IgM and IgA did not show immunoreactivity. Scale bar 1 mm. (H) Detection of antibodies to the GABA_A receptors (GABA_AR) using HEK293 cell-based assay. Patient's but not control serum detects GABA_AR. Human GABA_AR subunits transfected into HEK293 cells and stained via life cell staining (serum 1:40). Green human IgG, red commercial GABA_AR antibody. Scale bar 20 μ m. (I) Patient's but not control serum detects neuronal surface antigens. Non-permeabilized embryonic rat hippocampal neuron cultures DIV21 life cell stained with human IgG and nuclear counterstaining with DAPI (blue). Scale bar 20 μ m. (J) Postmortem herpes simplex virus antigen staining of the patient's hippocampus. Scale bar 5 mm. DAPI = 4',6-diamidino-2-phenylindole; IgG = immunoglobulin G.

Discussion

We describe an unusual case of CSF-qPCR-negative HSVE with concomitant GABA_AR antibodies. We confirmed presence and specificity of GABA_AR antibodies in serum and CSF

with high titers, typical staining on rat brain immunohistochemistry and neuronal synapses of live neurons in vitro.

Our patient was initially misdiagnosed with idiopathic GABA_AR encephalitis owing to detection of GABA_AR antibodies, 2

negative HSV-1 qPCR in CSF, and characteristic clinical presentation with severe encephalitis and refractory status epilepticus.^{3,4} HSVE was only diagnosed postmortem by demonstration of widespread viral replication in brain tissue. Coincidental development of HSVE and GABA_AR encephalitis is unlikely because of the low incidence of both diseases; rather breakdown of immunologic tolerance toward GABA_AR likely provoked by virus-induced destruction of neurons would be a plausible explanation.⁵

Previous post-HSVE autoimmune encephalitis cases predominantly had a biphasic course. However, development in contiguity with HSVE symptoms similar to our case has been described in adults,¹ and relapses have been observed as early as 7 days after HSVE in a 2-month-old boy.⁵

Furthermore, a case of post-HSVE GABA_AR encephalitis was recently described in a 15-month-old child occurring 8 weeks after herpes infection, and a second case occurred following HHV6 encephalitis.⁴ We are not aware of a case of post-HSVE GABA_AR encephalitis in an adult patient. However, because of the unavailability of initial CSF antibody testing, we cannot exclude the presence of “premorbid” GABA_AR antibodies as a coincidental finding related to previous lymphoma. Although, the “premorbid” presence of high-titer CSF GABA_AR antibodies without a history of seizures appears less plausible than our current hypothesis of continuous post-HSVE GABA_AR encephalitis.

Pathologic-proven HSVE without detectable HSV-1 DNA in CSF is another unusual feature and was rarely reported.^{6,7} Hypothetical explanations could be early lumbar puncture and analysis under pretreatment with acyclovir. The previous lymphoma could have predisposed toward (1) the development of HSVE as has been noted in inborn errors of pattern-recognition pathways and (2) a breakdown of immune tolerance and consecutively development of systemic GABA_AR antibodies predisposing to rapid development of CNS autoimmunity after HSVE.

In summary, this single case illustrates the occurrence of GABA_AR antibodies in PCR-negative HSVE in an immunosuppressed patient resulting in misdiagnosis of idiopathic GABA_AR encephalitis. Key clinical points are (1) the difficulty of ruling out HSVE with qPCR and immunoglobulin G antibody index, (2) the occurrence of GABA_AR antibodies in parallel to HSVE, and (3) the possibility of unusual clinical courses of HSVE and post-HSVE autoimmune encephalitis in patients with immune deficiencies.

Study funding

No targeted funding reported.

Disclosure

S. Schuster, L. Abrante, and J. Matschke report no conflict of interests. M. Lütgehetmann reports receiving speaker honoraria from Roche Diagnostics, DiaSorin, Swisslab, Abbott, and

BioMerieux. B. Holst reports no conflict of interests. M. Gelderblom reports receiving personal fees from Allergan and Merz. H. Braass reports funding for travel from Boehringer Ingelheim. F. Leypoldt reports receiving speaker honoraria from Grifols, Teva, Fresenius, Roche, Merck, Bayer, Sanofi, and Novartis. He serves on a local advisory board for Roche and Biogen and works for an academic institution participating in commercial antibody testing. He does not receive personal reimbursements from this antibody testing. T. Magnus reports receiving personal fees from Merck Serono, Boehringer Ingelheim, Bristol-Myers Squibb/Pfizer, Novartis, Biogen, Grifols, and CSL-Behring and grants from the European Union (FP-7, ERA-NET), German Research Foundation, Schilling Foundation, and Werner-Otto Foundation. Go to Neurology.org/NN for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* April 29, 2019. Accepted in final form August 6, 2019.

Appendix Authors

Name	Location	Role	Contribution
Simon Schuster, MD	University Hospital Hamburg-Eppendorf, Department of Neurology	Author	Study design, collecting the data, analysis and interpretation of data, and drafting and revising the manuscript
Ligia Abrante	University Hospital Schleswig-Holstein, Kiel/Lübeck, Neuroimmunology, Institute of Clinical Chemistry	Author	Analysis and interpretation of data and revising the manuscript
Jakob Matschke, MD	University Hospital Hamburg-Eppendorf, Institute of Neuropathology	Author	Analysis and interpretation of data and revising the manuscript
Marc Lütgehetmann, MD	University Hospital Hamburg-Eppendorf, Department of Medical Microbiology, Virology and Hygiene	Author	Analysis and interpretation of data and revising the manuscript
Brigitte Holst, MD	University Hospital Hamburg-Eppendorf, Department of Neuroradiology	Author	Analysis and interpretation of data and revising the manuscript
Mathias Gelderblom, MD	University Hospital Hamburg-Eppendorf, Department of Neurology	Author	Analysis and interpretation of data and revising the manuscript
Hanna Braass, MD	University Hospital Hamburg-Eppendorf, Department of Neurology	Author	Collecting the data, analysis and interpretation of data, and revising the manuscript

Continued

Appendix *(continued)*

Name	Location	Role	Contribution
Frank Leypoldt, MD	University Hospital Schleswig-Holstein, Kiel/Lübeck, Neuroimmunology, Institute of Clinical Chemistry, and Department of Neurology	Author	Study design, analysis and interpretation of data, and drafting and revising the manuscript
Tim Magnus, MD	University Hospital Hamburg-Eppendorf, Department of Neurology	Author	Interpretation of data and revising the manuscript

References

1. Armangue T, Moris G, Cantarin-Extremera V, et al. Autoimmune post-herpes simplex encephalitis of adults and teenagers. *Neurology* 2015;85:1736–1743.
2. Armangue T, Spatola M, Vlagea A, et al. Frequency, symptoms, risk factors, and outcomes of autoimmune encephalitis after herpes simplex encephalitis: a prospective observational study and retrospective analysis. *Lancet Neurol* 2018;17:760–772.
3. Petit-Pedrol M, Armangue T, Peng X, et al. Encephalitis with refractory seizures, status epilepticus, and antibodies to the GABAA receptor: a case series, characterisation of the antigen, and analysis of the effects of antibodies. *Lancet Neurol* 2014;13: 276–286.
4. Spatola M, Petit-Pedrol M, Simabukuro MM, et al. Investigations in GABAA receptor antibody-associated encephalitis. *Neurology* 2017;88:1012–1020.
5. Armangue T, Leypoldt F, Málaga I, et al. Herpes simplex virus encephalitis is a trigger of brain autoimmunity. *Ann Neurol* 2014;75:317–323.
6. Adler AC, Kadimi S, Apaloo C, Marcu C. Herpes simplex encephalitis with two false-negative cerebrospinal fluid PCR tests and review of negative PCR results in the clinical setting. *Case Rep Neurol* 2011;3:172–178.
7. Mendez AA, Bosco A, Abdel-Wahed L, Palmer K, Jones KA, Killoran A. A fatal case of herpes simplex encephalitis with two false-negative polymerase chain reactions. *Case Rep Neurol* 2018;10:217–222.

Neurology® Neuroimmunology & Neuroinflammation

Fatal PCR-negative herpes simplex virus-1 encephalitis with GABA_A receptor antibodies

Simon Schuster, Ligia Abrante, Jakob Matschke, et al.
Neurol Neuroimmunol Neuroinflamm 2019;6;
DOI 10.1212/NXI.0000000000000624

This information is current as of September 25, 2019

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services

including high resolution figures, can be found at:
<http://nn.neurology.org/content/6/6/e624.full.html>

References

This article cites 7 articles, 0 of which you can access for free at:
<http://nn.neurology.org/content/6/6/e624.full.html##ref-list-1>

Citations

This article has been cited by 1 HighWire-hosted articles:
<http://nn.neurology.org/content/6/6/e624.full.html##otherarticles>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Autoimmune diseases

http://nn.neurology.org/cgi/collection/autoimmune_diseases

Encephalitis

<http://nn.neurology.org/cgi/collection/encephalitis>

Post-infectious

http://nn.neurology.org/cgi/collection/postinfectious_

Status epilepticus

http://nn.neurology.org/cgi/collection/status_epilepticus

Viral infections

http://nn.neurology.org/cgi/collection/viral_infections

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://nn.neurology.org/misc/about.xhtml#permissions>

Reprints

Information about ordering reprints can be found online:

<http://nn.neurology.org/misc/addir.xhtml#reprintsus>

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.