Paraneoplastic cerebellar ataxia and antibodies to metabotropic glutamate receptor 2

Raquel Ruiz-García, PhD, Eugenia Martínez-Hernández, MD, PhD, Bastien Joubert, MD, Mar Petit-Pedrol, PhD, Elena Pajarón-Boix, MD, Verónica Fernández, MD, Lidia Salais, MD, María del Pozo, MD, Thais Armangué, MD, PhD, Lidia Sabater, PhD, Josep Dalmau, MD, PhD, and Francesc Graus, MD, PhD

Correspondence

Dr. Graus
francesc.graus@idibaps.org

Neurol Neuroimmunol Neuroinflamm 2020;7:e658. doi:10.1212/NXI.0000000000000658

Abstract

Objective

To report the presence of a new neuronal surface antibody against the metabotropic glutamate receptor 2 antibody (mGluR2-Ab) in 2 patients with paraneoplastic cerebellar ataxia.

Methods

mGluR2-Abs were initially characterized by immunohistochemistry on the rat brain and confirmed by immunofluorescence on HEK293 cells transfected with mGluR2. Additional studies included analysis of potential cross-reactivity with other mGluRs, expression of mGluR2 in patients' tumors, and the effects of mGluR2-Abs on cultures of rat hippocampal neurons.

Results

Patient 1 was a 78-year-old woman with progressive cerebellar ataxia with an initial relapsing-remitting course who developed a small-cell tumor of unknown origin. Patient 2 was a 3-year-old girl who presented a steroid-responsive acute cerebellitis preceding the diagnosis of an alveolar rhabdomyosarcoma. Patients' serum and CSF showed a characteristic immunostaining of the hippocampus and cerebellum in rat brain sections and immunolabeled the cell surface of live rat hippocampal neurons. HEK293 cells transfected with mGluR1, 2, 3, and 5 confirmed that patients' antibodies only recognized mGluR2. mGluR2-Abs were not detected in 160 controls, 120 with paraneoplastic, autoimmune, or degenerative ataxias, and 40 with autoimmune encephalitis and antibodies against mGluR5 or unknown antigens. Expression of mGluR2 in tumors was confirmed by immunohistochemistry using a commercial mGluR2-Ab. Incubation of live rat hippocampal neurons with CSF of patient 2 did not modify the density of surface mGluR2 clusters.

Conclusions

mGluR2-Abs are a novel biomarker of paraneoplastic cerebellar ataxia. The potential pathogenic effect of the antibodies is not mediated by downregulation or internalization of neuronal surface mGluR2.

From the Immunology Department (R.R.-G.), Centre Diagnòstic Biomèdic, Hospital Clínic; Neuroimmunology Program (E.M.-H., B.J., M.P.-P., T.A., L. Sabater, J.D., F.G.), Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS); Service of Neurology (E.M.-H.), Hospital Clínic; Service of Neurology (E.P.-B., L. Salais), Hospital General de Castellón, Spain; Service of Pediatrics (V.F.), Hospital General de Albacete, Spain; Onco-Hematology Unit (M.d.P.), Service of Pediatrics, Hospital General de Albacete; Institutió Catalana de Recerca i Estudis Avançats (ICREA) (J.D.), Barcelona, Spain; and Department of Neurology (J.D.), University of Pennsylvania, Philadelphia, PA.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

CBA = cell-based assay; mGluR2-Ab = metabotropic glutamate receptor 2 antibody.

In autoimmune encephalitis, there are many autoantibodies against neuronal cell surface or synaptic receptors that associate with a good response to immunotherapy.¹ However, in autoimmune cerebellar ataxias, the occurrence of similar autoantibodies is rare. Only 2 antibodies against neuronal surface antigens, and therefore potentially pathogenic, have been consistently reported: the antibodies against voltage-gated calcium channels in patients with small-cell lung cancer and the antibodies against the metabotropic glutamate receptor (mGluR) 1 in patients with autoimmune ataxia.² Here, we report 2 patients with cerebellar ataxia and cancer who presented mGluR2 antibodies (mGluR2-Abs).

Methods

Immunologic studies

Serum and CSF from the 2 patients were examined by immunohistochemistry on frozen sections of the rat brain and by indirect immunofluorescence on cell-based assays (CBAs) using unfixed HEK293 cells transfected with plasmids encoding mGluR1, 2, 3, and 5 as reported.^{3,4} In addition, serum or CSF from 160 patients, 120 with cerebellar ataxias (paraneoplastic: 58; pediatric cerebellitis: 32; degenerative: 20; with mGluR1 antibodies: 10), 10 patients with autoimmune encephalitis with mGluR5 antibodies, and 30 patients with autoimmune encephalitis and antibodies against unknown antigens were tested with CBA of mGluR2. mGluR2-Ab effects were examined in cultures of rat hippocampal neurons. CSF (dilution 1:10 in culture medium) of patient 2 or control CSF was added to the culture media for 48 hours. After removing the media, neurons were sequentially incubated with mGluR2-Ab-positive CSF (1:50) for 1 hour at 37°C, followed by a secondary anti-human Immunoglobulin G (IgG) Alexa Fluor 488 (Invitrogen A11013, 1:1,000) and the mGluR2 clusters quantified as previously described.⁵

To demonstrate the expression of mGluR2 in the tumor, paraffin sections were deparaffinized, incubated with a commercial mGluR2-Ab (Cell Signaling Technology), and developed with the avidin-biotin technique (Vector Labs). Specificity of the tumor immunoreactivity was confirmed by adsorption of the antibody with HEK293 cells transfected with mGluR2 or with plasmids without insert.

Standard protocol approvals, registrations, and patient consents

The Ethic Committee of the Hospital Clinic approved the study. All patients or proxies gave written informed consent for the storage and use samples and clinical information for research purposes.

Data availability

All data are reported within the article and available anonymized by request from qualified investigators.

Results

Patient 1

A 78-year-old woman presented with a 3-year history of cerebellar ataxia. During the first 2 years, she had relapsing episodes of gait instability and dysarthria that spontaneously improved in a few days. In the last year, she developed progressive ataxia that did not respond to IV immunoglobulins or corticosteroids and only partially to 1 cycle of rituximab. The brain MRI showed focal hyperintense cerebellar lesions in T2-weighted images that later evolved to diffuse involvement of the cerebellar white matter (figure 1). Three years after symptom onset, a right inguinal adenopathy was identified, which biopsy demonstrated a small-cell neuroendocrine cancer of unknown origin. Despite palliative chemotherapy, the patient died a few months after tumor diagnosis.

Patient 2

In September 2018, a 3-year-old girl presented with a 3-day history of fever, nausea, and vomiting, followed by somnolence and gait instability. Neurologic examination disclosed irritability, dysarthria, horizontal right-beating nystagmus, limb and truncal ataxia, and broad-based gait requiring bilateral support. The CSF was normal, and brain MRI showed patchy gadolinium enhancement in the cerebellar folia. She was treated with IV methylprednisolone and immunoglobulins with full recovery. In January 2019, she was diagnosed with alveolar rhabdomyosarcoma of the right axillary region with positive regional lymph nodes but without distant metastasis. The patient did not present a relapse of the cerebellar ataxia.

Immunologic findings

Serum and CSF of the 2 patients showed an identical staining of the cerebellum and hippocampus and immunolabeled the cell surface of live rat hippocampal neurons. In the cerebellum, the staining was limited to the granular cell layer with robust immunoreactivity of neuronal processes along with dendrites from cells immediately below the Purkinje cells extending into the molecular layer. In the hippocampus, there was a strong staining of the stratum moleculare with the exception of its innermost part, the stratum lacunosum moleculare, the stratum pyramidale of CA3, and CA4, and the hilus (figure 2, A–C). This immunohistochemical pattern had not been previously seen during the screening of more than 4,000 samples that we receive annually for the detection of neuronal antibodies. In contrast, we found that the immunostaining was very similar to that shown by a commercial mGluR2-Ab (figure 2, D and E).⁶ Serum and CSF of the 2 patients were reactive with HEK293 cells expressing mGluR2 but not with cells expressing mGluR1, 3, or 5 (figure 2, F–I). IgG subclass analysis showed that IgG1 was the predominant type in the 2

Figure 1 Brain MRI findings in patient 1

Brain MRI obtained in 2016 showed multiple FLAIR hyperintensities that did not enhance with gadolinium and involved the inferior part of the cerebellar hemispheres (A). A follow-up MRI in 2018 showed a more diffuse FLAIR hyperintensity in the white matter of the cerebellar hemispheres (B).

patients. None of the 160 controls were positive in the CBA of mGluR2.

Expression of mGluR2 was observed in both tumors as a granular immunoreactivity that was probably located in the membrane and cytoplasm (figure 3, B and D). This reactivity was not observed when the commercial mGluR2-Ab was preadsorbed with HEK293 cells transfected with mGluR2 (figure 3, A and C). Incubation of rat hippocampal neurons

with patients' CSF for 48 hours did not result in a change of surface density of mGluR2 clusters (data not shown).

Discussion

This study characterizes a novel neuronal surface antibody against mGluR2 in patients with cerebral ataxia and cancer. mGluR2-Abs probably recognize epitopes in the

Figure 2 Reactivity of patients' antibodies with rat brain and HEK293 cells transfected with metabotropic glutamate receptors (mGluR)

Sagittal section of rat brain immunolabeled with CSF of patient 1 (A and B). Note the intense staining of the molecular layer (Mo), stratum lacunosum molecular (SL), and hilus (Hi), as well as the stratum pyramidalis of CA3 and CA4 of the rat hippocampus. In the paraformaldehyde-perfused rat cerebellum, CSF of patient 1 shows reactivity with granular layer (Gr) neurons and dendrites extending into the molecular layer (Mo) (C). Identical immunoreactivity with the hippocampus and cerebellum is observed with a commercial mGluR2-Ab (D and E). Immunostaining of live HEK293 cells expressing mGluR2 with serum from patient 2 (P2) or a healthy control (HC) is shown in panels F and G. P2 serum intensely labels mGluR2-expressing cells (in red; F), whereas the HC serum is negative (G). P2 serum did not react with HEK293 cells transfected with mGluR1, mGluR3, or mGluR5, all 3 tagged with green fluorescent protein (GFP) (H–J).

Figure 3 Tumor expression of mGluR2

mGluR2 expression in tumors of patients 1 (A and B) and 2 (C and D). A commercial mGluR2-Ab shows a granular reactivity with the cytoplasm and probably the membrane of tumor cells as demonstrated in detail in the insets (A and C). This reactivity was abrogated when the antibody was preadsorbed with HEK293 cells transfected with mGluR2 (B and D).

extracellular region of the receptor as supported by the immunolabeling of live nonpermeabilized neuronal cultures. Unlike classical paraneoplastic cerebellar degeneration associated with onconeural antibodies against intracellular antigens, the clinical symptoms of patient 2 responded to immunotherapy suggesting a potential pathogenic role of mGluR2-Abs.

There are 8 subtypes of mGluRs divided in 3 subgroups according to amino acid homologies and interaction with intracellular second messengers. Group I includes mGluR1 and 5, and both are autoimmune targets in cerebellar ataxia and encephalitis associated with Hodgkin lymphoma, respectively.^{5,7} Group II comprises mGluR2 and 3, whereas group III includes the remaining subtypes.⁸ The main physiologic role of mGluR2 is to modulate glutamatergic and GABAergic synaptic transmission.⁸ mGluR2-Abs could potentially alter these functions resulting in cerebellar ataxia. Different from autoantibodies directed against ion channel coupled receptors (e.g., N-methyl D-aspartate receptor), which cause internalization of the corresponding targets, our patients' mGluR2-Ab did not cause reduction of the surface receptor. An alternative possibility is that the antibodies block the function of the channel as has been proposed for other metabotropic receptors such as the gamma-aminobutyric acid b receptor.¹ Future studies should determine whether passive transfer of antibodies to animals causes ataxia, as demonstrated with anti-mGluR1-Ab.⁷ Interestingly, rats lacking mGluR2 display behavioral symptoms but not cerebellar ataxia, suggesting that mGluR2-Ab and genetic deletion of mGluR2 may have different effects. There is precedence for

this; for example, genetic and immunologic models of anti-LGI1 have different phenotypes. Although the genetic abrogation of LGI1 associates with a lethal epileptic phenotype, animals infused with LGI1 antibodies develop severe memory deficits without clinical seizures.⁹

mGluRs are overexpressed in several types of cancer, particularly primary brain tumors, and they seem to be implicated in the regulation of cancer cell proliferation.¹⁰ Our studies show that mGluR2 was expressed by the tumor of both patients. In line with our findings, mGluR2 expression has also been found expressed in lung cancer and rhabdomyosarcoma cell lines.¹⁰

The task for the future is to confirm our findings with additional patients and elucidate the pathogenic role of mGluR2-Ab. However, the current study has practical clinical implications: (1) mGluR2-Abs may occur in patients with autoimmune cerebellar symptoms, (2) detection of these antibodies suggest that the syndrome may respond to immunotherapy, and (3) patients should be examined for the presence of an underlying tumor.

Acknowledgment

The authors thank Mrs. Eva Caballero for her excellent technical support.

Study funding

This study was supported in part by Fondo de Investigaciones Sanitarias, FEDER, Spain (FIS 14/00203, J.D.; FIS 18/00067 L.S.), NIH RO1NS077851 (J.D.), and Fundació Cellex (J.D.).

E. Martínez-Hernández is a recipient of a Juan Rodés grant (JR17/00012) from the Instituto de Salud Carlos III, Spain.

Disclosure

R. Ruiz-Garcia, B. Joubert, E. Martínez-Hernández, M. Petit-Pedrol, E. Pajarón-Boix, V. Fernández, L. Salais, M. del Pozo, T. Armangué, and L. Sabater report no disclosures. J. Dalmau receives royalties from Athena Diagnostics for the use of Ma2 as an autoantibody test and from Euroimmun for the use of NMDA, GABAB receptor, GABAA receptor, DPPX and IgLON5 as autoantibody tests. He is the editor of *Neurology: Neuroimmunology & Neuroinflammation*. F. Graus receives royalties from Euroimmun for the use of IgLON5 as an autoantibody test and honoraria for Assistant Editor of *MedLink Neurology*. Go to Neurology.org/NN for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* November 5, 2019. Accepted in final form November 18, 2019.

Appendix Authors

Name	Location	Role	Contribution
Raquel Ruiz-Gracia, PhD	Hospital Clínic, Barcelona, Spain	Author	Designed and conceptualized the study; analyzed the data; immunohistochemical studies; drafted the first version of the manuscript; and revised the manuscript for intellectual content
Eugenia Martínez-Hernández, MD, PhD	Hospital Clínic and IDIBAPS, Barcelona, Spain	Author	Acquisition of clinical data; analysis and interpretation of the data; and revised the manuscript for intellectual content
Bastien Joubert, MD	IDIBAPS, Barcelona, Spain	Author	Analysis of antibody effects and revised the manuscript for intellectual content
Mar Petit-Pedrol, PhD	IDIBAPS, Barcelona, Spain	Author	Analysis of antibody effects and revised the manuscript for intellectual content
Elena Pajarón-Boix, MD	Hospital General Castellón, Spain	Author	Acquisition of clinical data; analysis and interpretation of the data; and revised the manuscript for intellectual content
Verónica Fernández, MD	Hospital General Albacete, Spain	Author	Acquisition of clinical data; analysis and interpretation of the data; and revised the manuscript for intellectual content

Appendix (continued)

Name	Location	Role	Contribution
María Del Pozo, MD	Hospital General Albacete, Spain	Author	Acquisition of clinical data; analysis and interpretation of the data; and revised the manuscript for intellectual content
Lidia Salais, MD	Hospital General Castellón, Spain	Author	Acquisition of clinical data; analysis and interpretation of the data; and revised the manuscript for intellectual content
Thais Armangué, MD, PhD	IDIBAPS, Barcelona, Spain	Author	Acquisition of clinical data; analysis and interpretation of the data; and revised the manuscript for intellectual content
Lidia Sabater, PhD	IDIBAPS, Barcelona, Spain	Author	Acquisition of clinical data; analysis and interpretation of the data; obtained funding; and revised the manuscript for intellectual content
Josep Dalmau, MD, PhD	Hospital Clínic, ICREA Barcelona, Spain and University of Pennsylvania, PA, USA	Author	Acquisition of clinical data; review of clinical data; drafted the first version of the manuscript; revised the manuscript for intellectual content; and obtained funding
Francesc Graus, MD, PhD	IDIBAPS, Barcelona, Spain	Author	Designed and conceptualized the study; acquisition of clinical data; review of clinical data; analysis and interpretation of the data; revised the manuscript for intellectual content; and wrote the final version

References

1. Dalmau J, Geis C, Graus F. Autoantibodies to synaptic receptors and neuronal cell surface proteins in autoimmune diseases of the central nervous system. *Physiol Rev* 2017;97:839–887.
2. Joubert B, Rostásy K, Honnorat J. Immune-mediated ataxias. *Handb Clin Neurol* 2018;155:313–332.
3. Ances BM, Vitaliani R, Taylor RA, et al. Treatment-responsive limbic encephalitis identified by neuropil antibodies: MRI and PET correlates. *Brain* 2005;128:1764–1777.
4. Dalmau J, Gleichman AJ, Hughes EG, et al. Anti-NMDA-receptor encephalitis: case series and analysis of the effects of antibodies. *Lancet Neurol* 2008;7:1091–1098.
5. Spatola M, Sabater L, Planagumà J, et al. Encephalitis with mGluR5 antibodies: symptoms and antibody effects. *Neurology* 2018;90:e1964–e1972.
6. Shigemoto R, Kinoshita A, Wada E, et al. Differential presynaptic localization of metabotropic glutamate receptor subtypes in the rat hippocampus. *J Neurosci* 1997;17:7503–7522.
7. Sillevs Smitt P, Kinoshita A, De Leeuw B, et al. Paraneoplastic cerebellar ataxia due to autoantibodies against a glutamate receptor. *N Engl J Med* 2000;342:21–27.
8. Ferraguti F, Shigemoto R. Metabotropic glutamate receptors. *Cell Tissue Res* 2006;326:483–504.
9. Petit-Pedrol M, Sell J, Planaguma J, et al. LGI1 antibodies alter Kv1.1 and AMPA receptors changing synaptic excitability, plasticity and memory. *Brain* 2018;141:3144–3159.
10. Yu LJ, Wall BA, Wangari-Talbot J, Chen S. Metabotropic glutamate receptors in cancer. *Neuropharmacology* 2017;115:193–202.

Neurology® Neuroimmunology & Neuroinflammation

Paraneoplastic cerebellar ataxia and antibodies to metabotropic glutamate receptor 2

Raquel Ruiz-García, Eugenia Martínez-Hernández, Bastien Joubert, et al.

Neurol Neuroimmunol Neuroinflamm 2020;7;

DOI 10.1212/NXI.0000000000000658

This information is current as of December 11, 2019

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services

including high resolution figures, can be found at:
<http://nn.neurology.org/content/7/2/e658.full.html>

References

This article cites 10 articles, 1 of which you can access for free at:
<http://nn.neurology.org/content/7/2/e658.full.html##ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

Autoimmune diseases

http://nn.neurology.org/cgi/collection/autoimmune_diseases

Clinical neurology examination

http://nn.neurology.org/cgi/collection/clinical_neurology_examination

Diagnostic test assessment

http://nn.neurology.org/cgi/collection/diagnostic_test_assessment

Gait disorders/ataxia

http://nn.neurology.org/cgi/collection/gait_disorders_ataxia

Paraneoplastic syndrome

http://nn.neurology.org/cgi/collection/paraneoplastic_syndrome

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://nn.neurology.org/misc/about.xhtml#permissions>

Reprints

Information about ordering reprints can be found online:

<http://nn.neurology.org/misc/addir.xhtml#reprintsus>

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.