Understanding neurodegenerative changes of the afferent visual pathway in MS

Hanna G. Zimmermann, PhD, and Frederike Cosima Oertel, MD

Neurol Neuroimmunol Neuroinflamm 2020;7:e667. doi:10.1212/NXI.0000000000000667

Correspondence

Dr. Zimmermann

hanna.zimmermann@charite.de

RELATED ARTICLE

Article

MS optic neuritis-induced long-term structural changes within the visual pathway

Page e665

The chronic involvement of the afferent visual system in multiple sclerosis (MS) is a powerful model to understand the temporal course and functional relevance of neurodegeneration because it is very well distinguishable from neighboring structures and offers quantifiable structural and functional metrics. Optical coherence tomography (OCT) of the retina and diffusion tensor imaging (DTI)-based tractography of the optic radiation have proven to provide sensitive markers for neurodegeneration of the visual pathway in MS.^{1,2} OCT-derived neuroaxonal retinal damage, represented by thinning of the retinal nerve fiber layer (RNFL) and ganglion cell/inner plexiform layer (GCIPL), has been reported in patients with MS with a history of optic neuritis (ON) and—to a lesser extent—also independent of ON episodes.¹ RNFL and GCIPL show strong associations with visual function in MS, especially with low-contrast visual acuity.³ Both layer thicknesses are associated with brain atrophy: this association being stronger in the eyes without a history of ON.³

DTI facilitates the reconstruction and analysis of white matter tracts and can thereby reveal microstructural changes of the optic radiation as the most frequently affected white matter pathway in MS.² Although fractional anisotropy (FA) is considered as a marker of microstructural integrity, higher radial diffusivity (RD) can be caused by demyelination and axial diffusivity (AD) changes are suggested to be associated with axonal damage.²

In this issue of *Neurology® Neuroimmunology & Neuroinflammation*, Pawlitzki et al.⁴ describe retinal neuroaxonal loss and microstructural changes in the optic radiation in MS, using OCT and DTI-based tractography. In this cross-sectional study, 17 patients with a history of a single unilateral ON episode and a sex- and age-matched cohort of 11 patients without an ON episode were included. Pawlitzki et al. found more retinal atrophy and microstructural damage of the optic radiation in patients with a history of ON—measured by thinning of the RNFL and GCIPL as well as FA, AD, and RD changes in the optic radiation. This clearly describes the pronounced damage in the visual system after a single ON episode. Furthermore, by reporting associations of RNFL and GCIPL thickness with DTI parameters, the study is an important confirmation of long-term anterograde and retrograde neuroaxonal loss in the afferent visual system. Interestingly, the thickness of the primary visual cortex was neither associated with optic radiation DTI parameters nor with RNFL and GCIPL thinning.⁴

DTI and retinal atrophy have emerged as important outcome measures in clinical MS research.¹ However, to widely implement OCT and DTI for the individual monitoring of disease progression and treatment response, we still lack important pathologic and technical considerations.

The great advantages of OCT parameters include the rigorous and uniform quality control criteria, the reproducible correlation with visual function, and the measurability of even small degrees of atrophy.^{1,5} However, there is evidence that the test-retest reliability is associated with scan protocols, scan quality, rater bias, or physiologic variation of common OCT parameters^{6,7} while keeping in mind that OCT is not histopathology. Dead and nonfunctional ganglion cells

From the Experimental and Clinical Research Center (H.G.Z., F.C.O.), Max-Delbrück-Centrum für Molekulare Medizin and Charité–Universitätsmedizin Berlin, Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Germany; NeuroCure Clinical Research Center (H.G.Z., F.C.O.), Charité–Universitätsmedizin Berlin, Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Germany; and Multiple Sclerosis Center (F.C.O.), Department of Neurology, University of California San Francisco.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

might also have a volume effect; meanwhile, other cells such as astrocytes could contribute to thickness measurements. There is evidence that edema—associated with chronic inflammation—occurs in the RNFL, one of the most common OCT outcome parameters.⁸ Thus, the effect of pseudoatrophy, as we know it from MRI-based brain volume measurements, might also be present in the RNFL.

The same applies to DTI-based metrics.² In contrast to OCT, the protocols for DTI acquisition and postprocessing are still very heterogeneous, which substantially limits the use of DTI as a potential biomarker.^{2,9} Furthermore, the diffusion tensors in the optic radiation are influenced by optic radiation lesions and (Wallerian) neurodegeneration resulting from optic nerve damage.² So far, researchers have not succeeded in untangling these effects. In addition, the study of Pawlitzki et al.⁴ does not shed light on this enigma: the frequency of lesions in the optic radiation is lower in the non-ON group, and the extent of ON-independent visual pathway damage in this cohort could not be evaluated because the study did not include healthy controls.

The lack of normal reference values, confounded with the wide range of physiologic variation, challenge the interpretation of imaging markers and, thus, their application for individualized monitoring in MS. In the future, improving OCT- and DTI-based metrics and combining their application with newer methods such as the volumetric analyses of the lateral geniculate nucleus will implement a gapless assessment of structure and function in the afferent visual system and thereby new pathophysiologic insights.¹⁰ For example, it is currently controversially discussed whether ON-independent retinal atrophy in MS is a result of subclinical ON episodes, ongoing widespread inflammation and demyelination, trans-synaptic retrograde neurodegeneration from lesions in the white matter pathway or a primary retinal degeneration. None of these hypotheses is mutually exclusive,

and each of these may partially contribute to our observation. Closing the gap between these metrics by applying ongoing methodical advancements of afferent visual system research in longitudinal studies will expand our pathophysiologic understanding and the differentiation of normal and disease-related neurodegeneration in the afferent visual system of patients with MS.

Study funding

No targeted funding reported.

Disclosure

H.G.Z. reports research grants from Novartis, unrelated to this study. F.C.O. was employed by Nocturne, unrelated to this study. Go to Neurology.org/NN for full disclosures.

References

1. Petzold A, Balcer LJ, Calabresi PA, et al. Retinal layer segmentation in multiple sclerosis: a systematic review and meta-analysis. *Lancet Neurol* 2017;16:797–812.
2. Kuchling J, Brandt AU, Paul F, Scheel M. Diffusion tensor imaging for multilevel assessment of the visual pathway: possibilities for personalized outcome prediction in autoimmune disorders of the central nervous system. *EPMA J* 2017;8:279–294.
3. Oertel FC, Zimmermann HG, Brandt AU, Paul F. Novel uses of retinal imaging with optical coherence tomography in multiple sclerosis. *Expert Rev Neurother* 2019;19:31–43.
4. Pawlitzki M, Horbruegger M, Loewe K, et al. MS optic neuritis-induced long-term structural changes within the visual pathway. *Neurol Neuroimmunol Neuroinflamm* 2020;7:e665. doi: 10.1212/NX1.0000000000000665.
5. Tewarie P, Balk L, Costello F, et al. The OSCAR-IB consensus criteria for retinal OCT quality assessment. *PLoS One* 2012;7:e34823.
6. Oberwahrenbrock T, Weinhold M, Mikolajczak J, et al. Reliability of intra-retinal layer thickness estimates. *PLoS One* 2015;10:e0137316.
7. Oberwahrenbrock T, Traber GL, Lukas S, et al. Multicenter reliability of semi-automatic retinal layer segmentation using OCT. *Neurol Neuroimmunol Neuroinflamm* 2018;5:e449. doi: 10.1212/NX1.0000000000000449.
8. Oertel FC, Outteryck O, Knier B, et al. Optical coherence tomography in myelin-oligodendrocyte-glycoprotein antibody-seropositive patients: a longitudinal study. *J Neuroinflammation* 2019;16:154.
9. Kuchling J, Backner Y, Oertel FC, et al. Comparison of probabilistic tractography and tract-based spatial statistics for assessing optic radiation damage in patients with autoimmune inflammatory disorders of the central nervous system. *Neuroimage Clin* 2018;19:538–550.
10. Papadopoulou A, Gaetano L, Pfister A, et al. Damage of the lateral geniculate nucleus in MS: assessing the missing node of the visual pathway. *Neurology* 2019;92:e2240–e2249.

Neurology® Neuroimmunology & Neuroinflammation

Understanding neurodegenerative changes of the afferent visual pathway in MS

Hanna G. Zimmermann and Frederike Cosima Oertel

Neurol Neuroimmunol Neuroinflamm 2020;7;

DOI 10.1212/NXI.0000000000000667

This information is current as of January 22, 2020

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/7/2/e667.full.html
References	This article cites 10 articles, 2 of which you can access for free at: http://nn.neurology.org/content/7/2/e667.full.html##ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): DWI http://nn.neurology.org/cgi/collection/dwi Evoked Potentials/Visual http://nn.neurology.org/cgi/collection/evoked_potentials-visual MRI http://nn.neurology.org/cgi/collection/mri Multiple sclerosis http://nn.neurology.org/cgi/collection/multiple_sclerosis Optic neuritis; see Neuro-ophthalmology/Optic Nerve http://nn.neurology.org/cgi/collection/optic_neuritis
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Online ISSN: 2332-7812.