Encephalitis with radial perivascular emphasis

Not necessarily associated with GFAP antibodies

Jonathan Wickel, MD,* Ha-Yeun Chung, MD,* Klaus Kirchhof, MD, David Boeckler, MD, Stefan Merkelbach, MD, Peter Kuzman, MD, Wolf C. Mueller, MD, Christian Geis, MD,† and Albrecht Günther, MD†

Correspondence

Dr. Wickel
jonathan.wickel@med.uni-jena.de

Neurol Neuroimmunol Neuroinflamm 2020;7:e670. doi:10.1212/NXI.0000000000000670

Abstract

Objective

Autoimmune steroid-responsive meningoencephalomyelitis with linear perivascular gadolinium enhancement in brain MRI is regarded as glial fibrillary acidic protein (GFAP) astrocytopathy characterized by anti-GFAP antibodies (ABs). We questioned whether anti-GFAP ABs are necessarily associated with this syndrome.

Methods

Two patients with a strikingly similar disease course suggestive of autoimmune GFAP astrocytopathy are reported. Clinical examination, MRI, laboratory, and CSF analysis were performed. Neuropathologic examination of brain tissue was obtained from one patient. Serum and CSF were additionally tested using mouse brain slices, microglia-astrocyte cocultures, and a GFAP-specific cell-based assay.

Results

Both patients presented with subacute influenza-like symptoms and developed severe neurocognitive and neurologic deficits and impaired consciousness. MRIs of both patients revealed radial perivascular gadolinium enhancement extending from the lateral ventricles to the white matter suggestive of autoimmune GFAP astrocytopathy. Both patients responded well to high doses of methylprednisolone. Only one patient had anti-GFAP ABs with a typical staining pattern of astrocytes, whereas serum and CSF of the other patient were negative and showed neither reactivity to brain tissue nor to vital or permeabilized astrocytes. Neuropathologic examination of the anti-GFAP AB-negative patient revealed infiltration of macrophages and T cells around blood vessels and activation of microglia without obvious features of clasmotodendrosis.

Conclusions

The GFAP-AB negative patient had both a striking (para)clinical similarity and an immediate response to immunotherapy. This supports the hypothesis that the clinical spectrum of steroid-responsive meningoencephalomyelitis suggestive of autoimmune GFAP astrocytopathy may be broader and may comprise also seronegative cases.

*These authors contributed equally to the manuscript.

†Shared seniority.

From the Hans Berger Department of Neurology (J.W., H.-Y.C., C.G., A.G.), Section of Translational Neuroimmunology, Jena University Hospital Germany; Department of Neuroradiology (K.K.), Jena University Hospital, Germany; Department of Neurology (D.B.), Sana Hospital Borna, Germany; Department of Neurology (S.M.), Heinrich-Braun Hospital, Zwickau; and Department of Neuropathology (P.K., W.C.M.), Leipzig University Hospital, Germany.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

AB = antibody; GFAP = glial fibrillary acidic protein; IgG = Immunoglobulin G.

First described in 2016, autoimmune glial fibrillary acidic protein (GFAP) astrocytopathy has been characterized as a rare CNS disorder, typically manifesting as a steroid-responsive encephalitis, meningitis, myelitis, or meningoencephalomyelitis. Neurologic symptoms such as (sub)acute encephalopathy, blurred vision, postural tremor, and seizures often occur following an initial prodromal phase with influenza-like symptoms.^{1,2} A characteristic feature in brain MRIs is a linear perivascular gadolinium enhancement in the white matter extending radially outward from the ventricles. In addition, extensive lesions of the spinal gray matter may be detected.¹ No definite diagnostic criteria have been established yet.³ For diagnosis of autoimmune GFAP astrocytopathy, detection of GFAP antibodies (ABs) in the patient's CSF or serum is required.¹ CSF cell count and CSF protein level are usually abnormal. Disease onset has been reported after influenza-like symptoms, a preceding herpes simplex encephalitis, and varicella zoster encephalitis, respectively.^{1,4} Other ABs, e.g., NMDA receptor IgG and aquaporin-4 IgG may also be detected in autoimmune GFAP astrocytopathy. Furthermore, the disease can be related to an underlying malignancy, with ovarian teratoma being the most common.¹ The majority of patients improve after treatment with immunotherapy, especially corticosteroids.² Yet, the pathophysiology of autoimmune GFAP astrocytopathy is unknown. Here, we compare clinical, radiologic, and serologic findings of two patients with a very similar disease course suggestive of GFAP astrocytopathy. Despite intriguing similarity, only one patient harbored anti-GFAP ABs in serum and CSF. We here discuss that the characteristic clinical syndrome of autoimmune meningoencephalomyelitis with linear perivascular gadolinium enhancement may not necessarily be associated with anti-GFAP ABs.

Case report 1

During winter season, a 53-year-old Caucasian man was admitted to hospital due to influenza A infection. In the following weeks, the patient developed cognitive impairment, ataxia, tremor, and a left gaze-evoked nystagmus (for details, see table 1). Cranial MRI revealed areas with diffuse periventricular T2 hyperintensities and linear perivascular gadolinium enhancement in the supratentorial white matter extending radially outward from the ventricles (figure, A). CSF diagnostics revealed 86 cells/ μ L and 1,075 mg/L protein. High titers of anti-GFAP IgG ABs (titer 1:320) were found in CSF and serum. A typical staining pattern restricted to astrocytes could be detected after incubation of mouse brain slices and astrocyte and microglia cocultures with the patient's CSF. A confirmatory cell-based assay with the GFAP α isoform (Euroimmun, Lübeck, Germany) was positive when

incubated with the patient's CSF (figure, B). After other differential diagnoses were ruled out (table 2), autoimmune GFAP astrocytopathy was diagnosed. The patient was treated with methylprednisolone 1000 mg/d for 5 consecutive days. A rapid clinical improvement and a reduction of anti-GFAP IgG AB titers (CSF 1:100, serum 1:100) could be observed. MRI follow-up revealed regressive gadolinium enhancement and decreased periventricular T2 hyperintensities. In addition, EEG follow-up showed improvement with basal alpha activity. When the daily oral prednisolone dose was reduced to less than 20 mg during the following months, the patient's condition deteriorated again as he developed tremor. We therefore initiated 6 cycles of immunoadsorption and subsequently started rituximab treatment. This led to a clinical improvement with almost complete remission of clinical symptoms within 2 weeks.

Case report 2

In the same month, a 63-year-old Caucasian man presented with influenza-like symptoms. He later developed cognitive impairment, aggressive behavior, ataxia, and apraxia (for details, see table 1). MRI revealed pronounced T2 hyperintensities and gadolinium enhancement extending radially along the vessels within the supratentorial white matter (figure, C). CSF analysis revealed 53 cells/ μ L and 2,130 mg/L protein. Repetitive testing for antineuronal, anti-myelin oligodendrocyte glycoprotein, and anti-GFAP ABs was negative including incubation of both the patient's serum and CSF in mouse and monkey brain slices, vital respectively fixed glia cocultures after permeabilization, and a GFAP cell-based assay (figure, D). As most differential diagnoses were ruled out (table 2), the patient underwent frontal brain biopsy. Biopsy featured parenchymal blood vessel associated infiltration of macrophages and T cells. Brain tissue itself showed activation of microglia, infiltration of macrophages, and astrogliosis (figure, E). Although not specific, histopathologic findings are compatible with the few available histology reports of proven cases of autoimmune GFAP astrocytopathy in the literature.⁵ We did not observe direct signs of clasmotodendrosis or loss of aquaporin-4 (not shown), which is typically seen in anti-aquaporin-4 AB-positive neuromyelitis optica spectrum disorder.

Following treatment with IV immunoglobulins, the patient's condition deteriorated as he developed brainstem symptoms including dysarthria, dysphagia, and impaired consciousness. Artificial ventilation was required. Following methylprednisolone pulse therapy of 1000 mg/d for 5 consecutive days, a rapid clinical improvement was observed, and the patient could be extubated and was able to eat and walk again. Brain MRI confirmed a significant regression of T2 lesions and

Table 1 Clinical characteristics of case reports 1 and 2

Case report 1 (GFAP AB positive)	Case report 2 (GFAP AB negative)
53-year-old man	63-year-old man
Prodromal symptoms	
Upper respiratory tract infection with dry cough for several weeks (laboratory confirmed influenza A)	Upper respiratory tract infection with dry cough for 3 weeks during influenza season
Weight loss (10 kg in 4 weeks)	Weight loss (6 kg in 3 weeks)
Fever and night sweats	Fever
Fatigue	Fatigue
Neuropsychiatric symptoms	
Progressive personality change for several months with apathy	Progressive personality change for several months with apathy, confabulations, and aggressive behavior
Neurologic symptoms	
Left gaze-evoked nystagmus, ataxia in arms and legs, gait ataxia, intention and resting tremor of the upper limbs, and repetitive aphasia	Ataxia of arms and legs, gait ataxia, dysarthria, dysphagia, decreased consciousness, and apraxia
Neurocognitive symptoms	
Deficits in attention, word fluency, and memory	Deficits in attention, word fluency, abstraction ability, learning, and memory
Treatment	
Response to methylprednisolone	Deterioration after IVIG
Relapse after methylprednisolone tapering; stabilization additional treatment with immunoadsorption and rituximab	Response to methylprednisolone
	Relapse after methylprednisolone tapering
	Stabilization with immunoadsorption; additional treatment with azathioprine

Abbreviations: GFAP = glial fibrillary acidic protein; IVIG = intravenous immunoglobulin.

decreased radial perivascular gadolinium enhancement. Thus, oral prednisolone treatment was continued and tapered over the next months. EEG follow-up showed a significant improvement with basic alpha but still intermittent bifrontal delta activity. CSF measurement 3 months after first hospital admission revealed lower but still increased cell count (33 cells/ μ L) and protein levels (1,100 mg/L protein). Thereafter, 5 cycles of immunoadsorption were initiated with subsequent clinical improvement. However, after tapering prednisolone to less than 10 mg daily symptoms reoccurred, MRI demonstrated radial perivascular gadolinium enhancement, and CSF showed a persistent pleocytosis of 31 cells/ μ L. Additional treatment with azathioprine was initiated, and prednisolone was increased to 80 mg again and tapered over the following months.

Discussion

Detection of anti-GFAP ABs and typical MRI findings are regarded as essential features in the diagnosis of autoimmune GFAP astrocytopathy. We here compare two patients with a similar disease course suggestive of autoimmune GFAP astrocytopathy. Yet, only in one patient, anti-GFAP ABs were

detected. Recent reports questioned the relevance of anti-GFAP ABs in this clinical syndrome.^{3,6} In line with these reports, we here support the hypothesis that autoimmune meningoencephalomyelitis with characteristic MRI findings and steroid responsiveness may present with diverse immunologic findings, and the presence of anti-GFAP ABs is not obligatory. So far, it is not clear whether the presence of anti-GFAP-ABs in some patients with this disorder is just an immunologic accompaniment or whether these patients with anti-GFAP-ABs represent a particular subgroup with a specific pathophysiology targeting the astrocyte.

Because anti-GFAP ABs bind to astrocytic cytosolic intermediate filaments and ABs are not internalized, a direct pathophysiologic relevance of these ABs is unlikely. As known from autoimmune encephalitis and neuromyelitis optica spectrum disorders, pathophysiologically relevant ABs usually target membrane surface proteins.⁷ In contrast, in paraneoplastic disorders, specific onconeural ABs to intracellular antigens are not pathogenic but excellent biomarkers.⁷ In autoimmune GFAP astrocytopathy, a role of autoreactive T cells triggering a GFAP-specific autoimmune response has been suggested.³ In

Patient 1: (A.a) T2-weighted images demonstrate diffuse periventricular hyperintense lesions (thick arrows, A.a). Axial (A.b) and sagittal (A.c) T1-weighted images with gadolinium show linear perivascular enhancement extending radially through the periventricular white matter (thin arrows). (B) Characteristic staining pattern of GFAP-positive astrocytes in mouse brain slices incubated with CSF of patient 1 (B.a: hippocampus, coronal; magnification $\times 100$; B.b: Cerebellum, sagittal; magnification $\times 200$). Incubation of astrocyte-microglial cocultures with CSF of patient 1 showed characteristic staining of astrocytes in fixed cells after permeabilization (B.c; magnification $\times 400$), but not in vital cells (B.d; magnification $\times 200$). Incubation of a cell-based assay transfected with the GFAP α isoform with the patient's CSF revealed a positive staining pattern (B.e; magnification $\times 200$). Scale bars: 50 μ m each. **Patient 2:** (C) brain MRI shows similar features as shown in A with periventricular T2 lesions (thick arrows, C.a) and linear perivascular gadolinium enhancement (thin arrows, C.b, C.c). (D) Immunofluorescence stains with CSF of patient 2 revealed no specific staining in brain tissue (D.a: Hippocampus, coronal; magnification $\times 100$; D.b: cerebellum, sagittal; magnification $\times 200$) and no in permeabilized (D.c; magnification $\times 400$) and vital cells (D.d; magnification $\times 200$) in astrocyte-microglial coculture. Incubation of a cell-based assay transfected with the GFAP α isoform with the patient's CSF did not show binding (D.e; magnification $\times 200$). Scale bars: 50 μ m each. (E) Histological analysis of patient 2 brain biopsy showed blood vessel-associated infiltration by hematopoietic cells (E.a; hematoxylin and eosin stain; magnification $\times 200$; E.d; CD5-positive T lymphocytes; magnification $\times 200$). Only scattered infiltration by single cytotoxic T cells was observed (E.b; CD8; magnification $\times 200$). The majority of infiltrating cells were identified as macrophages (E.c; CD68; magnification $\times 200$). Brain tissue showed microglial activation (E.e; CD68; magnification $\times 200$) and astrogliosis (E.f; GFAP; magnification $\times 200$), but no obvious signs of clasmotodendrosis. Scale bars: 50 μ m each. GFAP = glial fibrillary acidic protein.

Table 2 Diagnostics of case reports 1 and 2 before treatment

Case report 1 (GFAP AB positive)	Case report 2 (GFAP AB negative)
Blood tests	
Negative, e.g., for vasculitis, viral and bacterial infections	Negative, e.g., for vasculitis, viral and bacterial infections (including Borna virus, <i>Chlamydia trachomatis</i> , <i>Mycoplasma pneumoniae</i> , and dengue virus), and toxoplasmosis
CSF	
Lymphocytic pleocytosis	Lymphocytic pleocytosis (predominantly T lymphocytes)
Increased albumin quotient (20.2)	Increased albumin quotient (19.8)
Increased lactate (2.5 mmol/L)	Increased lactate (3.4 mmol/L)
Initially, synthesis of IgG, IgM, und IgA; oligoclonal bands positive in CSF only	Initial intrathecal IgM and IgA synthesis; oligoclonal bands negative
Normal soluble IL2 receptor; negative viral DNA (CMV, HSV, VZV, EBV, and HHV6)	Normal soluble IL2 receptor; normal protein 14–33, JC virus PCR negative; negative viral DNA (CMV, HSV, VZV, EBV, and HHV6)
Autoimmune and onconeural ABs	
Negative (anti-mGluR1, anti-mGluR2, NMDA-R, GAD65, GABA-bR, Glycin-R, LGI1, CASPR2, amphiphysin, anti-Yo, Hu, Ri, anti-PNMA2, anti-SOX1, anti-recoverin, anti-CV2, anti-titin, and anti-DPPX)	Negative (anti-Hu, Ri, ANNA-3, Yo, Tr/DNER, Ma/Ta, GAD65, amphiphysin, aquaporin-4, MOG, NMDA-R, GABA-B receptor, LGI1, CASPR2, IGLON5, ZIC4, DPPX, CARP VIII, Glycin-R, mGluR1, mGluR5, GABA-aR, Rho GTPase activating protein 26, recoverin, GluRD2, flotillin, ITPR1, Homer 3, and neurochondrin)
PET/CT	
No tumor	No tumor
Electroencephalography	
Generalized, frontally emphasized theta activity with intermittent rhythmic delta activity	Bifrontal rhythmic delta activity

Abbreviations: AB = antibody; CMV = Cytomegalievirus; EBV = Epstein-Barr virus; GFAP = glial fibrillary acidic protein; HSV = Herpes simplex virus; IgA = Immunoglobulin A; IgM = Immunoglobulin M; VZV = Varicella-zoster virus.

a rat model, CD3⁺ T cells have been shown in close proximity to GFAP-positive astrocytes, thereby inducing an immune response with production of IgG ABs to GFAP.⁸ This might be a first direct link of how the humoral response is initiated. Of note, in our patients and also in others, an observational link exists between viral infections, e.g., influenza A, herpes simplex, and varicella zoster and subsequent GFAP autoimmunity.¹ The association of viral CNS infections and other forms of autoimmune encephalitis is well established. Here, antigen presentation due to neuronal damage caused by viral inflammation or virus-induced costimulation of immune cells is favored to be a pathophysiologic mechanism involved in promoting encephalitis.⁹ Although influenza A infection does not necessarily lead to neuronal damage in the CNS, it may be a predisposing factor to develop GFAP autoimmunity.

The specificity of anti-GFAP ABs for meningoencephalomyelitis needs to be clarified in further studies, as serum anti-GFAP ABs can also be found in other diseases, e.g., traumatic brain injury, vascular dementia, and astrocytoma, or even in healthy controls.¹⁰

Acknowledgment

The authors thank the patients for their participation in this study and Claudia Sommer for excellent technical assistance.

Furthermore, they thank Euroimmun (Lübeck, Germany) for their support in antibody diagnostics and Prof. Dr. Christine Stadelmann-Nessler and Dr. Jonas Franz (Institute of Neuropathology, University Medical Center Göttingen, Germany) for providing histological staining of aquaporin 1 and 4.

Study funding

This study was funded by the Schilling Foundation (to C. Geis); Center for Sepsis Control and Care (CSCC; to C. Geis and J. Wickel); and Interdisciplinary Center for Clinical Research Jena University Hospital (IZKF; to H.-Y. Chung and J. Wickel).

Disclosure

In the preparation of this manuscript, the funding sources played no role. J. Wickel, H.-Y. Chung, K. Kirchhof, and D. Boeckler report no disclosures. S. Merkelbach received travel funding, speaker, and/or advisory board honoraria from Bayer, Biogen, Daiichi Sankyo, Genzyme, and Merck Serono. P. Kuzman and W. C. Mueller report no disclosures. C. Geis received advisory board honoraria and speaker honoraria from Roche and Alexion and received research support from the Schilling Foundation, the German

Research Council, and the German Ministry of Education. A. Günther received travel funding, speaker, and/or advisory board honoraria from Merz, Ipsen, Boehringer Ingelheim, Pfizer, Daiichi Sankyo, and Bayer. Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* September 4, 2019. Accepted in final form November 26, 2019.

Appendix Authors

Name	Location	Role	Contribution
Jonathan Wickel, MD	Jena University Hospital, Jena	Author	Drafted the manuscript and involved in treatment of the patients
Ha-Yeun Chung, MD	Jena University Hospital, Jena	Author	Drafted the manuscript and involved in treatment of the patients
Klaus Kirchhof, MD	Jena University Hospital, Jena	Author	Interpreted MRI data
David Boeckler, MD	Sana Hospital Borna, Borna	Author	Involved in treatment of the patients
Stefan Merkelbach, MD	Heinrich-Braun Hospital, Zwickau	Author	Involved in treatment of the patients
Peter Kuzman, MD	Leipzig University Hospital, Leipzig	Author	Provided neuropathology findings

Appendix (continued)

Name	Location	Role	Contribution
Wolf C. Mueller, MD	Leipzig University Hospital, Leipzig	Author	Provided neuropathology findings
Christian Geis, MD	Jena University Hospital, Jena	Author	Supervised the project, involved in treatment of the patients, and writing of the manuscript
Albrecht Günther, MD	Jena University Hospital, Jena	Author	Supervised the project, involved in treatment of the patients, and writing of the manuscript

References

1. Flanagan EP, Hinson SR, Lennon VA, et al. Glial fibrillary acidic protein immunoglobulin G as biomarker of autoimmune astrocytopathy: analysis of 102 patients. *Ann Neurol* 2017;81:298–309.
2. Iorio R, Damato V, Evoli A, et al. Clinical and immunological characteristics of the spectrum of GFAP autoimmunity: a case series of 22 patients. *J Neurol Neurosurg Psychiatry* 2018;89:138–146.
3. Shan F, Long Y, Qiu W. Autoimmune glial fibrillary acidic protein astrocytopathy: a review of the literature. *Front Immunol* 2018;9:2802.
4. Fang B, McKeon A, Hinson SR, et al. Autoimmune glial fibrillary acidic protein astrocytopathy: a novel meningoencephalomyelitis. *JAMA Neurol* 2016;73:1297–1307.
5. Shu Y, Long Y, Chang Y, et al. Brain immunohistopathology in a patient with autoimmune glial fibrillary acidic protein astrocytopathy. *Neuroimmunomodulation* 2018;25:1–6.
6. Xie L, Liu T, Yao H, et al. Autoimmune inflammatory meningoencephalitis in a patient negative for glial fibrillary acidic protein-specific immunoglobulin G. *Mult Scler Relat Disord* 2019;28:167–171.
7. Dalmau J, Geis C, Graus F. Autoantibodies to synaptic receptors and neuronal cell surface proteins in autoimmune diseases of the central nervous system. *Physiol Rev* 2017;97:839–887.
8. Park ES, Uchida K, Nakayama H. Establishment of a rat model for canine necrotizing meningoencephalitis (NME). *Vet Pathol* 2014;51:1151–1164.
9. Armangue T, Leypoldt F, Málaga I, et al. Herpes simplex virus encephalitis is a trigger of brain autoimmunity. *Ann Neurol* 2014;75:317–323.
10. Wang KK, Yang Z, Yue JK, et al. Plasma anti-glial fibrillary acidic protein autoantibody levels during the acute and chronic phases of traumatic brain injury: a transforming research and clinical knowledge in traumatic brain injury pilot study. *J Neurotrauma* 2016;33:1270–1277.

Neurology® Neuroimmunology & Neuroinflammation

Encephalitis with radial perivascular emphasis: Not necessarily associated with GFAP antibodies

Jonathan Wickel, Ha-Yeun Chung, Klaus Kirchhof, et al.
Neurol Neuroimmunol Neuroinflamm 2020;7;
DOI 10.1212/NXI.0000000000000670

This information is current as of February 4, 2020

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/7/2/e670.full.html
References	This article cites 10 articles, 1 of which you can access for free at: http://nn.neurology.org/content/7/2/e670.full.html#ref-list-1
Citations	This article has been cited by 3 HighWire-hosted articles: http://nn.neurology.org/content/7/2/e670.full.html#otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Immunology http://nn.neurology.org/cgi/collection/all_immunology Autoimmune diseases http://nn.neurology.org/cgi/collection/autoimmune_diseases MRI http://nn.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.