Erdheim-Chester disease among neuroinflammatory syndromes: the case for precision medicine

Macarena I. de la Fuente, MD, Marc K. Rosenblum, MD, Eli L. Diamond, MD, Viviane S. Tabar, MD, and Antonio Omuro, MD

Correspondence

Dr. Omuro
antonio.omuro@yale.edu

Neurol Neuroimmunol Neuroinflamm 2020;7:e686. doi:10.1212/NXI.0000000000000686

Erdheim-Chester disease (ECD) is a rare, non-Langerhans histiocytosis characterized by xanthogranulomatous infiltration typically affecting long bones, cardiovascular system, retroperitoneum, and lung, and that involves the CNS in 25%–50% of patients (table).^{1,2} Historically, establishing the diagnosis has been challenging, particularly in the absence of systemic abnormalities. Recent genomic studies have uncovered that approximately 50% of ECD tissue samples harbor a mutation in the *BRAF* gene,³ termed *BRAF*^{V600E}, and that pointed to a neoplastic, rather than inflammatory, nature of the disease.^{4,5}

We describe a case of ECD with isolated CNS presentation emphasizing the diagnostic challenges and how a precision medicine approach provided a path to successful treatment.

Clinical case

A 51-year-old right-handed man presented with a 3-month history of diplopia and falls. Neurologic examination revealed right VI nerve palsy and mildly ataxic gait, with no other findings. A brain MRI revealed multifocal FLAIR hyperintensities mainly in the posterior fossa, with nodular enhancement (figure 1, A and B). Extensive workup was unrevealing (lactate dehydrogenase, erythrocyte sedimentation rate, B2-microglobulin, angiotensin-converting enzyme, CSF analysis with flow cytometry, spine MRI, HIV, syphilis, and rheumatologic/inflammatory panels). Whole body fluorodeoxyglucose-positron emission tomography showed minor increased uptake within the pons and no systemic abnormalities. CT of the legs was normal.

The patient underwent stereotactic needle biopsy of a leading cerebellar lesion. Pathology showed fragments of normal cerebellum and no mutations on next-generation targeted sequencing of 422 genes (NGS). High-dose steroids were tried without clinical improvement. ECD was considered but felt unlikely because of the absence of systemic involvement.

Three weeks after corticosteroid tapering, an open biopsy was performed, but pathology and NGS were again unrevealing. Given mild but continuous worsening of symptoms, a decision was made to perform a third cerebellar biopsy. Pathology showed non-granulomatous lymphohistiocytic infiltrates, with CD3-labeling T cell infiltrates, no loss of pan-T antigen expression, and rare B cells. A very large number of CD163-labeling, CD1a/Langerin-negative mononuclear elements were present. Findings were reviewed by 3 pathologists; the possibility of an infection was raised, and infectious diseases genomic studies suggested. However, given normal CSF, and on discussions with an experienced pathologist (M.R.), tissue was prioritized for genomic studies focusing on neoplastic etiologies and ECD. Immunohistochemistry for *BRAF*^{V600E} mutation and RNA sequencing

From the Department of Neurology, Sylvester Comprehensive Cancer Center (M.I.d.l.F., A.O.), University of Miami, Miami, FL; and Department of Pathology (M.K.R.), Department of Neurology (E.L.D.), and Department of Neurosurgery (V.S.T.), Memorial Sloan Kettering Cancer Center, New York, NY.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by Yale Cancer Center.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Table Clinical manifestations of ECD

Organ involved	Manifestation	Incidence, %	Comment
Bone	Long bone osteosclerosis	80–95%	Presents with pain or asymptomatic.
Cardiovascular	Periaortic infiltration	46–62%	Usually asymptomatic
	Right atrium pseudotumor	9–57%	
	Coronary infiltration	27%	Risk for coronary stenosis and myocardial infarction
	Pericardial involvement	10–31%	Risk for tamponade
Pulmonary	Involvement of pleura, lung parenchyma or both	25–50%	Asymptomatic or manifest as dyspnea and/or cough
Retroperitoneum	Mass-like, perirenal infiltrative lesion	58–65%	“Hairy kidneys”
Central nervous system	Periorbital involvement	22–27%	Exophthalmos
	CNS involvement	37–38%	Cognitive impairment, neuropsychiatric and pyramidal syndrome
	Pituitary involvement	28–47%	Central diabetes insipidus
	Cerebellar involvement	17%	
	Dura involvement		Differential diagnosis with meningioma

Abbreviation: ECD = Erdheim-Chester disease.

panel (ARCHER) showed no abnormalities (no *BRAF*/*KIAA* gene fusions). NGS was initially reported as negative, but review of results and comparison of pathology accession numbers indicated DNA had been inadvertently extracted using tissue from a previous biopsy. NGS was repeated using the correct tissue and finally demonstrated a *BRAF*^{V600E} mutation, as well as mutations in *CHEK2*, *DOT1L*, *KDMSA*, and *MSH6* and deletions in *ROS1* and *ATXN2*. The integrated pathology diagnosis was ECD. The disease course timeline was summarized in figure 2.

Further staging included normal echocardiogram and cardiac MRI; repeat fluorodeoxyglucose-positron emission tomography/CT showed right knee mild hypermetabolism, questioning osseous involvement. Treatment with vemurafenib, an Food and Drug Administration-approved drug for ECD, was considered. However, based on the literature on *BRAF*^{V600E} melanoma suggesting improved efficacy and decreased toxicity with combined *BRAF* and *MEK1/2* inhibition, a regimen with dabrafenib and trametinib was favored. After insurance denial and a successful appeal process supported by the genomic findings, treatment was initiated, resulting in significant and early clinical and radiographic improvement (figure 1, C and D). The patient was still in remission 18+ months later.

Discussion

This report illustrates how a combination of precision medicine, perseverance, and clinical judgment may result in

successful diagnoses and treatments of rare diseases. Although this patient’s MRI did show findings suggestive of ECD, these are not pathognomonic and differential was broad. The absence of systemic findings further confounded the diagnosis.

Two biopsies and NGS were negative, and it would be tempting to categorize this case among unspecified CNS inflammatory disorders. However, the mismatch between MRI suggesting an active process and histology showing no signs of active inflammatory or neoplastic diseases suggested biopsies and NGS were not representative of the ongoing process. The lack of response to corticosteroids was another red flag for an inflammatory disease or CNS lymphoma diagnosis, prompting a third, and eventually representative biopsy. Histology remained inconclusive after multiple reviews, but genetic analysis finally provided the crucial missing information. Interestingly, *BRAF*^{V600E} was negative by immunohistochemistry.^{6,7} IHC is a sensitive and specific tool for detection of *BRAF*-V600E. However, negative or low staining cases should undergo genetic analysis, based on clinical and histopathologic features. It is also noteworthy that the first 2 gene sequencing attempts were negative because the tissue was not representative of the active process, containing normal cerebellum and no neoplastic cells to allow for detection of this somatic mutation.⁸

The genomic findings provided a path for utilization of dabrafenib and trametinib, possibly a better treatment of ECD than single-agent vemurafenib, but that is not Food and Drug Administration-approved, with only one treated

Figure 1 MRI imaging

(A and B) Baseline MRI. Axial FLAIR (A) sequence showing FLAIR signal hyperintensity lesions in the pons, middle cerebellar peduncles, and cerebellum (arrows), with corresponding solid enhancement on T1 postcontrast images (arrow heads). A follow-up MRI (C and D) performed 10 weeks into treatment, showing near complete resolution of the lesions.

patient reported in the literature.² We provide a second case successfully treated off-label with this combination. Overall, our patient illustrates how precision medicine can result in

successful diagnosis and therapies for previously untreatable conditions, although clinical judgment remains irreplaceable for realizing the full potential of emerging technologies.

Figure 2 Disease course timeline

Study funding

Study sponsor: Sylvester Comprehensive Cancer Center and Yale Brain Tumor Center.

Disclosure

M.I. de la Fuente served as a member on an ad hoc scientific advisory board for Puma Biotechnology, Agios Pharmaceuticals, and Forma Therapeutics and as a consultant for Foundation Medicine. M.K. Rosenblum has nothing to disclose. E.L. Diamond has nothing to disclose. V.S. Tabar is a scientific cofounder of Blue Rock Therapeutics and receives research support from the same. She is a scientific advisor for Robeauté. A. Omuro served as a member on an ad hoc scientific advisory board for BTG and Merck. Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures.

Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* December 16, 2019. Accepted in final form January 30, 2020.

Appendix Authors

Name	Location	Contribution
Macarena I. de la Fuente, MD	University of Miami	Data acquisition; drafting/revising the manuscript; analysis or interpretation of the data
Marc K. Rosenblum, MD	Memorial Sloan-Kettering Cancer Center	Data acquisition; drafting/revising the manuscript; analysis or interpretation of the data

Appendix (continued)

Name	Location	Contribution
Eli L. Diamond, MD	Memorial Sloan-Kettering Cancer Center	Drafting/revising the manuscript; analysis or interpretation of the data
Viviane S. Tabar, MD	Memorial Sloan-Kettering Cancer Center	Data acquisition
Antonio Omuro, MD	Yale Brain Tumor Center	Data acquisition; drafting/revising the manuscript; study concept or design; analysis or interpretation of the data

References

1. Diamond EL, Dagna L, Hyman DM, et al. Consensus guidelines for the diagnosis and clinical management of Erdheim-Chester disease. *Blood* 2014;124:483–492.
2. Al Bayati A, Plate T, Al Bayati M, Yan Y, Lavi ES, Rosenblatt JD. Dabrafenib and trametinib treatment for erdheim-chester disease with brain stem involvement. *Mayo Clin Proc Innov Qual Outcomes* 2018;2:303–308.
3. Haroche J, Charlotte F, Arnaud L, et al. High prevalence of BRAF V600E mutations in Erdheim-Chester disease but not in other non-Langerhans cell histiocytoses. *Blood* 2012;120:2700–2703.
4. Haroche J, Cohen-Aubart F, Emile JF, et al. Dramatic efficacy of vemurafenib in both multisystemic and refractory Erdheim-Chester disease and Langerhans cell histiocytosis harboring the BRAF V600E mutation. *Blood* 2013;121:1495–1500.
5. Cohen Aubart F, Emile JF, Carrat F, et al. Targeted therapies in 54 patients with Erdheim-Chester disease, including follow-up after interruption: the LOVE study. *Blood* 2017;130:1377–1380.
6. Breton Q, Plouhinec H, Prunier-Mirebeau D, et al. BRAF-V600E immunohistochemistry in a large series of glial and glial-neuronal tumors. *Brain Behav* 2017;7:e00641.
7. Anwar MA, Murad F, Dawson E, Abd Elmageed ZY, Tsumagari K, Kandil E. Immunohistochemistry as a reliable method for detection of BRAF-V600E mutation in melanoma: a systematic review and meta-analysis of current published literature. *J Surg Res* 2016;203:407–415.
8. Estrella JS, Tetzlaff MT, Bassett RL Jr, et al. Assessment of BRAF V600E status in colorectal carcinoma: tissue-specific discordances between immunohistochemistry and sequencing. *Mol Cancer Ther* 2015;14:2887–2895.

Neurology® Neuroimmunology & Neuroinflammation

Erdheim-Chester disease among neuroinflammatory syndromes: the case for precision medicine

Macarena I. de la Fuente, Marc K. Rosenblum, Eli L. Diamond, et al.

Neurol Neuroimmunol Neuroinflamm 2020;7;

DOI 10.1212/NXI.0000000000000686

This information is current as of March 2, 2020

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

Updated Information & Services

including high resolution figures, can be found at:
<http://nn.neurology.org/content/7/3/e686.full.html>

References

This article cites 8 articles, 5 of which you can access for free at:
<http://nn.neurology.org/content/7/3/e686.full.html##ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

All Clinical Neurology

http://nn.neurology.org/cgi/collection/all_clinical_neurology

All Medical/Systemic disease

http://nn.neurology.org/cgi/collection/all_medical_systemic_disease

All Oncology

http://nn.neurology.org/cgi/collection/all_oncology

Chemotherapy-tumor

<http://nn.neurology.org/cgi/collection/chemotherapytumor>

Primary brain tumor

http://nn.neurology.org/cgi/collection/primary_brain_tumor

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:

<http://nn.neurology.org/misc/about.xhtml#permissions>

Reprints

Information about ordering reprints can be found online:

<http://nn.neurology.org/misc/addir.xhtml#reprintsus>

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.