

# Serum antibodies to phosphatidylcholine in MS

Maria Cruz Sádaba, PhD, Veit Rothhammer, MD, PhD, Úrsula Muñoz, PhD, Cristina Sebal, PhD, Esther Escudero, PhD, Pia Kivisäkk, MD, PhD, Maria Isabel Garcia Sanchez, PhD, Guillermo Izquierdo, MD, PhD, Stephen L. Hauser, MD, Sergio E. Baranzini, PhD, Jorge R. Oksenberg, PhD, Roberto Álvarez-Lafuente, PhD, Rohit Bakshi, MD, Howard L. Weiner, MD, and Francisco J. Quintana, PhD

## Correspondence

Dr. Sádaba  
 mariacruz.sadabaarigaiz@ceu.es  
 or Dr. Quintana  
 fquintana@rics.bwh.harvard.edu

*Neurol Neuroimmunol Neuroinflamm* 2020;7:e765. doi:10.1212/NXI.0000000000000765

## Abstract

### Objective

To evaluate the value of serum immunoglobulin G (IgG) and immunoglobulin M (IgM) antibodies reactive with phosphatidylcholine (PC) and lactosylceramide (LC) as biomarkers in MS.

### Methods

We developed an ultrasensitive ELISA technique to analyze serum IgG and IgM antibodies to LC and PC, which we used to analyze samples from 362 patients with MS, 10 patients with non-MS myelin diseases (Non-MSMYDs), 11 patients with nonmyelin neurologic diseases (Non-MYNDs), and 80 controls. MS serum samples included clinically isolated syndrome (CIS, n = 17), relapsing-remitting MS (RRMS, n = 62), secondary progressive MS (SPMS, n = 50), primary progressive MS (PPMS, n = 37), and benign MS (BENMS, n = 36).

### Results

We detected higher levels of serum IgM antibodies to PC (IgM-PC) in MS than control samples; patients with CIS and RRMS showed higher IgM-PC levels than patients with SPMS, PPMS, and BENMS and controls. MS and control samples did not differ in serum levels of IgM antibodies reactive with LC, nor in IgG antibodies reactive with LC or PC.

### Conclusions

Serum IgM-PC antibodies are elevated in patients with MS, particularly during the CIS and RRMS phases of the disease. Thus, serum IgM-PC is a candidate biomarker for early inflammatory stages of MS.

### Classification of evidence

This study provides Class III evidence that serum antibodies to PC are elevated in patients with MS. The study is rated Class III because of the case control design and the risk of spectrum bias: antibody levels in patients with MS were compared with healthy controls.

## MORE ONLINE

### → Class of Evidence

Criteria for rating therapeutic and diagnostic studies

[NPub.org/coe](https://www.npub.org/coe)

From the Ann Romney Center for Neurologic Diseases (M.C.S., V.R., P.K., R.B., H.L.W., F.J.Q.), Brigham and Women's Hospital, Harvard Medical School, Boston, MA; Facultad de Medicina (M.C.S., U.M., C.S., E.E.), Instituto de Medicina Molecular Aplicada (INMA), Universidad San Pablo-CEU, CEU Universities, Madrid; Molecular Biology Service and MS Unit (M.I.G.S., G.I.), University of Sevilla; Department of Neurology (S.L.H., S.E.B., J.R.O.), University of California, San Francisco; Instituto de Investigación Sanitaria San Carlos (IdISSC) (R.A.-L.), Hospital Clínico San Carlos, Madrid, Spain; and The Broad Institute of Harvard and MIT (F.J.Q.), Cambridge, MA.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the NIH.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

## Glossary

**BCR** = B-cell receptor; **BENMS** = benign MS; **CIS** = clinically isolated syndrome; **IgG** = immunoglobulin G; **IgG-PC** = IgG to PC; **IgG-LC** = IgG to LC; **IgM** = immunoglobulin M; **IgM-LC** = IgM to LC; **IgM-PC** = IgM to PC; **LC** = lactosylceramide; **Non-MSMYD** = non-MS myelin disease; **Non-MYND** = nonmyelin neurologic disease; **PC** = phosphatidylcholine; **PPMS** = primary progressive MS; **RRMS** = relapsing-remitting MS; **SPMS** = secondary progressive MS.

The high prevalence of immunoglobulin G (IgG) and immunoglobulin M (IgM) oligoclonal bands in the CSF has guided multiple studies focused on the identification of antibody targets in MS.<sup>1,2</sup> Lipids are a major component of myelin, the main target of the autoimmune response in MS.<sup>3</sup> New antigen microarray techniques have allowed the detection of lipid-reactive antibodies in the CSF of patients with MS.<sup>4,5</sup> Indeed, IgG antibodies to sulfatide, ganglioside GM4, and galactocerebroside<sup>6–9</sup> worsen disease pathogenesis in MS experimental models.<sup>4</sup> In addition, patients with MS harboring CSF IgM reactive with phosphatidylcholine (PC) develop a more severe disease course.<sup>6,7,10–12</sup> Moreover, serum IgG antibodies to lactosylceramide (LC) are associated with cerebral tissue damage in patients with MS.<sup>13</sup> These data suggest that lipid-reactive antibodies may contribute to disease pathogenesis and constitute useful as biomarkers in MS.

In this study, we evaluated the potential of lipid-reactive antibodies as biomarkers in MS. We detected increased levels of serum IgM antibodies to PC (IgM-PC) in patients with MS. In contrast, we did not detect differences in serum IgG antibodies to PC (IgG-PC), nor IgM or IgG to LC (IgM-LC and IgG-LC, respectively) between MS and controls. Thus, serum IgM-PC is a candidate biomarker for MS.

## Methods

### Study design

We developed an ELISA technique to study IgG and IgM antibodies reactive with LC and PC in patients with MS. This Class III retrospective cohort control study included individuals from different cohorts (United States and Spain). The study included a narrow spectrum of persons with MS, non-MS myelin diseases (Non-MSMYDs) and nonmyelin neurologic diseases (Non-MYNDs) with a retrospective longitudinal follow-up, and without neurologic disease (controls). The MS patient and control samples analyzed are described in table 1. The samples were analyzed by ELISA by an investigator blinded to disease status; the diagnostic test results and disease status were determined by different team members. Clinical information was provided from a clinical database of the different patients with associated biobanks (blood and CSF samples).

### Patients

Samples were obtained from the Comprehensive Longitudinal Investigation of Multiple Sclerosis (CLIMB, Boston,

MA) and EPIC (San Francisco, CA), as part of the SUMMIT consortium<sup>14,15</sup>; from the Servicio de Neurología Hospital Virgen del Rocío (Sevilla, Spain); from the Servicio de Inmunología del Hospital Clínico San Carlos (Madrid, Spain); and from the Servicio de Neurología del Hospital Universitario Quirónsalud (Madrid, Spain). The classification and clinical data of all samples analyzed are summarized in table 1.

Peripheral blood samples from patients with MS (362), patients with Non-MSMYD (10), patients with Non-MYND (11), and from 80 individuals without neurologic diseases (control group) were collected. Patients were free of relapses at the time of blood sample collection. Sera samples were aliquoted and stored at  $-80^{\circ}\text{C}$  until analyzed.

### ELISA assay

To detect IgG and IgM antibodies reactive with LC (Matreya, State College, PA) and PC (Sigma-Aldrich, St. Louis, MO), 96-well plates were first coated with LC diluted in dimethyl sulfoxide and PC diluted in methanol at  $10\ \mu\text{g}/\text{mL}$ . After washing 3 times with phosphate-buffered saline, the wells were treated with blocking solution. Serum samples were diluted (1/50, 1/100, 1/200) in blocking solution and pipetted into the wells because we obtained the higher signal and a lower background. IgG or IgM antibodies were detected with anti-human IgG biotin and anti-human IgM biotin, respectively (Jackson ImmunoResearch, West Grove, PA), followed by avidin–horse peroxidase (Sigma-Aldrich). Finally, we used TMB-one (Kementec, Taastrup, Denmark) as substrate. Plates were read at 450 nm using an Infinite 200 PRO spectrophotometer (Tecan instrument, Zürich, Switzerland). We classified a serum as positive when the optic density was higher than  $\text{mean} \pm 2 \times \text{SEM}$  of the control group.

### Statistics

Statistical analyzes were performed with GraphPad Prism (version 6.0) and IBM SPSS 24 statistical packages. *p* Values  $<0.05$  were considered statistically significant. We used the Mann-Whitney *U* test for comparisons between 2 groups and the Kruskal-Wallis test for multiple group comparisons. To analyze the fraction of individuals harboring lipid-reactive antibodies, we used the Fisher test for comparisons between MS and control group and the Pearson  $\chi^2$  test for comparisons among different MS types. OR and 95% CI were used to estimate the association between the presence of antibodies and the disease.

**Table 1** Demographic and clinical data from patients with MS, Non-MSMYD, and Non-MYND and control group

Figure	Cohorts	Females	Age	Disease duration	EDSS score	Treatment
1, 2	Controls (80)					0 (0%)
	United States (60)	47 (78.3%)	44 (25–63)			
	Spanish (20)	17 (85%)	58 (28–88)			
3, 4	MS (362)	246 (70.7%)	48 (19–92)	13 (0–57)	3.0 (0–9)	120 (33%)
	United States (235)	173 (73.6%)	47 (19–92)	13 (0–51)	2.2 (0–9)	113 (48%)
	Spanish (127)	73 (57.5%)	49 (19–77)	15 (2–57)	4.4 (0–9)	7 (6%)
3, 4	CIS (17)	10 (58.8%)	44 (19–76)	10 (2–36)	1.9 (0–6)	0 (0%)
	RR (63)	43 (70.5%)	49 (23–92)	15 (2–56)	2.0 (0–7)	4 (6%)
	SP (50)	31 (62.0%)	59 (37–79)	22 (8–57)	6.4 (3–9)	7 (14%)
	PP (37)	18 (48.6%)	54 (31–75)	12 (4–27)	6.0 (2–9)	0 (0%)
	Benign (36)	20 (55.6%)	48 (34–61)	22 (4–42)	2.7 (0–8)	6 (17%)
	Control (80)	64 (80%)	48 (25–88)			0 (0%)
5	Non-MSMYD (10)	4 (40%)	41 (25–59)			
	Myelitis (4)					
	Polyneuropathy (2)					
	Optic neuritis (2)					
	Autoimmune myelopathy (1)					
5	Non-MYND (11)	8 (72.7%)	38 (2–78)			
	Intracranial hypertension (2)					
	Lymphocytic meningitis (1)					
	Schizophrenia (1)					
	Cerebellar syndrome (1)					
	Non-Hodgkin lymphoma (1)					
	Ataxia telangiectasia (1)					
	Cephalea (2)					
	ELA (1)					
	Meningoencephalitis (1)					

Abbreviations: CIS = clinically isolated syndrome; EDSS = Expanded Disability Status Scale; IFN = interferon; Non-MSMYD = non-MS myelin disease; Non-MNYD = nonmyelin neurologic disease; PP = primary progressive; RR = relapsing-remitting; SP = secondary progressive. Females: number of cases analyzed and the percentage in the group. Age, disease duration, and EDSS score: mean (minimum and maximum). Treatment: number and percentage of treated patients. Patients in figure 1 were treated with IFN-beta (84), glatiramer acetate (24), and natalizumab (6). Treatment used for the RR patients was IFN (5); SP patients were treated with IFN (1), glatiramer acetate (3), and natalizumab (3), and benign patients were treated with IFN (3) and natalizumab (3). PP patients were not treated.

## Standard protocol approvals, registrations, and patient consents

The study was approved by the Bioethics Committees of the participant hospitals. All the participants gave verbal and written informed consent for sample collection.

## Data availability

The corresponding authors will provide anonymized data of this study on reasonable request from any

qualified investigator, following relevant data protection regulations.

## Results

### Increased serum IgM antibodies to PC in MS

We first analyzed serum IgG and IgM antibodies to PC and LC in patients with MS and controls. We detected higher IgM

reactivity against PC (IgM-PC) in MS serum samples (95% CI 0.03–0.15; 95% CI 0.04–0.17; United States and Spain cohorts, respectively) (figure 1A). Serum IgM-PC levels were not related to sex, age, or disease duration (table e-1, links. [lww.com/NXI/A265](http://lww.com/NXI/A265)). In a preliminary study, we detected lower serum IgM-PC levels in patients with MS treated with natalizumab than in those untreated (data not shown). We did not detect significant differences in serum IgM to LC (IgM-LC) nor in IgG reactivity to PC (IgG-PC) or LC (IgG-LC) between patients with MS and controls (figure 1B and not shown).

### IgM-PC antibodies are associated with disease course and subtype

We then analyzed serum IgM-PC levels at different MS phases. We detected higher serum IgM-PC levels in clinically isolated syndrome (CIS) (95% CI 0.17–0.31; 95% CI 0.12–0.29), relapsing-remitting MS (RRMS) (95% CI 0.16–0.25; 95% CI 0.10–0.23), secondary progressive MS (SPMS) (95% CI 0.09–0.22; 95% CI 0.03–0.20), and primary progressive MS (PPMS) (95% CI 0.00–0.06; 95% CI 0.04–0.14) samples compared with benign MS (BENMS) or controls, respectively (figure 2A). Moreover, we detected higher serum IgM-PC antibody levels in CIS (95% CI –0.02 to 0.19; 95% CI 0.04–0.19) and RRMS (95% CI –0.01 to 0.10; 95% CI 0.02–0.13) samples than in SPMS or PPMS samples; controls harbored higher serum IgM-PC levels than BENMS (95% CI 0.00–0.07). We did not observe differences in IgM-PC serum antibodies between CIS and RRMS samples, nor between SPMS and PPMS samples. We detected increased IgM-PC levels (above percentile 50) in 23% of control, 89% of CIS, 87% of RRMS, 48% of SPMS, 59% of PPMS, and 8% of BENMS samples. We did not detect differences in IgM-LC levels when

we analyzed samples taken from different MS subtypes (figure 2B).


To further investigate the value of IgM-PC antibodies as biomarkers, we analyzed the fraction of IgM-PC positive samples, which was higher in the MS group than in controls (OR 4.66; 95% CI 2.40–9.07) (table 2). The fraction of IgM-PC positive samples was higher in the CIS (OR 22.50; 95% CI 4.53–111.90), RRMS (OR 24.00; 95% CI 8.78–65.63), SPMS (OR 4.14; 95% CI 1.78–9.62), and PPMS (OR 4.40; 95% CI 1.77–10.91) groups than in controls. We also observed significant differences between CIS (OR 60.00; 95% CI 9.87–364.90), RRMS (OR 64.00; 95% CI 17.38–235.60), SPMS (OR 11.05; 95% CI 1.39–36.02), or PPMS (OR 11.73; 95% CI 3.43–40.13) and BENMS. CIS (OR 5.43; 95% CI 1.12–26.34) and RRMS (OR 5.79; 95% CI 2.20–15.22) showed the highest fraction of IgM-PC compared with SPMS. We also detected significant differences in the frequency of IgM-PC positive serums between CIS (OR 5.11; 95% CI 1.02–25.72) or RRMS (OR 5.45; 95% CI 1.95–15.19) and PPMS (figure 3).

### IgM-PC antibodies in other neurologic diseases

Finally, we analyzed serum IgM-PC levels in samples from patients affected by other neurologic diseases. Patients with MS exhibited higher IgM-PC levels compared with the Non-MSMYD group (95% CI 0.02–0.23); these differences were larger compared with CIS samples (95% CI 0.09–0.34); similar IgM-PC levels were detected between MS or CIS samples and Non-MNYD (figure 4A).


We detected IgM-PC (above percentile 50) in 30% of Non-MSMYD and 63.7% Non-MNYD samples. In addition, we

**Figure 1** Serum IgM-PC and IgM-LC in patients with MS and controls



(A) Serum IgM-PC controls. Boxes represent the median of OD ± percentiles 25–75, and whiskers include 100% of the patients. Percentile 25 = 0.0559. Percentile 75 = 0.2416. (B) Serum IgM-LC. Boxes represent the median of OD ± percentiles 25–75, and whiskers include 100% of the patients. Percentile 25 = 0.0559. Percentile 75 = 0.2416. The Mann-Whitney *U* test was used to compare different groups. C = control group; IgM-LC = IgM to lactosylceramide; IgM-PC = IgM to phosphatidylcholine; NS = nonsignificant; OD = optic density.

**Figure 2** IgM-PC and IgM-LC in different MS phases



(A) Serum IgM-PC. Boxes represent the median of OD  $\pm$  percentiles 25–75, and whiskers include 100% of the patients. Percentile 25 = 0.0559. Percentile 75 = 0.2416. (B) Serum IgM-LC. Boxes represent the median of OD  $\pm$  percentiles 25–75, and whiskers include 100% of the patients. Percentile 25 = 0.0559. Percentile 75 = 0.2416. The Kruskal-Wallis test was used for multiple group comparisons, and when significances were detected, Mann-Whitney *U* test to compare 2 groups. BEN = benign; C = control group; CIS = clinically isolated syndrome; IgM-LC = IgM to lactosylceramide; IgM-PC = IgM to phosphatidylcholine; NS = nonsignificant; PP = primary progressive; RR = relapsing-remitting; SP = secondary progressive.

also detected significant differences in the fraction of IgM-PC-positive MS (OR 5.179; 95% CI 1.40–19.18) or CIS (OR 25.00; 95% CI 3.52–177.60) and Non-MSMYD samples (figure 4B).

**Table 2** IgM-PC and IgM-LC in patients with MS, Non-MSMYD, and Non-MYND and control group

	IgM-PC	IgM-LC	IgM-LIP
<b>CIS</b>	15 (88.2%)	5 (29.4%)	16 (94.1%)
<b>RR</b>	55 (88.7%)	4 (6.5%)	55 (88.7%)
<b>SP</b>	29 (58.0%)	10 (20.0%)	29 (58.0%)
<b>PP</b>	22 (59.5%)	6 (16.2%)	23 (62.2%)
<b>BEN</b>	4 (11.1%)	3 (8.3%)	7 (19.4%)
<b>Control</b>	13 (25.0%)	7 (13.5%)	16 (30.8%)
<b>Non-MSMYD</b>	3 (30.0%)		
<b>Non-MYND</b>	7 (63.6%)		

Abbreviations: BEN = benign; CIS = clinically isolated syndrome; IgM-LC = sera positive for IgM-LC; IgM-LIP = sera positive for IgM-PC or IgM-LC; IgM-PC = sera positive for IgM-PC; LC = lactosylceramide; LIP = lipids; Non-MSMYD = non-MS myelin disease; Non-MYND = nonmyelin neurologic disease; PC = phosphatidylcholine; PP = primary progressive; RR = relapsing-remitting; SP = secondary progressive.

## Discussion


In this work, we report the detection by ELISA of IgM serum antibodies reactive with PC in patients with MS. Microarray-based assays detected lipid-reactive serum antibodies in patients with MS, potentially associated with different disease processes.<sup>13</sup> Serum antibodies to gangliosides have been described in 13–50% of patients with MS and about 40% of patients affected by other autoimmune diseases.<sup>16</sup> IgM antibodies reactive with cholesterol have also been detected in patients with MS, but their roles as biomarkers were not investigated.<sup>17</sup> Our study differs from these reports in that we analyzed serum antibody reactivity directed against antigens different from those previously investigated; in particular, we focused on IgM serum antibodies reactive with PC detected by ELISA. In addition, we analyzed different disease stages and found that most of the patients in the first relapse or during the relapsing-remitting phase present IgM-PC.

Technical problems related to antigen solubility, accessibility of reactive groups, and the requirement of auxiliary lipids are linked to the detection of lipid-reactive antibodies.<sup>17,18</sup> Moreover, the detection of IgG oligoclonal bands is cumbersome, among other reasons because it requires CSF collection, and the sensitivity achieved varies between laboratories, ranging from 91% to 69%.<sup>19</sup> We detected increased serum IgM-PC in most MS patient samples, with a sensitivity comparable to that





**Figure 4** IgM-PC in patients with MS, Non-MSMYD, and Non-MYND and control group



(A) Serum levels of IgM-PC in MS, clinically isolated syndrome, Non-MSMYDs, and Non-MYNDs. Boxes represent the median of OD  $\pm$  percentiles 25–75, and whiskers include 100% of the patients. Percentile 25 = 0.0559. Percentile 75 = 0.2416. (B) Percentage of positive IgM-PC serums in patients with MS, Non-MSMYD, and Non-MYND. Bars represent the percentage of positives or negatives in relation to the total number of individuals analyzed in each group. Dot bars: IgM-PC negative. Crossed bars: IgM-PC positive. The Fisher test was used for comparisons between MS or CIS groups and patients with other neurologic diseases. CIS = clinically isolated syndrome; IgM-PC = IgM to phosphatidylcholine; Non-MSMYD = non-MS myelin disease; Non-MYND = nonmyelin neurologic disease; NS = nonsignificant; OD = optic density.

levels in patients with CIS and RRMS than patients with SPMS and PPMS, suggesting that serum IgM-PC reflects ongoing inflammation and myelin destruction driven by peripheral immune cells. These data are in agreement with a central role for CNS intrinsic mechanisms during the progressive phase of MS.<sup>22,23</sup> Indeed, we previously reported that lipid-reactive IgM OCBs in CSF persist during disease course,<sup>10</sup> and we also detected IgM and IgG antibodies deposited on axons and oligodendrocytes in MS CNS samples,<sup>24</sup> suggesting that these antibodies are produced locally within the CNS, probably in meningeal follicle-like structures.<sup>25–28</sup> Although pathogenic activities have been assigned to some lipid-reactive antibodies,<sup>4</sup> the role of IgM-PC antibodies in MS pathogenesis is still unclear, and they may simply reflect the release of target lipids as a result of myelin destruction.<sup>4,29–31</sup>

PC-reactive IgMs are part of the natural antibody repertoire, which provides protection against multiple pathogens.<sup>32–34</sup> Natural autoantibodies are produced in a T cell-independent manner by B-1 cells<sup>35,36</sup> and usually do not undergo isotype switching to IgG.<sup>37</sup> One mechanism thought to drive the generation of natural antibodies is the ability of their target antigens to trigger signaling via the B-cell receptor (BCR) as well as additional stimulatory pathways in B cells.<sup>38</sup> Because PC is reported to modulate multiple aspects of B-cell biology,<sup>39,40</sup> it is possible that PC released from myelin in the context of inflammation simultaneously activates PC-reactive B-1 cells via the BCR and additional pathways to expand preexisting natural antibody responses. Indeed, increased serum levels of the myelin components GM1 and GD1a are detected in patients with RRMS.<sup>41</sup> Moreover, our finding of increased serum IgM-PC suggests that B-1 cell hyperactivity may also contribute to MS pathology,<sup>42</sup> as

reported in other autoimmune diseases such as rheumatoid arthritis and idiopathic thrombocytopenic purpura.<sup>43</sup> Indeed, the number of B-1 cells in the CSF is linked to poor prognosis in patients with MS.<sup>12</sup>

In conclusion, our results identify IgM-PC antibodies as potential biomarkers for MS. Future studies should extend these observations to additional patient cohorts, analyze the effect of disease-modifying therapies on serum concentration of IgM-PC antibodies, and investigate the role of these immunoglobulins in MS pathogenesis.

### Study funding

This work was supported by grants NS087867, ES02530, AI126880, and AI093903 from the NIH and RSG-14-198-01-LIB from the American Cancer Society to F.J.Q., grant PCON11/2016 from Banco Santander to M.C.S., and grants USP-BS-PPC16/2012 and MEMERG-1 from Instituto de Medicina Molecular Aplicada, Universidad San Pablo-CEU, and PI14/01620 from Banco Santander to U.M. V.R. was supported by a fellowship from the German Research Foundation (DFG RO4866 1/1) and an educational grant from Mallinckrodt Pharmaceuticals (A219074). M.C.S. was supported by a fellowship from the CEU-Banco (XII Convocatoria de ayudas a la movilidad investigadora CEU-BANCO SANTANDER) Santander. C.S. was supported by a fellowship from Fundación Universitaria San Pablo-CEU–Instituto de Medicina Molecular Aplicada. S.E.B. holds the Heidrich Family and Friends endowed chair in Neurology at UCSF. The SUMMIT consortium is supported by NMSS grant RR 2005-A-13 (RG-1510-06785).

### Disclosure

The authors declare no conflict of interest. Go to Neurology.org/NN for full disclosures.

## Publication history

Received by *Neurology: Neuroimmunology & Neuroinflammation* January 28, 2019. Accepted in final form April 9, 2020.

## Appendix Authors

Name	Location	Contribution
<b>María Cruz Sádaba, PhD</b>	Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA; Facultad de Medicina, Instituto de Medicina Molecular Aplicada (INMA), Universidad San Pablo-CEU, CEU Universities, Madrid, Spain	Performed experiments and wrote the manuscript
<b>Veit Rothhammer, MD, PhD</b>	Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA	Performed experiments and wrote the manuscript
<b>Úrsula Muñoz, PhD</b>	Facultad de Medicina, Instituto de Medicina Molecular Aplicada (INMA), Universidad San Pablo-CEU, CEU Universities, Madrid, Spain	Provided unique reagents and discussed and interpreted findings
<b>Cristina Sebal, PhD</b>	Facultad de Medicina, Instituto de Medicina Molecular Aplicada (INMA), Universidad San Pablo-CEU, CEU Universities, Madrid, Spain	Performed experiments
<b>Esther Escudero, PhD</b>	Facultad de Medicina, Instituto de Medicina Molecular Aplicada (INMA), Universidad San Pablo-CEU, CEU Universities, Madrid, Spain	Provided unique reagents and discussed and interpreted findings
<b>Pia Kivisäkk, MD, PhD</b>	Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA	Provided unique reagents and discussed and interpreted findings
<b>María Isabel García Sanchez, PhD</b>	Molecular Biology Service and MS Unit, University of Sevilla, Sevilla, Spain	Provided unique reagents, discussed and interpreted findings
<b>Guillermo Izquierdo, MD, PhD</b>	Molecular Biology Service and MS Unit, University of Sevilla, Sevilla, Spain	Provided unique reagents and discussed and interpreted findings
<b>Stephen L. Hauser, MD</b>	Department of Neurology, University of California, San Francisco	Provided unique reagents and discussed and interpreted findings
<b>Sergio E. Baranzini, PhD</b>	Department of Neurology, University of California, San Francisco	Provided unique reagents and discussed and interpreted findings
<b>Jorge R. Oksenberg, PhD</b>	Department of Neurology, University of California, San Francisco	Provided unique reagents and discussed and interpreted findings

## Appendix (continued)

Name	Location	Contribution
<b>Roberto Álvarez-Lafuente, PhD</b>	Instituto de Investigación Sanitaria San Carlos (IdISSC), Hospital Clínico San Carlos, Madrid, Spain	Provided unique reagents and discussed and interpreted findings
<b>Rohit Bakshi, MD</b>	Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA	Provided unique reagents and discussed and interpreted findings
<b>Howard L. Weiner, MD</b>	Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA	Provided unique reagents and discussed and interpreted findings
<b>Francisco J. Quintana, PhD</b>	Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, MA; The Broad Institute of Harvard and MIT, Cambridge, MA	Provided unique reagents, discussed and interpreted findings, and wrote the manuscript

## References

- Sádaba MC, González Porqué P, Masjuan J, Alvarez-Cermeño JC, Bootello A, Villar LM. An ultrasensitive method for the detection of oligoclonal IgG bands. *J Immunol Methods* 2004;284:141–145.
- Villar LM, Masjuan J, Sádaba MC, et al. Early differential diagnosis of multiple sclerosis using a new oligoclonal band test. *Arch Neurol* 2005;62:574–577.
- O'Brien JS, Sampson EL. Fatty acid and fatty aldehyde composition of the major brain lipids in normal human gray matter, white matter, and myelin. *J Lipid Res* 1965;6:545–551.
- Kanter JL, Narayana S, Ho PP, et al. Lipid microarrays identify key mediators of autoimmune brain inflammation. *Nat Med* 2006;12:138–143.
- Yeste A, Quintana FJ. Antigen microarrays for the study of autoimmune diseases. *Clin Chem* 2013;59:1036–1044.
- Brennan KM, Galban-Horcajo F, Rinaldi S, et al. Lipid arrays identify myelin-derived lipids and lipid complexes as prominent targets for oligoclonal band antibodies in multiple sclerosis. *J Neuroimmunol* 2011;238:87–95.
- Kasai N, Pachner AR, Yu RK. Anti-glycolipid antibodies and their immune complexes in multiple sclerosis. *J Neurol Sci* 1986;75:33–42.
- Quintana FJ, Patel B, Yeste A, et al. Epitope spreading as an early pathogenic event in pediatric multiple sclerosis. *Neurology* 2014;83:2219–2226.
- Quintana FJ, Yeste A, Weiner HL, Covacu R. Lipids and lipid-reactive antibodies as biomarkers for multiple sclerosis. *J Neuroimmunol* 2012;248:53–57.
- Villar L, García-Barragán N, Espiño M, et al. Influence of oligoclonal IgM specificity in multiple sclerosis disease course. *Mult Scler* 2008;14:183–187.
- Villar LM, Masterman T, Casanova B, et al. CSF oligoclonal band patterns reveal disease heterogeneity in multiple sclerosis. *J Neuroimmunol* 2009;211:101–104.
- Villar LM, Sádaba MC, Roldán E, et al. Intrathecal synthesis of oligoclonal IgM against myelin lipids predicts an aggressive disease course in MS. *J Clin Invest* 2005;115:187–194.
- Bakshi R, Yeste A, Patel B, et al. Serum lipid antibodies are associated with cerebral tissue damage in multiple sclerosis. *Neurol Neuroimmunol Neuroinflamm* 2016;3:e200. doi: 10.1212/NXI.0000000000000200.
- Bove R, Chitnis T, Cree BA, et al. SUMMIT (Serially Unified Multicenter Multiple Sclerosis Investigation): creating a repository of deeply phenotyped contemporary multiple sclerosis cohorts. *Mult Scler* 2018;24:1485–1498.
- Gauthier SA, Glanz BI, Mandel M, Weiner HL. A model for the comprehensive investigation of a chronic autoimmune disease: the multiple sclerosis CLIMB study. *Autoimmun Rev* 2006;5:532–536.
- Cross AH, Trotter JL, Lyons J. B cells and antibodies in CNS demyelinating disease. *J Neuroimmunol* 2001;112:1–14.
- Jurewicz A, Domowicz M, Galazka G, Raine CS, Selmaj K. Multiple sclerosis: presence of serum antibodies to lipids and predominance of cholesterol recognition. *J Neurosci Res* 2017;95:1984–1992.
- Ravindranath MH, Muthugounder S, Saravanan TS, Presser N, Morton DL. Human anti-ganglioside autoantibodies: validation of ELISA. *Ann NY Acad Sci* 2005;1050:229–242.
- Schäffler N, Köpke S, Winkler L, et al. Accuracy of diagnostic tests in multiple sclerosis: a systematic review. *Acta Neurol Scand* 2011;124:151–164.



20. Hohmann C, Milles B, Schinke M, et al. Categorization of multiple sclerosis relapse subtypes by B cell profiling in the blood. *Acta Neuropathol Commun* 2014;2:138.
21. Sabatino JJ, Zamvil SS, Hauser SL. B-cell therapies in multiple sclerosis. *Cold Spring Harb Perspect Med* 2019;9:a032037.
22. Thompson AJ, Baranzini SE, Geurts J, Hemmer B, Ciccarelli O. Multiple sclerosis. *Lancet* 2018;391:1622–1636.
23. Baecher-Allan C, Kaskow BJ, Weiner HL. Multiple sclerosis: mechanisms and immunotherapy. *Neuron* 2018;97:742–768.
24. Sádaba MC, Tzartos J, Paíno C, et al. Axonal and oligodendrocyte-localized IgM and IgG deposits in MS lesions. *J Neuroimmunol* 2012;247:86–94.
25. Magliozzi R, Howell O, Vora A, et al. Meningeal B-cell follicles in secondary progressive multiple sclerosis associate with early onset of disease and severe cortical pathology. *Brain* 2007;130:1089–1104.
26. Serafini B, Rosicarelli B, Magliozzi R, Stigliano E, Aloisi F. Detection of ectopic B-cell follicles with germinal centers in the meninges of patients with secondary progressive multiple sclerosis. *Brain Pathol* 2004;14:164–174.
27. Zurawski J, Lassmann H, Bakshi R. Use of magnetic resonance imaging to visualize leptomeningeal inflammation in patients with multiple sclerosis: a review. *JAMA Neurol* 2017;74:100–109.
28. Zurawski J, Tauhid S, Chu R, Khalid F, Healy BC, Weiner HL, Bakshi R. Leptomeningeal enhancement in the brain on 7T MRI scans is common and is associated with cortical lesions in relapsing-remitting multiple sclerosis. *Mult Scler J* 2018;24:88.
29. Farez MF, Quintana FJ, Gandhi R, Izquierdo G, Lucas M, Weiner HL. Toll-like receptor 2 and poly(ADP-ribose) polymerase 1 promote central nervous system neuroinflammation in progressive EAE. *Nat Immunol* 2009;10:958–964.
30. Mayo L, Trauger SA, Blain M, et al. Regulation of astrocyte activation by glycolipids drives chronic CNS inflammation. *Nat Med* 2014;20:1147–1156.
31. Quintana FJ, Farez MF, Vigiotta V, et al. Antigen microarrays identify unique serum autoantibody signatures in clinical and pathologic subtypes of multiple sclerosis. *Proc Natl Acad Sci USA* 2008;105:18889–18894.
32. Corti D, Lanzavecchia A. Broadly neutralizing antiviral antibodies. *Annu Rev Immunol* 2013;31:705–742.
33. Haigwood NL, Stamatatos L. Role of neutralizing antibodies in HIV infection. *AIDS* 2003;17(suppl 4):S67–S71.
34. Ochsenbein AF, Fehr T, Lutz C, et al. Control of early viral and bacterial distribution and disease by natural antibodies. *Science* 1999;286:2156–2159.
35. Berland R, Wortis HH. Origins and functions of B-1 cells with notes on the role of CD5. *Annu Rev Immunol* 2002;20:253–300.
36. Panda S, Ding JL. Natural antibodies bridge innate and adaptive immunity. *J Immunol* 2015;194:13–20.
37. Fagarasan S. T-independent immune response: new aspects of B cell biology. *Science* 2000;290:89–92.
38. Quintana FJ, Solomon A, Cohen IR, Nussbaum G. Induction of IgG3 to LPS via Toll-like receptor 4 co-stimulation. *PLoS One* 2008;3:e3509.
39. Brewer JW, Solodushko V, Aragon I, Barrington RA. Phosphatidylcholine as a metabolic cue for determining B cell fate and function. *Cell Immunol* 2016;310:78–88.
40. Fagone P, Gunter C, Sage CR, Gunn KE, Brewer JW, Jackowski S. CTP:phosphocholine cytidyltransferase alpha is required for B-cell proliferation and class switch recombination. *J Biol Chem* 2009;284:6847–6854.
41. Zaprianova E, Deleva D, Ilinov P, et al. Serum ganglioside patterns in multiple sclerosis. *Neurochem Res* 2001;26:95–100.
42. Li R, Patterson KR, Bar-Or A. Reassessing B cell contributions in multiple sclerosis. *Nat Immunol* 2018;19:696–707.
43. Yamanishi S, Iizumi T, Watanabe E, et al. Implications for induction of autoimmunity via activation of B-1 cells by *Helicobacter pylori* urease. *Infect Immun* 2006;74:248–256.

# Neurology<sup>®</sup> Neuroimmunology & Neuroinflammation

## Serum antibodies to phosphatidylcholine in MS

Maria Cruz Sádaba, Veit Rothhammer, Úrsula Muñoz, et al.

*Neurol Neuroimmunol Neuroinflamm* 2020;7;

DOI 10.1212/NXI.0000000000000765

**This information is current as of June 9, 2020**

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://nn.neurology.org/content/7/4/e765.full.html">http://nn.neurology.org/content/7/4/e765.full.html</a>
<b>References</b>	This article cites 43 articles, 10 of which you can access for free at: <a href="http://nn.neurology.org/content/7/4/e765.full.html##ref-list-1">http://nn.neurology.org/content/7/4/e765.full.html##ref-list-1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Autoimmune diseases</b> <a href="http://nn.neurology.org/cgi/collection/autoimmune_diseases">http://nn.neurology.org/cgi/collection/autoimmune_diseases</a> <b>Multiple sclerosis</b> <a href="http://nn.neurology.org/cgi/collection/multiple_sclerosis">http://nn.neurology.org/cgi/collection/multiple_sclerosis</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://nn.neurology.org/misc/about.xhtml#permissions">http://nn.neurology.org/misc/about.xhtml#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://nn.neurology.org/misc/addir.xhtml#reprintsus">http://nn.neurology.org/misc/addir.xhtml#reprintsus</a>

*Neurol Neuroimmunol Neuroinflamm* is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

