

Amyotrophic Lateral Sclerosis Survival Associates With Neutrophils in a Sex-specific Manner

Benjamin J. Murdock, PhD, Stephen A. Goutman, MD, MS, Jonathan Boss, BS, MS, Sehee Kim, PhD, and Eva L. Feldman, MD, PhD, FAAN, FANA

Correspondence

Dr. Goutman
sgoutman@med.umich.edu

Neurol Neuroimmunol Neuroinflamm 2021;8:e953. doi:10.1212/NXI.0000000000000953

Abstract

Objective

To determine whether neutrophils contribute to amyotrophic lateral sclerosis (ALS) progression, we tested the association of baseline neutrophil count on ALS survival, whether the effect was sex specific, and whether neutrophils accumulate in the spinal cord.

Methods

A prospective cohort study was conducted between June 22, 2011, and October 30, 2019. Blood leukocytes were isolated from ALS participants and neutrophil levels assessed by flow cytometry. Participant survival outcomes were analyzed by groups ($<2 \times 10^6$, $2-4 \times 10^6$, and $>4 \times 10^6$ neutrophils/mL) with adjustments for relevant ALS covariates and by sex. Neutrophil levels were assessed from CNS tissue from a subset of participants.

Results

A total of 269 participants with ALS within 2 years of an ALS diagnosis were included. Participants with baseline neutrophil counts over 4×10^6 /mL had a 2.1 times higher mortality rate than those with a neutrophil count lower than 2×10^6 /mL (95% CI: 1.3–3.5, $p = 0.004$) when adjusting for age, sex, and other covariates. This effect was more pronounced in females, with a hazard ratio of 3.8 (95% CI: 1.8–8.2, $p = 0.001$) in the $>4 \times 10^6$ /mL vs $<2 \times 10^6$ /mL group. Furthermore, ALS participants ($n = 8$) had increased neutrophils in cervical ($p = 0.049$) and thoracic ($p = 0.022$) spinal cord segments compared with control participants ($n = 8$).

Conclusions

Higher neutrophil counts early in ALS associate with a shorter survival in female participants. Furthermore, neutrophils accumulate in ALS spinal cord supporting a pathophysiologic correlate. These data justify the consideration of immunity and sex for personalized therapeutic development in ALS.

Classification of Evidence

This study provides Class III evidence that in female participants with ALS, higher baseline neutrophil counts are associated with shorter survival.

MORE ONLINE**→ Class of Evidence**

Criteria for rating therapeutic and diagnostic studies

[NPub.org/coe](https://www.npub.org/coe)

From the Department of Neurology (B.J.M., S.A.G., E.L.F.), and Department of Biostatistics (J.B., S.K.), School of Public Health, University of Michigan, Ann Arbor.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

ALS = amyotrophic lateral sclerosis; **ALSFRS-R** = ALS Functional Rating Scale, Revised; **FBS** = fetal bovine serum; **PBS** = phosphate-buffered saline.

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease¹ with incompletely understood mechanisms and treatment options. The immune system is a therapeutic target in ALS. Early attempts to treat ALS using global immune suppression were ineffective or exacerbated disease.^{2,3} However, an increasing body of literature suggests that specific immune cell types may have beneficial *or* detrimental effects on disease outcomes^{4,5}; therefore, global immune suppression would suppress beneficial cell function in parallel with destructive function. In response, the next generation of immune-based ALS therapeutic trials is more precise, targeting specific cell populations or immune pathways.

We used our existing immunophenotyping pipeline to explore the potential impact of sex and immunity on ALS survival. Previously, we have shown an association between neutrophil levels and ALS progression,^{6,7} findings supported by Choi et al.⁸ who report a connection between the ratio of neutrophils to lymphocytes and ALS survival. We therefore examined total neutrophil levels and how differences in peripheral neutrophil levels associate with survival in men and women, as there are significant differences in ALS rates between sexes.⁹ As it is still unclear whether peripheral neutrophil changes actively contribute to ALS progression or are simply a marker of disease, we also compared neutrophil levels in postmortem CNS tissue from ALS and control participants to determine whether neutrophils accumulate in the CNS during disease. We also examined whether cellular accumulation is affected by sex.

Methods

Survival Study Participants

All patients seen at the University of Michigan Pranger ALS clinic were invited to participate in this study. Participants provided written informed consent, and the study was approved by the institutional review board. Study protocols were previously published.^{6,7,10,11} Participants with sampling within 2 years of diagnosis were included; furthermore, 2 participants with a prolonged symptom onset to diagnosis interval (9.8 and 15.8 years) were also excluded. Flow cytometry data were obtained between June 22, 2011, and October 20, 2019.

Postmortem Tissue Participants

Persons with ALS seen at the University of Michigan Pranger ALS clinic and control volunteers (no neurologic disease, Alzheimer disease, probable Alzheimer, or possible Parkinson disease) consented to donate postmortem CNS tissue to the University of Michigan Brain Bank.

Enrichment of Leukocytes From Peripheral Blood

Study participants provided blood during clinical encounters. Following peripheral venipuncture, blood was collected into a BD Vacutainer sodium heparin tube (BD Biosciences, San Jose, CA), placed at 4°C, transferred to the laboratory on ice, and processed within 3 hours of collection. One milliliter of whole blood was split into 2 tubes and lysed with red blood cell lysing buffer (0.8% NH₄Cl, 0.098% KHCO₃, 0.1 mM EDTA, and 13.8 mM HEPES) using gentle rocking for 12 minutes. Cells were then pelleted, washed twice with flow buffer (phosphate-buffered saline [PBS], 4% fetal bovine serum [FBS], and 0.1% sodium azide), and counted using a hemocytometer (Hausser Scientific, Horsham, PA).

Isolation of Leukocytes From Human CNS Tissue

Postmortem tissue was collected from ALS and control participants between February 27, 2019, and January 1, 2020. After death postmortem, the spinal cord was divided into cervical, thoracic, and lumbar sections and transferred for laboratory analysis, at which point the immunologist (B.J.M.) was blinded to the diagnosis. Spinal cord tissue was physically dissociated using surgical scissors and enzymatically dissociated with collagenase for 90 minutes with gentle mixing at 15-minute intervals in RPMI-1640 medium (supplemented with 5% FBS [both Thermo Fisher Scientific, Waltham, MA], 50 µg/mL penicillin, 100 µg/mL streptomycin, and 20 mg/mL *Clostridium histolyticum* collagenase [all 3 from Sigma-Aldrich, St. Louis, MO]). The resulting suspension was placed on a 70-µm cell strainer (Corning, Corning, NY) over a 50-mL conical tube (Corning) and dissociated further by grinding with a sterile 3-mL syringe plunger (BD Biosciences), resuspended in 30% stock isotonic Percoll (90% Percoll [GE Healthcare, Chicago, IL] and 10% 10X Hanks' balanced salt solution without Ca²⁺ or Mg²⁺ [Thermo Fisher Scientific]), layered on the top of 70% stock Percoll, and centrifuged at 500g for 30 minutes. After removal of neuronal debris, the resulting interface was collected, washed, resuspended, and counted using a hemocytometer before analysis by flow cytometry.

Flow Cytometry

Human leukocyte suspensions were plated in round bottom 96-well plates (Corning) at $\leq 10^6$ cells/25 µL flow buffer and blocked using 10 µg/mL human TruStain FcX blocking solution (BioLegend, San Diego, CA). Cells were stained for 30 minutes in the dark at 4°C in a 50 µL final volume using a cocktail of fluorescently labeled antibodies against cell surface markers suspended in flow buffer (1X PBS, 2% FBS, and 1% sodium azide). After staining, cells were washed with 200 µL

flow buffer, pelleted, resuspended in 185 μ L of BD stabilizing fixative (BD Biosciences), and transferred for analysis to polystyrene tubes (12 \times 75 mm) (Becton Dickinson, Franklin Lakes, NJ). Cells were analyzed on a BD FACSCanto or LSRFortessa flow cytometer with FACSDiva software (BD Biosciences) and analyzed by FlowJo (FlowJo, Ashland, OR). Fluorophore-conjugated antibodies were CD45-BV421 (BioLegend, catalogue #304032), CD16-PE (BioLegend, catalogue # 302056), CD11b-PerCP-Cy/5.5 (BioLegend, catalogue #301328), control IgG-PE-Cy7 (BioLegend, catalogue # 400125), and CD15-PE-Cy7 (BD, catalogue # 560827). Peripheral neutrophil levels were gated as previously described⁷; CNS neutrophils were gated using CD45, CD11b, and CD15. A nonspecific IgG control was used for CD15 staining to subtract nonspecific events.

Survival Analysis

Descriptive statistics were produced for demographics and ALS disease characteristics. Study population differences were compared between male and female participants and participants with differing neutrophil counts by analysis of variance tests and chi-square tests.

Cox proportional hazards models assessed the association between neutrophil levels and ALS survival endpoints, defined as the time from diagnosis to death. Associations were adjusted for potential confounders, e.g., age, sex, onset segment, El Escorial criteria at diagnosis, and time from symptom onset to diagnosis. A delay in blood sample collection (median duration of 7 months) occurred in some participants, potentially leading to a selection bias if ignored. To account for this gap, a left-truncated survival analysis was performed.¹² To determine a functional form of continuous neutrophil counts, we first fit penalized spline regression models and found a nonlinear effect of neutrophils. Continuous neutrophil counts were categorized into 3 groups (neutrophil counts: <2 , $2-4$, and $>4 \times 10^6$ neutrophils/mL of blood) based on the nonlinear effect of neutrophils identified by penalized spline regression (figure 1). Neutrophil grouping was also confirmed by a sensitivity analysis in which the recategorized neutrophil variables into one-unit-change bins were examined, and bins of similar effect sizes were combined. Linearity of continuous adjustment factors (age and time from symptom onset to diagnosis) was checked, and no violation was found for their linearity assumption. The proportional hazards assumptions were checked using global and individual Schoenfeld tests with graphical assessment of the rescaled Schoenfeld residuals over time, and there were no concerning violations. Estimated, covariate-adjusted survival curves were plotted based on the Cox proportional hazards model using the *survminer* R package following published methods.^{13,14} To perform this method, the data are replicated 6 times (once for each sex by the neutrophil group), and each replicated data set serves as a pseudopopulation where sex by neutrophil groupings are assigned to be the same, keeping all other covariates unchanged. This ensures that plotted survival differences are due to the association with neutrophils and not

a result of imbalances in other clinical factors such as sex, age, and onset segment. Analyses were performed with R version 3.6.2.

CNS Infiltration Analysis

Neutrophil level differences were compared using Prism (GraphPad, San Diego, CA). The Shapiro-Wilk test assessed the normality of distribution. One or more data sets did not adhere to Gaussian distribution in each tissue section; Mann-Whitney was therefore used to assess significance.

Data Availability

Deidentified data will be shared on request from a qualified investigator.


Results

Survival Analysis

To examine the impact of sex on immune-based therapies in ALS, we used flow cytometry to measure neutrophil levels in the peripheral blood of 271 study participants who had provided samples within 2 years of initial diagnosis. Two participants were excluded for missing covariates, one with uncertain onset segment and the other with uncertain onset date (see table 1 for population demographics). Thus, this analysis included 269 participants with a median age of 67.6 years, 45% female, a median diagnostic delay (time from symptom onset to diagnosis) of 1.02 years, median time from symptom onset to flow cytometry of 1.75 years, and a median Revised ALS Functional Rating Scale (ALSFRS-R) of 33. The median time to death was 1.5 years with a median time to censoring of 1.59 years, indicating that observation time was longer in the censored participants compared with those who died. These data indicate the participants were still early in their ALS disease course with preserved functional abilities at study entry. Furthermore, male and female groups were comparable in terms of baseline neutrophil value, age, family history of ALS, race, El Escorial criteria, ALSFRS-R, time from symptom onset to diagnosis, and time from diagnosis and symptom onset to neutrophil collection; the only significant difference was onset segment favoring bulbar onset in females (table 1). When divided by neutrophil levels and sex, there were no significant differences in age, family history of ALS, race, El Escorial criteria, time from symptom onset to diagnosis, and time from diagnosis and symptom onset to neutrophil collection. There was a significant difference in baseline ALSFRS-R scores that were lower in the higher neutrophil group and in onset segment, with females showing a higher proportion of bulbar onset and males showing a higher proportion of cervical onset (table 2). Two separate statistical survival models were constructed to associate peripheral neutrophil levels with participant survival and sex.

In the first survival model (model 1), data from male and female study participants were combined and adjusted for

Figure 1 Estimated Functional Form of the Association Between Neutrophils and Survival



Plots show the estimated functional form of the association between neutrophils and survival (the larger the value on the y axis, the higher the hazard) for (A) all participants (model 1), (B) male participants (model 2), and (C) female participants (model 2). (A) Model 1 shows an overall flat line between the 2×10^6 /mL and 4×10^6 /mL neutrophil counts with an upward sloping line on either side, therefore justifying the cutoffs at $<2 \times 10^6$ /mL, $2-4 \times 10^6$ /mL, and $>4 \times 10^6$ /mL. Similar trends in the functional form of neutrophils and survival were observed for males (B) and females (C). Each tick mark at the bottom of the graph represents a single participant sample. As illustrated, the decreased risk for the highest neutrophil count is driven by 1 participant.

potential confounders such as sex, age, onset segment, El Escorial criteria at diagnosis, and time from symptom onset to diagnosis. Study participants were grouped based on peripheral neutrophil levels: $<2 \times 10^6$ neutrophils/mL of blood, $2-4 \times 10^6$ /mL, and $>4 \times 10^6$ /mL for survival analysis. In the second survival model (model 2), participant data were similarly grouped based on peripheral neutrophil levels, and data were adjusted based on potential cofounders. Rather than adjusting for sex, participants were stratified into male and female groups.

After constructing each model and adjusting for covariates, we examined the association between peripheral neutrophil levels and ALS survival before and after stratifying for sex. In model 1 (combined male and female participants), neutrophil levels were inversely associated with survival: participants with low peripheral neutrophils had the longest survival (median 2.11 years), followed by participants with moderate neutrophil levels (median 1.59 years) and participants with high neutrophil levels (median 1.15 years) (figure 2A). The mortality rate of participants with high neutrophil levels was 2.11 times higher than that of participants with low neutrophil levels ($p = 0.004$; table 3), when adjusting for sex, age, onset segment and other covariates. There was a trend toward reduced survival in participants with a moderate number of peripheral neutrophils compared with low neutrophils not reaching statistical significance. These data indicate that increasing levels of peripheral neutrophils are associated with reduced survival in ALS. Although as we show below, this association was likely driven by sex.

Survival Analysis by Sex

Next, we used the second statistical model where the neutrophil-survival association was stratified by sex. In model 2, median survival times for those with $>4 \times 10^6$ neutrophils/mL were 1.23 years in males and 1.21 years in females (figure 2B). Median survival times for those with $2-4 \times 10^6$ neutrophils/mL were 1.66 years in males and 1.55 years in females. However, the absolute difference in median survival times was most pronounced in those with $<2 \times 10^6$ neutrophils/mL resulting in 1.34 years in males and 3.34 years in females. The 5 of 6 neutrophil and sex groups had a statistically significant higher mortality rate when compared with the one $<2 \times 10^6$ /mL neutrophil female group (table 4) when adjusting for age, onset segment, and other covariates. Similar associations were not seen when comparing the hazard ratio and p value with each of the other neutrophil and sex groups acting as the reference category (table 5). These data suggest that low neutrophil levels have a different impact in males and females.

As effect modification by sex was of particular interest, a likelihood ratio test comparing the marginal (model 1) and interaction (model 2) models assessed how well the data set fit each statistical model. The likelihood ratio test yielded a p value of 0.026, indicating that the sex-based interaction model (model 2) significantly fit the data better. Importantly, these

Table 1 Peripheral Blood Participant Demographics

Covariate	Overall (N = 269)	Female (N = 120)	Male (N = 149)	p Value
Age (y)	67.6 (59.0–74.3)	68.3 (60.8–73.8)	67.5 (56.6–74.9)	0.084
Neutrophils ($\times 10^6$ /mL)	2.87 (2.19–3.85)	2.70 (2.09–4.01)	2.95 (2.22–3.77)	0.884
Last contact event				0.779
Death	160 (59.5)	73 (60.8)	87 (58.4)	
Censored	109 (40.5)	47 (39.2)	62 (41.6)	
Median time to event (y)				
Death	1.5 (1.0–2.37)	1.53 (1.0–2.3)	1.42 (1.0–2.3)	0.469
Censored	1.59 (1.0–2.4)	1.68 (1.3–2.2)	1.44 (0.9–2.5)	0.870
Family history of ALS				0.759
No	227 (84.4)	102 (85.0)	125 (83.9)	
Yes	28 (10.4)	14 (11.7)	14 (9.4)	
Missing	14 (5.2)	4 (3.3)	10 (6.7)	
Race				NA
African American	5 (1.9)	2 (1.7)	3 (2.0)	
White	254 (94.4)	116 (96.7)	138 (92.6)	
Other	3 (1.1)	0 (0.0)	3 (2.0)	
Missing	7 (2.6)	2 (1.7)	5 (3.4)	
El Escorial criteria				0.114
Possible/suspected	46 (17.1)	14 (11.7)	32 (21.5)	
Probable, LS	74 (27.5)	31 (25.8)	43 (28.9)	
Probable	86 (32.0)	43 (35.8)	43 (28.9)	
Definite	63 (23.4)	32 (26.7)	31 (20.8)	
Onset segment				<0.001
Bulbar	82 (30.5)	50 (41.7)	32 (21.5)	
Cervical	94 (34.9)	24 (20.0)	70 (47.0)	
Lumbar	93 (34.6)	46 (38.3)	47 (31.5)	
ALSFRS-R ^a	33.0 (27.0–38.0)	34.0 (26.0–37.0)	32.0 (27.0–38.0)	0.587
Time between symptom onset and neutrophil measure (y)	1.75 (1.20–2.52)	1.79 (1.33–2.33)	1.66 (1.12–2.66)	0.288
Time between diagnosis and neutrophil measure (y)	0.58 (0.35–0.94)	0.60 (0.34–0.95)	0.57 (0.37–0.94)	0.916
Time between symptom onset and diagnosis (y)	1.02 (0.69–1.67)	1.06 (0.78–1.46)	1.01 (0.62–1.85)	0.189

Abbreviations: ALS = amyotrophic lateral sclerosis; ALSFRS-R = Amyotrophic Lateral Sclerosis Functional Rating Scale, Revised; LS = lab supported. Table of descriptive statistics for the ALS cases. For continuous variables, median (25th–75th percentile), and for categorical variables, n (%). *p* Values for continuous and categorical variables correspond to analysis of variance tests and χ^2 tests, respectively.

^a Closest ALSFRS-R score to neutrophil measurement. Participants without an ALSFRS-R score within 1 month of neutrophil measurement were not included in this calculation (N = 250).

models showed no violation of proportional hazards with the global Schoenfeld test, yielding *p* values of 0.29 and 0.14 for the marginal and interaction models, respectively. Thus, both models are valid, but model 2—which stratifies by sex—is more accurate.

CNS Neutrophil Levels in ALS

Finally, we examined whether neutrophils accumulate in the CNS during ALS and whether CNS neutrophil levels were affected by sex. We measured neutrophil levels in CNS postmortem tissue of 8 control and 8 ALS participants.

Table 2 Peripheral Blood Participant Demographics by Neutrophil Group

Covariate	Neutrophils 0-2		Neutrophils 2-4		Neutrophils >4		p Value
	Female (N = 26)	Male (N = 30)	Female (N = 64)	Male (N = 89)	Female (N = 30)	Male (N = 30)	
Age (y)	68.2 (62.4-73.5)	59.8 (51.5-72.2)	68.0 (60.8-73.3)	67.6 (59.9-75.7)	68.3 (60.9-74.2)	69.6 (57.8-73.0)	0.275
Time to event (y)							
Death	2.21 (1.62-2.73)	1.11 (0.77-1.93)	1.45 (0.98-1.98)	1.65 (1.15-2.53)	1.67 (0.87-2.19)	1.34 (0.85-2.18)	0.058
Censored	1.62 (1.29-3.26)	2.03 (0.88-3.03)	1.84 (1.28-2.25)	1.30 (0.89-2.36)	1.42 (0.95-1.51)	1.83 (1.34-2.98)	0.773
Last contact event							0.008
Death	10 (38.5)	15 (50.0)	39 (60.9)	49 (55.1)	24 (80.0)	23 (76.7)	
Censored	16 (61.5)	15 (50.0)	25 (39.1)	40 (44.9)	6 (20.0)	7 (23.3)	
Education							NA
HS/GED or less	5 (19.2)	8 (26.7)	17 (26.6)	17 (19.1)	12 (40.0)	7 (23.3)	
Some college/associate's	5 (19.2)	7 (23.3)	21 (32.8)	25 (28.1)	11 (36.7)	6 (20.0)	
Bachelor's	5 (19.2)	6 (20.0)	7 (10.9)	21 (23.6)	4 (13.3)	4 (13.3)	
Master's/professional	8 (30.8)	2 (6.7)	8 (12.5)	12 (13.5)	1 (3.3)	4 (13.3)	
Missing	3 (11.5)	7 (23.3)	11 (17.2)	14 (15.7)	2 (6.7)	9 (30.0)	
Family history of ALS							NA
No	25 (96.2)	25 (83.3)	56 (87.5)	72 (80.9)	21 (70.0)	28 (93.3)	
Yes	1 (3.9)	3 (10.0)	6 (9.4)	9 (10.1)	7 (23.3)	2 (6.7)	
Missing	0 (0.0)	2 (6.7)	2 (3.1)	8 (9.0)	2 (6.7)	0 (0.0)	
Race							NA
African American	2 (7.7)	2 (6.7)	0 (0.0)	1 (1.1)	0 (0.0)	0 (0.0)	
White	23 (88.5)	25 (83.3)	63 (98.4)	84 (94.4)	30 (100.0)	29 (96.7)	
Other	0 (0.0)	2 (6.7)	1 (1.6)	1 (1.1)	0 (0.0)	0 (0.0)	
Missing	1 (3.9)	1 (3.3)	0 (0.0)	3 (3.4)	0 (0.0)	1 (3.3)	
Ethnicity							NA
Hispanic/Latino	0 (0.0)	1 (3.3)	0 (0.0)	2 (2.3)	0 (0.0)	0 (0.0)	
Not Hispanic/Latino	24 (92.3)	29 (96.7)	62 (96.9)	85 (95.5)	30 (100.0)	30 (100.0)	
Missing	2 (7.7)	0 (0.0)	2 (3.1)	2 (2.3)	0 (0.0)	0 (0.0)	
El Escorial criteria							NA
Possible/suspected	7 (26.9)	6 (20.0)	4 (6.3)	17 (19.1)	3 (10.0)	9 (30.0)	
Probable, LS	6 (23.1)	11 (36.7)	18 (28.1)	27 (30.3)	7 (23.3)	5 (16.7)	
Probable	9 (34.6)	9 (30.0)	25 (39.1)	25 (28.1)	9 (30.0)	9 (30.0)	
Definite	4 (15.4)	4 (13.3)	17 (26.6)	20 (22.5)	11 (36.7)	7 (23.3)	
Onset segment							<0.001
Bulbar	10 (38.5)	4 (13.3)	30 (46.9)	23 (25.8)	10 (33.3)	5 (16.7)	
Cervical	4 (15.4)	17 (56.7)	15 (23.4)	40 (44.9)	5 (16.7)	13 (43.3)	
Lumbar	12 (46.2)	9 (30.0)	19 (29.7)	26 (29.2)	15 (50.0)	12 (40.0)	

Continued

Table 2 Peripheral Blood Participant Demographics by Neutrophil Group (continued)

Covariate	Neutrophils 0-2		Neutrophils 2-4		Neutrophils >4		p Value
	Female (N = 26)	Male (N = 30)	Female (N = 64)	Male (N = 89)	Female (N = 30)	Male (N = 30)	
ALSFRS-R ^a	37.0 (35.0–40.0)	36 (29.5–41.0)	33.0 (26.5–36.5)	32.0 (27.0–36.8)	30.0 (22.0–35.0)	29.5 (23.8–35.2)	<0.001
Time between symptom onset and neutrophil measure (y)	1.80 (1.36–2.12)	1.54 (1.17–2.56)	1.71 (1.33–2.36)	1.77 (1.12–2.77)	2.13 (1.34–2.50)	1.43 (1.12–2.47)	0.733
Time between diagnosis and neutrophil measure (y)	0.57 (0.35–0.84)	0.44 (0.30–0.83)	0.63 (0.35–0.95)	0.62 (0.37–0.98)	0.59 (0.35–1.25)	0.59 (0.45–0.75)	0.735
Time between symptom onset and diagnosis (y)	1.09 (0.78–1.24)	0.94 (0.63–1.85)	1.10 (0.82–1.45)	1.07 (0.60–1.92)	0.98 (0.70–1.62)	0.97 (0.68–1.65)	0.686

Abbreviations: ALS = amyotrophic lateral sclerosis; ALSFRS-R = ALS Functional Rating Scale, Revised; HS = high school; LS = lab supported. Table of descriptive statistics for the ALS cases. For continuous variables, median (25th–75th percentile), and for categorical variables, n (%). *p* Values for continuous and categorical variables correspond to analysis of variance tests and χ^2 tests, respectively. NA indicates that cell counts were too small to perform the χ^2 test.

^a Closest ALSFRS-R score to neutrophil measurement. Participants without an ALSFRS-R score within 1 month of neutrophil measurement were not included in this calculation (N = 250).

Mean death age for cases and controls was 72.5 and 85.6 years, respectively. ALS cases were 50% male and controls were 100% male. Final neuropathologic diagnosis in all cases was ALS and, in controls, was Alzheimer 12.5%, probable Alzheimer 50%, possible Parkinson 12.5%, and no diagnosis 25%. We previously found that neutrophils accumulate in the spinal cord of ALS mice,¹⁵ so in human participants we examined neutrophil levels in 3 distinct spinal cord sections (cervical, thoracic, and lumbar). When assessed using flow cytometry, we detected increased neutrophil levels in the spinal cord of ALS compared with control participants (figure 3A). When quantified, mean neutrophil counts were significantly increased in the cervical ($p = 0.049$) and thoracic ($p = 0.022$) sections of ALS participants compared with control participants (figure 3B). However, significant differences in neutrophil levels between ALS male and female participants were not observed (figure 3C), indicating that neutrophils accumulate in the CNS of ALS participants without a clear difference between sexes.


Discussion

We demonstrate that the immune system has differing effects in male and female ALS participants and that sex should be accounted for during the development of ALS therapeutics and early phase clinical trials, particularly those focused on the immune system. Our analysis was performed by associating peripheral neutrophil levels with participant survival. Two separate survival models examined the potential role of neutrophils in ALS and to determine whether sex affects immunity in ALS. When data from male and female study participants were combined, we found increasing levels of peripheral neutrophils were associated with increased mortality. However, when stratified by sex, we found that neutrophil levels had drastically different associations with ALS mortality.

These observations are of particular importance given the recent interest in ALS immune factors and the growing number of clinical trials targeting the immune system.

Our first statistical model combined data from male and female participants to elucidate the overall impact of neutrophils in ALS, to set a baseline of comparison for the second statistical model, and to ensure our analysis was consistent with previously published studies. When participants were grouped by peripheral neutrophil levels, we found that increasing peripheral neutrophils were associated with increased mortality in ALS. These data are consistent with our previous studies that have linked neutrophil levels to disease progression,⁶ linked neutrophil levels to disease progression rates,⁷ and have demonstrated CNS accumulation in an ALS mouse model.¹⁵ They are also highly consistent with a recent study, which found that a high ratio of neutrophils to lymphocytes was associated with reduced survival in ALS.⁸ Furthermore, they are consistent with another recent study showing that higher neutrophil levels associate with a faster disease progression as measured by the change in ALSFRS-R.¹⁶ Although to date these studies have been correlative, other studies have demonstrated a central role for neutrophils in neurologic damage. Neutrophils are some of the first responders to neurologic damage,¹⁷ and although they can facilitate repair in the spinal cord, they are equally capable of inducing damage.¹⁸ For example, in MS, neutrophils facilitate further damage by activating other immune cell types,¹⁹ and targeting neutrophils attenuates disease.^{19,20} The accumulation of neutrophils in the CNS supports their involvement in disease progression, although the underlying mechanism or mechanisms driving neutrophil accumulation in the CNS of ALS participants remain unclear. Multiple proinflammatory cytokines are upregulated in the plasma of ALS participants,^{21–23} and these cytokines could be increasing neutrophil activity and trafficking. In addition, other immune factors linked with ALS can also affect neutrophil function. Leukotriene B4, platelet-activating factor, and C5a, for

Figure 2 Neutrophil Impact on ALS Survival in Male and Female Participants



Estimated survival curves using 2 separate models for analysis. (A) In model 1, survival curves were adjusted for age, sex, onset segment, El Escorial criteria at diagnosis, and time between symptom onset and diagnosis. Study ALS participants were categorized into 3 neutrophil groups ($< 2 \times 10^6$, $2-4 \times 10^6$, and $> 4 \times 10^6$ neutrophils/mL of blood). Survival curves were generated by creating a pseudopopulation with identical population characteristics as the study participants for each neutrophil group and calculating their expected survival based on model 1 (table 3). Dashed lines indicate the median survival for each neutrophil group. (B) In model 2, female and male survival curves were displayed separately after adjusting for age, onset segment, El Escorial criteria at diagnosis, and time between symptom onset and diagnosis were estimated from the interaction model. Male and female data sets were categorized into 3 neutrophil groups per sex ($< 2 \times 10^6$, $2-4 \times 10^6$, and $> 4 \times 10^6$ neutrophils/mL of blood). Survival curves were generated by creating a pseudopopulation with identical population characteristics as the study participants for each neutrophil and sex group and calculating their expected survival based on model 2 (table 4). Dashed lines indicate the median survival for each neutrophil and sex group. ALS = amyotrophic lateral sclerosis.

example, have all been associated with ALS^{24–27} and can enhance the activity, trafficking, and survival of neutrophils.^{28,29}

However, although our data and previous studies suggest neutrophils may directly contribute to increased mortality

during ALS, our second statistical model suggests that the role of neutrophils is complicated by sex. When the neutrophil-survival association was stratified by sex, we found that low peripheral neutrophil levels had completely different effects in males and females. In female participants, low peripheral

Table 3 Model 1: Cox Proportional Hazard Survival Model for All Participants

Term	HR	95% CI	p Value
Neutrophils $2-4 \times 10^6$/mL	1.403	0.882–2.229	0.152
Neutrophils $>4 \times 10^6$/mL	2.112	1.278–3.493	0.004
Male	1.169	0.829–1.648	0.373
Onset segment			
Cervical	0.660	0.433–1.007	0.054
Lumbar	0.740	0.490–1.117	0.152
Age (y)	1.026	1.010–1.041	0.001
El Escorial criteria			
Definite	2.778	1.626–4.745	<0.001
Probable	1.515	0.915–2.511	0.107
Probable, LS	1.194	0.703–2.027	0.513
Time between symptom onset and diagnosis (y)	0.803	0.668–0.965	0.019

Abbreviations: HR = hazard ratio; LS = lab supported.

Estimated median survival time for individuals with neutrophils $< 2 \times 10^6$ /mL is 2.11 years, estimated median survival time for those with neutrophils $2-4 \times 10^6$ /mL is 1.59 years, and estimated median survival time for those with neutrophils $> 4 \times 10^6$ /mL is 1.15 years.

Table 4 Model 2: Cox Proportional Hazard Survival Model by Sex

Term	HR	95% CI	p Value
Female, neutrophils 2–4 × 10 ⁶ /mL	2.685	1.299–5.549	0.008
Female, neutrophils >4 × 10 ⁶ /mL	3.829	1.777–8.249	0.001
Male, neutrophils <2 × 10 ⁶ /mL	3.420	1.445–8.092	0.005
Male, neutrophils 2–4 × 10 ⁶ /mL	2.561	1.267–5.174	0.009
Male, neutrophils >4 × 10 ⁶ /mL	3.755	1.758–8.022	0.001
Onset segment			
Cervical	0.630	0.410–0.967	0.035
Lumbar	0.785	0.518–1.188	0.252
Age (y)	1.029	1.013–1.045	<0.001
EI Escorial criteria			
Definite	2.495	1.452–4.287	0.001
Probable	1.366	0.819–2.280	0.232
Probable, LS	1.042	0.609–1.784	0.881
Time between symptom onset and diagnosis (y)	0.782	0.651–0.944	0.010

Abbreviations: HR = hazard ratio; LS = lab supported.

Estimated median survival time for females with neutrophils between <2 × 10⁶/mL is 3.34 years, estimated median survival time for females with neutrophils 2–4 × 10⁶/mL is 1.55 years, estimated median survival time for females with neutrophils >4 × 10⁶/mL is 1.21 years, estimated median survival time for males with neutrophils <2 × 10⁶/mL is 1.34 years, estimated median survival time for males with neutrophils 2–4 × 10⁶/mL is 1.66 years, and estimated median survival time for males with neutrophils >4 × 10⁶/mL is 1.23 years.


neutrophil levels are associated with a longer median survival. By contrast, low neutrophil levels in male participants did not associate with increased survival. The mechanism for this discrepancy is currently not clear, although several possibilities exist. The first is the direct effects of sex hormones on cellular activity: males and females display altered immune activation states in multiple cell types,^{30–32} including neutrophils,^{33–35} because of the immunomodulatory effects of male and female hormones.³⁶ A second possibility is that sex-based differences contribute to altered neutrophil levels or altered CNS infiltration. This is

unlikely, however, as we saw no differences in peripheral or spinal cord neutrophil levels between male and female ALS participants. A third possibility is that sex fundamentally alters the immune environment within the CNS. Previous data support this last possibility. In rodent models, for example, the microglia of healthy males and females have different transcriptomic signatures,³⁷ and hormone levels create different immune environments during sexual development.^{38,39} The ability of neutrophils to initiate tissue repair or drive further damage has also been linked to the local microenvironment.⁴⁰

Table 5 Hazard Ratios and p Values Corresponding to Different Reference Categories in the Final Cox Proportional Hazards Model

	Reference (Ref) group					
	Female, 0–2	Female, 2–4	Female, >4	Male, 0–2	Male, 2–4	Male, >4
	HR (p value)	HR (p value)	HR (p value)	HR (p value)	HR (p value)	HR (p value)
Female, 0–2	Ref	0.372 (0.008)	0.261 (<0.001)	0.292 (0.005)	0.391 (0.009)	0.266 (<0.001)
Female, 2–4	2.685 (0.008)	Ref	0.701 (0.185)	0.785 (0.456)	1.049 (0.832)	0.715 (0.235)
Female, >4	3.829 (<0.001)	1.426 (0.185)	Ref	1.120 (0.756)	1.495 (0.133)	1.020 (0.950)
Male, 0–2	3.420 (0.005)	1.274 (0.456)	0.893 (0.756)	Ref	1.336 (0.349)	0.911 (0.790)
Male, 2–4	2.561 (0.009)	0.954 (0.832)	0.669 (0.133)	0.749 (0.349)	Ref	0.682 (0.145)
Male, >4	3.756 (<0.001)	1.399 (0.235)	0.981 (0.950)	1.098 (0.790)	1.467 (0.145)	Ref

Figure 3 Accumulation of Neutrophils in the CNS During ALS



(A) Spinal cord tissue from control and ALS participants (cervical sections, collected postmortem) was analyzed for neutrophils using flow cytometry. (B) Total neutrophils were quantified in postmortem tissue of the cervical, thoracic, and lumbar spinal cord sections from control and ALS participants. Mean neutrophil counts were: cervical cord, ALS = $10.9 \times 10^3/\text{g}$, control = $1.6 \times 10^3/\text{g}$, $p = 0.049$; thoracic cord, ALS = $3.8 \times 10^3/\text{g}$, control = $0.7 \times 10^3/\text{g}$, $p = 0.002$; and lumbar cord, ALS = $8.4 \times 10^3/\text{g}$, control = $1.5 \times 10^3/\text{g}$, $p = 0.194$. (C) Neutrophil accumulation in the CNS of male and female ALS study participants is compared in the cervical, thoracic, and lumbar sections of the spinal cord. Mean neutrophil counts were: cervical cord, males $12.2 \times 10^3/\text{g}$, females $9.7 \times 10^3/\text{g}$, $p = 0.686$; thoracic cord, males $4.9 \times 10^3/\text{g}$, females $2.6 \times 10^3/\text{g}$, $p = 0.343$; lumbar cord, males $10.6 \times 10^3/\text{g}$, females $6.2 \times 10^3/\text{g}$, $p = 0.314$. * $p < 0.05$. ALS = amyotrophic lateral sclerosis; FSC = forward scatter.

Although the underlying mechanism remains to be determined, our data demonstrate that sex should be considered when examining the role of the immune system in ALS. Although few ALS studies have specifically examined the effect of sex on disease, others have detected immune discrepancies as well. For instance, 1 study examining the role of fractalkine receptor function in the microglia of ALS mice found that knockout of the receptor accelerated disease progression only in male mice.⁴¹ Another study showed that axonal sprouting varies between male and female SOD1^{G37R} mice.⁴² A more recent prospective study found that

prediagnostic eosinophil levels associated with ALS risk differently in males and females.⁴³ Finally, low-density lipoprotein-related receptor protein 4 antibodies are more commonly detected in female vs male ALS participants.⁴⁴ Combined with our observations, these studies suggest that multiple immune cell types, not just neutrophils, are affected by sex during ALS. We contend that any ALS study addressing immunity should account for sex.

This is particularly important because the ALS field becomes increasingly interested in immune-based therapies. A recent

phase I trial, for instance, isolated regulatory T cells through leukapheresis and expanded the cell population *ex vivo* before reintroducing them into the trial participant.⁴⁵ In addition, the upcoming HEALEY ALS Platform Trial will include immunomodulatory drugs targeting complement or myeloperoxidase.⁴⁶ Despite this new focus on immunity in ALS, sex-specific treatments are frequently not addressed, and sex was not specified as an important variable in the most recent ALS clinical trial guideline revisions⁴⁷ despite higher proportions of ALS in men than in women, particularly in younger study participants.⁹ Our data suggest that separate immune mechanisms may exist in men and women and that future immune-based ALS studies should account for sex.

These findings should not be surprising given that autoimmune diseases, such as systemic lupus erythematosus and MS, disproportionately affect females.³⁶ Furthermore, there are inherent differences in neutrophil activity between males and females,^{33–35} and neutrophil extracellular traps differ between males and females with relapsing remitting MS.⁴⁸ Overall, sex and immune system differences are well understood in non-neurologic disease and MS, and our data suggest the same may be true for ALS.

Our study does have limitations. It is primarily correlative, as it did not use any intervention to assess the impact of neutrophils on disease progression. Similarly, the accumulation of neutrophils in the CNS could be a byproduct of neuronal damage, rather than a contributing factor. Peripheral neutrophil levels were not measured in participants until they visited the ALS clinic, meaning participants in the earliest stages of observable disease who were yet to be diagnosed were missed; however, the immune system may also be involved before observable symptoms. When participants were divided by neutrophil count (table 2), the group with the highest neutrophils did have a lower ALSFRS-R score. Importantly, there were no differences in (1) time from symptom onset to neutrophil measure; (2) time between diagnosis and neutrophil measure; and (3) time between symptom onset and diagnosis. Therefore, we believe these ALSFRS-R differences by subgroup reflect the severity of this group as opposed to capturing these participants later in disease. All 8 control participants who donated CNS tissue were male and older, on average, than ALS participants, meaning that innate immune differences between the sexes or based on age were not accounted for in the CNS analysis. Although a strength of our study is the use of an inclusive clinic population, in contrast to a clinical trial population that largely excludes patients,⁴⁹ replication of our findings in other cohorts is needed to address generalizability.

In summary, the heterogeneous nature of ALS complicates the identification of therapeutic targets and the outcomes of early phase clinical studies.⁵⁰ However, despite the well-established differences in ALS rates between males and females,⁹ few studies take sex into consideration as an independent variable. Our current data indicate that not only do

differences in immunity exist between males and females during ALS but also that these differences link to different disease outcomes. Our observations suggest that immunomodulatory therapies for the treatment of ALS should be assessed separately by sex in both preclinical and early phase studies to better account for patient-specific immune signatures. Such a shift moves us closer to more personalized therapeutics for the treatment of ALS.

Acknowledgment

Blake Swihart, BA, MA, Jayna Duell, BS, RN, Amanda Williams, BS, Thomas Dent, BS, Adam Patterson, BS, Crystal Pacut, BS, Matthew Perkins, MD, Joshua Famie, BS, Caroline Piecuch, BS, Faye Mendelson, BS, Shayna Mason, Maegan Tabbey, BS, and Carey Backus, BA, University of Michigan, provided technical support. Stacey Sakowski Jacoby, PhD, and Masha G. Savelieff, PhD, University of Michigan, provided editing assistance. These contributors receive a salary for their work but were not directly compensated for this specific project.

Study Funding

This study was supported by R21 NS102960, the University of Michigan NeuroNetwork for Emerging Therapies, the Robert Nederlander Program, the A. Alfred Taubman Medical Research Institute, the University of Michigan MCubed program, contract N021258-00 from Target ALS, contract NS102960 from the NIH, contract 200-2013-56856 from the Centers for Disease Control and Prevention/Agency for Toxic Substances and Disease Registry, the NIH (NIH K23ES027221), and contract N021258 from the CReATe Consortium (the CReATe consortium [U54NS092091] is part of the Rare Diseases Clinical Research Network [RDCRN], an initiative of the Office of Rare Diseases Research [ORDR], and the National Center for Advancing Translational Sciences [NCATS]). CReATe is funded through a collaboration between the NCATS and the NINDS).

Disclosure

No authors declare conflicts of interest relevant to this manuscript. Go to Neurology.org/NN for full disclosures.

Publication History

Received by *Neurology: Neuroimmunology & Neuroinflammation* August 13, 2020. Accepted in final form November 13, 2020.

Appendix Authors

Name	Location	Contribution
Benjamin J. Murdock, PhD	University of Michigan, Ann Arbor	Design and conceptualized the study; analyzed the data; and drafted the manuscript for intellectual content
Stephen A. Goutman, MD, MS	University of Michigan, Ann Arbor	Design and conceptualized the study; analyzed the data; and drafted the manuscript for intellectual content

Continued

Appendix (continued)

Name	Location	Contribution
Jonathan Boss, BS, MS	University of Michigan, Ann Arbor	Interpreted the data and revised the manuscript for intellectual content
Sehee Kim, PhD	University of Michigan, Ann Arbor	Interpreted the data and revised the manuscript for intellectual content
Eva L. Feldman, MD, PhD, FAAN, FANA	University of Michigan, Ann Arbor	Design and conceptualized the study; analyzed the data; and revised the manuscript for intellectual content

References

- Goutman SA. Diagnosis and clinical management of amyotrophic lateral sclerosis and other motor neuron disorders. *Continuum (Minneapolis)* 2017;23:1332–1359.
- Gordon PH, Moore DH, Miller RG, et al. Efficacy of minocycline in patients with amyotrophic lateral sclerosis: a phase III randomised trial. *Lancet Neurol* 2007;6:1045–1053.
- Meininger V, Asselain B, Guillet P, et al. Pentoxifylline in ALS: a double-blind, randomized, multicenter, placebo-controlled trial. *Neurology* 2006;66:88–92.
- Murdock BJ, Bender DE, Segal BM, Feldman EL. The dual roles of immunity in ALS: injury overrides protection. *Neurobiol Dis* 2015;77:1–12.
- Zhao W, Beers DR, Appel SH. Immune-mediated mechanisms in the pathogenesis of amyotrophic lateral sclerosis. *J Neuroimmune Pharmacol* 2013;8:888–899.
- Murdock BJ, Bender DE, Kashlan SR, et al. Increased ratio of circulating neutrophils to monocytes in amyotrophic lateral sclerosis. *Neuro Neuroimmunol Neuroinflamm* 2016;3:e242.
- Murdock BJ, Zhou T, Kashlan SR, Little RJ, Goutman SA, Feldman EL. Correlation of peripheral immunity with rapid amyotrophic lateral sclerosis progression. *JAMA Neurol* 2017;74:1446–1454.
- Choi SJ, Hong YH, Kim SM, Shin JY, Suh YJ, Sung JJ. High neutrophil-to-lymphocyte ratio predicts short survival duration in amyotrophic lateral sclerosis. *Sci Rep* 2020;10:428.
- Manjaly ZR, Scott KM, Abhinav K, et al. The sex ratio in amyotrophic lateral sclerosis: a population based study. *Amyotroph Lateral Scler* 2010;11:439–442.
- Goutman SA, Boss J, Patterson A, Mukherjee B, Batterman S, Feldman EL. High plasma concentrations of organic pollutants negatively impact survival in amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 2019;90:907–912.
- Su FC, Goutman SA, Chernyak S, et al. Association of environmental toxins with amyotrophic lateral sclerosis. *JAMA Neurol* 2016;73:803–811.
- Wu F, Kim S, Qin J, Saran R, Li Y. A pairwise likelihood augmented Cox estimator for left-truncated data. *Biometrics* 2018;74:100–108.
- Alimujiang A, Wiensch A, Boss J, et al. Association between life purpose and mortality among US adults older than 50 years. *JAMA Netw Open* 2019;2:e194270.
- Therneau TM, Crowson CS, Atkinson EJ. Adjusted survival curves. 2015.
- Figuerola-Romero C, Guo K, Murdock BJ, et al. Temporal evolution of the microbiome, immune system and epigenome with disease progression in ALS mice. *Dis Model Mech* 2019;13:dmm041947.
- Woodruff TM, McCombe PA, Henderson RD, et al. Monocytes and neutrophils are associated with clinical features in amyotrophic lateral sclerosis. *Brain Commun* 2020;2:fcaa013.
- Kim CF, Moalem-Taylor G. Detailed characterization of neuro-immune responses following neuropathic injury in mice. *Brain Res* 2011;1405:95–108.
- Neirinckx V, Coste C, Franzen R, Gothot A, Rogister B, Wislet S. Neutrophil contribution to spinal cord injury and repair. *J Neuroinflammation* 2014;11:150.
- Steinbach K, Piedavent M, Bauer S, Neumann JT, Friese MA. Neutrophils amplify autoimmune central nervous system infiltrates by maturing local APCs. *J Immunol* 2013;191:4531–4539.
- Khaw YM, Cunningham C, Tierney A, Sivaguru M, Inoue M. Neutrophil-selective deletion of Cxcr2 protects against CNS neurodegeneration in a mouse model of multiple sclerosis. *J Neuroinflammation* 2020;17:49.
- Aebischer J, Moumen A, Sazdovitch V, Seilhean D, Meininger V, Raoul C. Elevated levels of IFN γ and LIGHT in the spinal cord of patients with sporadic amyotrophic lateral sclerosis. *Eur J Neurol* 2012;19:752–759, e745–e756.
- Cereda C, Baiocchi C, Bongioanni P, et al. TNF and sTNFR1/2 plasma levels in ALS patients. *J Neuroimmunol* 2008;194:123–131.
- Rentzos M, Rombos A, Nikolaou C, et al. Interleukin-15 and interleukin-12 are elevated in serum and cerebrospinal fluid of patients with amyotrophic lateral sclerosis. *Eur Neurol* 2010;63:285–290.
- Malaspina A, Puentes F, Amor S. Disease origin and progression in amyotrophic lateral sclerosis: an immunology perspective. *Int Immunol* 2015;27:117–129.
- Woodruff TM, Costantini KJ, Crane JW, et al. The complement factor C5a contributes to pathology in a rat model of amyotrophic lateral sclerosis. *J Immunol* 2008;181:8727–8734.
- Lee JD, Levin SC, Willis EF, Li R, Woodruff TM, Noakes PG. Complement components are upregulated and correlate with disease progression in the TDP-43(Q331K) mouse model of amyotrophic lateral sclerosis. *J Neuroinflammation* 2018;15:171.
- Briones MRS, Snyder AM, Ferreira RC, Neely EB, Connor JR, Broach JR. A possible role for platelet-activating factor receptor in amyotrophic lateral sclerosis treatment. *Front Neurol* 2018;9:39.
- Webster RO, Hong SR, Johnston RB Jr, Henson PM. Biological effects of the human complement fragments C5a and C5ades Arg on neutrophil function. *Immunopharmacology* 1980;2:201–219.
- Khreiss T, Jozsef L, Chan JS, Filep JG. Activation of extracellular signal-regulated kinase couples platelet-activating factor-induced adhesion and delayed apoptosis of human neutrophils. *Cell Signal* 2004;16:801–810.
- Hou J, Zheng WF. Effect of sex hormones on NK and ADCC activity of mice. *Int J Immunopharmacol* 1988;10:15–22.
- Retzew JA, Huet-Hudson YM, Marriott I. Testosterone reduces macrophage expression in the mouse of toll-like receptor 4, a trigger for inflammation and innate immunity. *Biol Reprod* 2008;78:432–437.
- Potluri T, Fink AL, Sylvia KE, et al. Age-associated changes in the impact of sex steroids on influenza vaccine responses in males and females. *NPJ Vaccin* 2019;4:29.
- Buyon JP, Korchak HM, Rutherford LE, Ganguly M, Weissmann G. Female hormones reduce neutrophil responsiveness in vitro. *Arthritis Rheum* 1984;27:623–630.
- Miller AP, Feng W, Xing D, et al. Estrogen modulates inflammatory mediator expression and neutrophil chemotaxis in injured arteries. *Circulation* 2004;110:1664–1669.
- Marin DP, Bolin AP, dos Santos Rde C, Curi R, Otton R. Testosterone suppresses oxidative stress in human neutrophils. *Cell Biochem Funct* 2010;28:394–402.
- Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol* 2016;16:626–638.
- Villa A, Gelosa P, Castiglioni L, et al. Sex-specific features of microglia from adult mice. *Cell Rep* 2018;23:3501–3511.
- Lenz KM, Nugent BM, Haliyur R, McCarthy MM. Microglia are essential to masculinization of brain and behavior. *J Neurosci* 2013;33:2761–2772.
- Lenz KM, Wright CL, Martin RC, McCarthy MM. Prostaglandin E(2) regulates AMPA receptor phosphorylation and promotes membrane insertion in preoptic area neurons and glia during sexual differentiation. *PLoS One* 2011;6:e18500.
- Butterfield TA, Best TM, Merrick MA. The dual roles of neutrophils and macrophages in inflammation: a critical balance between tissue damage and repair. *J Athl Train* 2006;41:457–465.
- Cardona AE, Pioro EP, Sasse ME, et al. Control of microglial neurotoxicity by the fractalkine receptor. *Nat Neurosci* 2006;9:917–924.
- Martineau E, Di Polo A, Vande Velde C, Robitaille R. Sex-specific differences in motor-unit remodeling in a mouse model of ALS. *eNeuro* 2020;7:ENEURO.0388-19.2020.
- Yazdani S, Mariosa D, Hammar N, et al. Peripheral immune biomarkers and neurodegenerative diseases: a prospective cohort study with 20 years of follow-up. *Ann Neurol* 2019;86:913–926.
- Rivner MH, Liu S, Quarles B, et al. Agrin and low-density lipoprotein-related receptor protein 4 antibodies in amyotrophic lateral sclerosis patients. *Muscle Nerve* 2017;55:430–432.
- Thonhoff JR, Beers DR, Zhao W, et al. Expanded autologous regulatory T-lymphocyte infusions in ALS: a phase I, first-in-human study. *Neuro Neuroimmunol Neuroinflamm* 2018;5:e465.
- HEALEY ALS Platform Trial—Master Protocol. Available at: ClinicalTrials.gov/show/NCT04297683. Accessed May 19, 2020.
- van den Berg LH, Sorenson E, Gronseth G, et al. Revised Airlie House consensus guidelines for design and implementation of ALS clinical trials. *Neurology* 2019;92:e1610–e1623.
- Tillack K, Naegele M, Haeus C, et al. Gender differences in circulating levels of neutrophil extracellular traps in serum of multiple sclerosis patients. *J Neuroimmunol* 2013;261:108–119.
- van Eijk RPA, Westeneng HJ, Nikolakopoulos S, et al. Refining eligibility criteria for amyotrophic lateral sclerosis clinical trials. *Neurology* 2019;92:e451–e460.
- Goyal NA, Berry JD, Windebank A, et al. Addressing heterogeneity in amyotrophic lateral sclerosis CLINICAL TRIALS. *Muscle Nerve* 2020;62:156–166.

Neurology[®] Neuroimmunology & Neuroinflammation

Amyotrophic Lateral Sclerosis Survival Associates With Neutrophils in a Sex-specific Manner

Benjamin J. Murdock, Stephen A. Goutman, Jonathan Boss, et al.
Neurol Neuroimmunol Neuroinflamm 2021;8;
DOI 10.1212/NXI.0000000000000953

This information is current as of February 2, 2021

Updated Information & Services	including high resolution figures, can be found at: http://nn.neurology.org/content/8/2/e953.full.html
References	This article cites 48 articles, 8 of which you can access for free at: http://nn.neurology.org/content/8/2/e953.full.html##ref-list-1
Citations	This article has been cited by 2 HighWire-hosted articles: http://nn.neurology.org/content/8/2/e953.full.html##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Amyotrophic lateral sclerosis http://nn.neurology.org/cgi/collection/amyotrophic_lateral_sclerosis_
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://nn.neurology.org/misc/about.xhtml#permissions
Reprints	Information about ordering reprints can be found online: http://nn.neurology.org/misc/addir.xhtml#reprintsus

Neurol Neuroimmunol Neuroinflamm is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2021 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

