

# Assessing Neurofilaments as Biomarkers of Neuroprotection in Progressive Multiple Sclerosis

From the MS-STAT Randomized Controlled Trial

Thomas E. Williams, BA, MB BChir, MRCP, Katherine P. Holdsworth, MSc, Jennifer M. Nicholas, PhD, Arman Eshaghi, PhD, Theodora Katsanouli, MSc, Henrietta Wellington, PhD, FRCP, Amanda Heslegrave, PhD, Henrik Zetterberg, PhD, Chris Frost, PhD, and Jeremy Chataway, PhD, FRCP

**Correspondence**  
Dr. Williams  
thomas.e.williams@ucl.ac.uk

*Neurol Neuroimmunol Neuroinflamm* 2022;9:e1130. doi:10.1212/NXI.0000000000001130

## Abstract

### Background and Objectives

Improved biomarkers of neuroprotective treatment are needed in progressive multiple sclerosis (PMS) to facilitate more efficient phase 2 trial design. The MS-STAT randomized controlled trial supported the neuroprotective potential of high-dose simvastatin in secondary progressive MS (SPMS). Here, we analyze serum from the MS-STAT trial to assess the extent to which neurofilament light (NfL) and neurofilament heavy (NfH), both promising biomarkers of neuroaxonal injury, may act as biomarkers of simvastatin treatment in SPMS.

### Methods

The MS-STAT trial randomized patients to 80 mg simvastatin or placebo. Serum was analyzed for NfL and NfH using Simoa technology. We used linear mixed models to investigate the treatment effects of simvastatin compared with placebo on NfL and NfH. Additional models examined the relationships between neurofilaments and MRI and clinical measures of disease severity.

### Results

A total of 140 patients with SPMS were included. There was no evidence for a simvastatin treatment effect on NfL or NfH: compared with placebo, NfL was 1.2% lower (95% CI 10.6% lower to 9.2% higher;  $p = 0.820$ ) and NfH was 0.4% lower (95% CI 18.4% lower to 21.6% higher;  $p = 0.969$ ) in the simvastatin treatment group. Secondary analyses suggested that higher NfL was associated with greater subsequent whole brain atrophy, higher T2 lesion volume, and more new/enlarging T2 lesions in the previous 12 months, as well as greater physical disability. There were no significant associations between NfH and MRI or clinical variables.

### Discussion

We found no evidence of a simvastatin treatment effect on serum neurofilaments. While confirmation of the neuroprotective benefits of simvastatin is awaited from the ongoing phase 3 study (NCT03387670), our results suggest that treatments capable of slowing the rate of whole brain atrophy in SPMS, such as simvastatin, may act via mechanisms largely independent of neuroaxonal injury, as quantified by NfL. This has important implications for the design of future phase 2 clinical trials in PMS.

### Trial Registration Information

MS-STAT: NCT00647348.

## MORE ONLINE

**Class of Evidence**  
Criteria for rating therapeutic and diagnostic studies  
[NPub.org/coe](http://NPub.org/coe)

From the Queen Square Multiple Sclerosis Centre (T.E.W., A.E., J.C.), Department of Neuroinflammation, UCL Queen Square Institute of Neurology, Faculty of Brain Sciences, University College London; National Hospital of Neurology and Neurosurgery (T.E.W., J.C.), London; London School of Hygiene and Tropical Medicine (K.P.H., J.M.N., T.K., C.F.); and UK Dementia Research Institute at UCL (H.W., A.H., H.Z.), United Kingdom.

Go to [Neurology.org/NN](http://Neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

## Glossary

**9HPT** = 9-hole peg test; **25FW** = 25-foot timed walk; **CoV** = coefficient of variation; **EDSS** = Expanded Disability Status Scale; **IQR** = interquartile range; **LLoQ** = lower limit of quantification; **MS** = multiple sclerosis; **NfH** = neurofilament heavy; **NfL** = neurofilament light; **PMS** = progressive MS; **SPMS** = secondary progressive MS; **T2LV** = T2 lesion volume.

## Classification of Evidence

This study provides class I evidence that simvastatin treatment does not have a large impact on either serum NfL or NfH, as quantified in this study, in SPMS.

Few treatments are available for progressive multiple sclerosis (PMS), with most restricted to those with active inflammatory disease. Treatments for nonactive PMS<sup>1</sup> may require an alternative approach, utilizing neuroprotective mechanisms alone or in combination with immunomodulation. One such candidate neuroprotective treatment is simvastatin, which in the MS-STAT trial demonstrated a 43% reduction in annualized brain atrophy, in addition to benefits on secondary physical and cognitive outcomes. No differences were noted on a panel of immunologic markers, supporting a neuroprotective rather than immunomodulatory mechanism.<sup>2,3</sup> The phase 3 MS-STAT2 study is ongoing (NCT03387670).

Blood neurofilament light (NfL) is a biomarker of neuroaxonal injury and an appealing surrogate outcome measure for trials in relapsing-remitting MS.<sup>4</sup> It is less well studied in PMS.<sup>5</sup> Initiation of immunosuppressive disease-modifying treatment is associated with a reduction in blood NfL in PMS cohorts, but an important question remains over the ability of NfL to capture the treatment effect of purportedly neuroprotective therapies.<sup>6-9</sup> Clarifying this is an important goal of the progressive MS alliance, with implications on future phase 2 trial design.<sup>10</sup> Data for blood neurofilament heavy (NfH) are more limited, but it may have utility as a biomarker of neuroprotection.<sup>11,12</sup>

Here, we analyze serum NfL and NfH from the MS-STAT trial. We use the known positive effect of simvastatin on whole brain atrophy to interrogate our primary research question: whether serum NfL and NfH may act as biomarkers of neuroprotective treatment with simvastatin in secondary progressive MS (SPMS). We additionally perform further analyses to examine the relationships between serum neurofilaments and established MRI and clinical variables.

## Methods

### MS-STAT Trial and Sample Processing

The MS-STAT study protocol has been outlined previously.<sup>2</sup> Briefly, patients with SPMS, aged 18–65 years with Expanded Disability Status Scale (EDSS) score 4.0–6.5, were eligible. Key exclusion criteria included primary progressive MS; relapse or steroid use within 3 months; and the use of immunosuppressive or disease-modifying therapy within the last 6 months. A total of

140 patients were randomized, 1:1, to simvastatin 80 mg or placebo. Baseline characteristics are shown in Table 1. Blood samples were acquired at baseline and months 6, 12, and 24. Serum was separated and stored at –80°C until time of analysis.

### Neurofilament Quantification

NfL and NfH are components of a neurone-specific intermediate filament, differing in their C-terminal domain and phosphorylation state. Both are released following neuroaxonal injury into the CSF and blood, where they may be quantified. Serum NfL and NfH were measured by Simoa technology on a HD-1 analyzer, according to the manufacturer's instructions (Quanterix, Billerica, MA). The Simoa NF-Light Advantage and Simoa pNF-heavy Discovery Kits (Quanterix) were used. Briefly, serum samples were thawed at 21°C, vortexed, and centrifuged at 10,000 RCF for 5 minutes at 21°C. On-board the HD-1, samples were diluted 1:4 with sample diluent and bound to paramagnetic beads coated with a capture antibody specific for human NfL or NfH. Antibody-coated beads were incubated with a biotinylated anti-NfL or anti-NfH detection antibodies, that in turn were labeled with a streptavidin- $\beta$ -galactosidase complex. Following the addition of the  $\beta$ -galactosidase substrate resorufin  $\beta$ -D-galactopyranoside, a fluorescent signal proportional to the concentration of neurofilament present in the sample was generated in the antigen-containing microwells of the Simoa plates.

Duplicate measurements were taken of each sample. Sample concentrations were extrapolated from a standard curve, fitted using a 4-parameter logistic algorithm. The lower limit of quantification (LLoQ) for NfL is 0.174 pg/mL and for NfH is 2.88 pg/mL.<sup>13,14</sup> Values below the LLoQ were assigned the value of half the LLoQ. The coefficient of variation (CoV) between sample replicates tends to be higher for lower value results. To avoid bias, all data were therefore included in the primary statistical analysis regardless of the CoV. Each assay was run in the same or consecutive batches by the same operator, who was blinded to treatment allocation.

### MRI Processing

The imaging data have been previously published and were acquired as previously described.<sup>2,15</sup> Briefly, 3D T1-weighted, double-echo proton density, and T2-weighted MRI was obtained at baseline, month 12, and month 25. Whole brain

**Table 1** Characteristics of the MS-STAT Trial Cohort and Descriptive Statistics for Serum NfL and Serum NfH Across Time Points

	Placebo (n = 70)	Simvastatin (n = 70)	All (N = 140)
<b>Baseline characteristics</b>			
Sex, n (%), female	48 (69)	49 (70)	97 (69)
Ethnicity, n (%), White	63 (90)	69 (99)	132 (94)
Relapse in last 24 mo, n (%)	18 (26)	8 (11)	26 (19)
Age, y, mean (SD)	51.1 (6.8)	51.5 (7.0)	51.3 (6.9)
MS duration, y, mean (SD)	20.3 (8.8)	22.1 (8.3)	21.2 (8.6)
SPMS duration, y, mean (SD)	7.1 (4.8)	7.3 (5.6)	7.2 (5.2)
EDSS score, median (IQR)	6 (5.5–6.5)	6 (5.5–6.5)	6 (5.5–6.5)
Previous use of interferon, n (%)	12 (17)	10 (14)	22 (16)
<b>Serum NfL, pg/mL</b>			
Baseline, median (IQR)	15.3 (10.2–22.2)	13.9 (10.9–19.7)	14.6 (10.8–20.2)
N	63	65	128
Month 6, median (IQR)	14.8 (12.0–23.0)	15.0 (11.5–21.8)	14.8 (11.7–22.6)
N	53	59	112
Month 12, median (IQR)	16.6 (12.1–22.3)	16.8 (13.3–23.1)	16.7 (12.8–22.9)
N	53	59	112
Month 24, median (IQR)	16.9 (12.4–22.6)	16.0 (11.7–21.4)	16.0 (11.9–22.2)
N	48	64	112
<b>Serum NfH, pg/mL</b>			
Baseline, median (IQR)	64.2 (24.0–136.0)	67.4 (23.9–116.0)	65.5 (24.0–118.5)
N	63	62	125
Month 6, median (IQR)	71.4 (25.7–135.4)	58.0 (22.7–112.1)	62.5 (22.7–121.9)
N	49	53	102
Month 12, median (IQR)	66.5 (22.9–135.5)	67.2 (25.2–107.1)	67.0 (25.2–113.6)
N	52	57	109
Month 24, median (IQR)	71.9 (26.6–111.8)	60.0 (27.0–125.6)	69.7 (27.0–119.5)
N	48	64	112

Abbreviations: EDSS = Expanded Disability Status Scale; IQR = interquartile range; MS = multiple sclerosis; NfH = neurofilament heavy; NfL = neurofilament light; SPMS = secondary progressive MS.

atrophy was determined using the boundary shift integral method and expressed as percentage change in whole brain volume. T2 new/enlarging lesions were expressed as a count and T2 lesion volume (T2LV) in milliliters.

### Statistical Analysis

The prespecified primary analysis was to examine the effect of simvastatin (80 mg) vs placebo on levels of serum NfL at 24 months. The primary analysis was conducted on the intention to treat population regardless of treatment adherence. An exploratory analysis was undertaken to examine the treatment effect using a

per-protocol data set, which included patients who complied with treatment and completed follow-up to 25 months. Participants were considered compliant with treatment if they reported taking, on average, at least 90% of their tablets at the protocol dose of 2 tablets per day.

The prespecified secondary analysis examined the relationship between serum NfL and whole brain atrophy rate. Further analyses of the association of NfL with other MRI and clinical variables and all analyses of NfH data were exploratory. Neurofilament data were skewed, and hence, all analyses were

**Table 2** Effect of High-Dose Simvastatin on Serum NfL and Serum NfH

	Difference in geometric mean (%) <sup>a</sup>	95% CI	p Value
<b>Serum NfL (pg/mL)</b>			
Month 6	-3.29	-16.2 to 11.6	0.646
Month 12	5.35	-7.4 to 19.9	0.429
Month 24	-5.21	-17.4 to 8.8	0.448
Mean treatment effect	-1.16	-10.6 to 9.2	0.820
<b>Serum NfH (pg/mL)</b>			
Month 6	-7.22	-27.9 to 19.4	0.560
Month 12	4.74	-15.1 to 29.3	0.666
Month 24	1.69	-18.4 to 26.7	0.881
Mean treatment effect	-0.39	-18.4 to 21.6	0.969

Abbreviations: EDSS = Expanded Disability Status Scale; IQR = interquartile range; NfH = neurofilament heavy; NfL = neurofilament light.  
<sup>a</sup> Percentage difference in geometric mean serum neurofilaments, simvastatin vs placebo, adjusted for age, sex, dichotomized EDSS scores (4.0–5.5, 6.0–6.5) and study site. Differences are shown for each follow-up visit and also the average treatment effect across all follow-up visits.

performed following log<sub>2</sub> transformation. Clinical data included EDSS scores, 25-foot timed walk (25FW) expressed as speed (inverse of completion time in seconds), and 9-hole peg test (9HPT) expressed as a speed (1,000 × inverse of completion time in seconds). Analyses were conducted in Stata 15.1 or later.

### Analysis of Simvastatin Treatment Effect on NfL and NfH

A mixed effect model was used to estimate the simvastatin treatment effect on serum neurofilaments at 6, 12, and 24 months, for NfL and NfH. A single model was used to estimate a separate treatment effect at each visit (6, 12, and 24 months) by including a categorical variable for visit and an interaction between visit and treatment group. Baseline neurofilament data were included as an additional end point, but with the treatment effect here constrained to be 0. This is essentially equivalent to adjusting for the baseline neurofilament level.<sup>16</sup> The model included an unstructured residual covariance matrix for the residuals (hence allowing a different variance at each visit and different covariances between each pair of measurements on the same participant). The following baseline variables, which were used as minimization factors in the randomization, were adjusted for by including them and their interactions with visit as fixed effects: age, sex, dichotomized EDSS scores (4.0–5.5, 6.0–6.5), and study site. The mean of the estimated treatment effects across all follow-up visits is also presented. In addition, an exploratory analysis was conducted which adjusted for the minimization factors in the randomization and the following baseline variables: T2LV

and number of relapses in the previous 24 months. In these models, we did not perform adjustments for covariates measured after randomization as they may be influenced by the treatment allocation and hence introduce bias into the analysis of treatment effect.<sup>17</sup>

### Analysis Relating NfL and NfH to MRI and Clinical Variables


This analysis used data on baseline neurofilament levels and the rates of change in neurofilament levels during the trial as predictors of MRI and clinical outcomes. A 2-stage analysis was performed.

In the first stage, a summary measure for rate of change in NfL and NfH for each participant was calculated as the slope from a simple linear regression model relating each participant's repeated measures of log NfL, and separately log NfH, to time from baseline using ordinary least squares.<sup>18</sup> In the second stage, each participant's estimated rate of change in log NfL or log NfH was used as a predictor variable, along with baseline-centered log NfL or log NfH, in a series of separate models for MRI and clinical outcomes. For each outcome, a separate model was fitted for log NfL and log NfH.

The model for whole brain atrophy was an extension of a previously described linear mixed model for directly measured change between each pair of MRI visits (baseline to 12, baseline to 25, and 12–25 months) as the outcome.<sup>19</sup> It included participant-level random slopes for time between scans and random effects for visit. All predictor variables were included as interactions with time between scans in order to model the associations with atrophy rate. The model included baseline neurofilaments and change in neurofilaments, as well as the following baseline variables: age, sex, MRI site, SPMS duration, relapse in the previous 24 months, previous use of interferon, and baseline treatments for fatigue, depression, neuropathic pain, spasticity, and bladder urgency.

The model for T2LV was a linear mixed model with T2LV at each MRI visit (baseline, 12 months, and 25 months) as the outcome. Analysis of 25FW and 9HPT used a linear mixed model with speed at each visit (baseline, 12 months, and 24 months) as the outcome. These models included participant level random slopes for change over time and random intercepts, allowing for correlation between these random effects. In each model, baseline log neurofilament was included as a predictor, along with its interaction with time since baseline. Analysis of T2LV additionally included change in log neurofilament and its interaction with time. The models for T2LV included the same variables that were adjusted for in the whole brain atrophy model on their own as well as an interaction with time. The models for clinical variables included these same adjustments, with the exception of MRI site. As T2LV and clinical variables violated normality assumptions, inference from these models is based on nonparametric bias-corrected and accelerated 95% and 99% CIs calculated from 10,000 bootstrap replications clustered on participant. *p*

**Figure 1** Estimated Mean Serum NfL (A) and Serum NfH (B) in Simvastatin and Placebo Treatment Groups



Values are marginal adjusted geometric means from linear mixed models with treatment effect at baseline constrained to 0. There was no evidence for a treatment effect on either NfL or NfH. NfH = neurofilament heavy; NfL = neurofilament light.

values are therefore not calculated for these models, but the ranges in which they lie can be inferred from the CIs.

For the EDSS score, the linear mixed model did not converge. Instead, a linear regression model for score at each visit (baseline, 12 months, and 24 months) was used. To allow for the nonindependence of measures from the same participant, nonparametric bias-corrected and accelerated CIs clustered on participant were used as described above. The models for the EDSS score included the same adjustment variables as for the other clinical outcomes.

Although T2LV may reflect overall disease burden, the identification of active lesions (either gadolinium-enhancing lesions on a single scan or new/enlarging T2 lesions when comparing 2 time points) is the key MRI measure of ongoing neuroinflammation.<sup>20</sup> The MS-STAT cohort did not include gadolinium-enhanced imaging, and new/enlarging T2 lesions cannot be determined at baseline. To further explore the known relationship between serum NfL and neuroinflammation in this SPMS cohort, we therefore performed additional exploratory linear regression modeling using month 24 log NfL as the dependent variable. This allowed inclusion of recent active lesions (new/enlarging T2 lesions during month 0 to month 12 and month 12 to month 25) and concurrent T2LV (month 25) as predictors. Models were fitted including each of the MRI variables on their own and then together in a mutually adjusted multivariable model. These models included the same covariates as the T2LV models.

### Standard Protocol Approvals, Registrations, and Patient Consents

The study was conducted in accordance with Good Clinical Practice and the Declaration of Helsinki.<sup>21</sup> The MS-STAT protocol was approved by each study site's institutional review board and a national ethics committee; all patients gave written informed consent before entering the study; and ethical approval for the retrospective analysis of serum

samples was received. The MS-STAT clinical trial identification number is NCT00647348.

### Data Availability

Anonymized NfL and NfH data are provided as a supplementary data file ([links.lww.com/NXI/A679](https://links.lww.com/NXI/A679)).

## Results

### NfL and NfH Data

Data on NfL were available from at least 1 visit for 138 patients (69 in each treatment group), with 128 patients having NfL data from at least 1 follow-up visit (61 placebo; 67 simvastatin). For NfH, data were available from 137 patients (69 placebo; 68 simvastatin), with 127 patients having NfH data from at least 1 follow-up visit (60 placebo; 67 simvastatin). No NfL measures were below the LLoQ, and all sample replicates had a CoV <20%. For NfH, 8 samples (1.8%) were below the LLoQ and 39 samples (8.6%) had a CoV >20%. At baseline, median NfL was 14.6 pg/mL (interquartile range [IQR] 10.8–20.2 pg/mL), and median NfH was 65.5 pg/mL (IQR 24.0–118.5 pg/mL) (Table 1). Characteristics of the per-protocol dataset are included in eTable 1 ([links.lww.com/NXI/A679](https://links.lww.com/NXI/A679)).

### Analysis of Simvastatin Treatment Effect on NfL and NfH

There was no evidence of a simvastatin treatment effect on either NfL or NfH at any time point (Table 2), with adjusted marginal mean NfL and NfH levels being similar in the 2 treatment groups at each follow-up visit (Figure 1). Taking the mean of the treatment effects across all follow-up time points, the geometric mean NfL was 1.2% lower in the simvastatin group than in the placebo group (95% CI 10.6% lower to 9.2% higher;  $p = 0.820$ ), whereas the geometric mean NfH was 0.4% lower in the simvastatin group than on placebo

**Table 3** Relationship Between Serum NfL, NfH, and Imaging Variables

	Parameter estimate	95% CI	p Value
<b>Whole brain atrophy</b>			
<b>Relationship with rate of whole brain atrophy (% per year)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	0.207	0.072 to 0.341	0.003
Rate of change in NfL (doublings per year <sup>a</sup> )	0.093	-0.201 to 0.387	0.535
Baseline NfH (per doubling)	0.048	-0.005 to 0.102	0.078
Rate of change in NfH (doublings per year <sup>a</sup> )	0.016	-0.105 to 0.136	0.795
<b>T2 lesion volume</b>			
<b>Relationship with baseline T2 lesion volume (mL)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	7.68	3.66 to 12.48	<0.01
Rate of change in NfL (doublings per year <sup>a</sup> )	15.08	6.67 to 26.40	<0.01
Baseline NfH (per doubling)	0.736	-0.921 to 2.631	>0.05
Rate of change in NfH (doublings per year <sup>a</sup> )	0.413	-6.696 to 8.001	>0.05
<b>Relationship with change in T2 lesion volume (mL/y)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	0.29	-0.14 to 0.63	>0.05
Rate of change in NfL (doublings per year <sup>a</sup> )	0.39	-0.60 to 1.02	>0.05
Baseline NfH (per doubling)	-0.070	-0.218 to 0.051	>0.05
Rate of change in NfH (doublings per year <sup>a</sup> )	-0.032	-0.535 to 0.541	>0.05

Abbreviations: NfH = neurofilament heavy; NfL = neurofilament light.

The results of 4 separate models are presented, with whole brain atrophy or T2 lesion volume as the dependent variables and NfL or NfH data as the predictor variables. In all analyses, neurofilament data were log<sub>2</sub> transformed. Results are from covariate adjusted models as indicated in the Methods section. For T2 lesion volume, the p value bounds (</>0.05 and </>0.01) can only be inferred from the 95% and 99% bias-corrected and accelerated cluster bootstrap (10,000 replications) CIs.

<sup>a</sup> A 1-unit increase in the number of doublings per year corresponds to a change from stable levels to a doubling per year or from doubling once every year to doubling every 6 months (2 doublings per year).

(95% CI 18.4% lower to 21.6% higher;  $p = 0.969$ ). The results from the exploratory per-protocol analysis were similar to those found for the intention-to-treat analysis, with no evidence of a simvastatin treatment effect on either NfL or NfH at any time point (eTable 2, [links.lww.com/NXI/A679](https://links.lww.com/NXI/A679)).

Sensitivity analyses found that the results were not materially altered following the exclusion of 2 individuals with outlying neurofilament values. In addition, an exploratory analysis found that results were essentially unchanged with adjustment for baseline lesion volume and relapses within the last 24 months (eTable 3, [links.lww.com/NXI/A679](https://links.lww.com/NXI/A679)).


### Association of NfL and NfH With MRI Variables

The relationships between both NfL and NfH and each of whole brain atrophy and T2LV are shown in Table 3 and Figure 2. There was evidence for an association between higher baseline NfL and faster whole brain atrophy rate and

between higher baseline NfL and greater T2LV: a twofold increase in baseline NfL was associated with a 0.21%/year increase in brain atrophy and with a 7.7 mL higher baseline T2LV. The rate of change in NfL was not associated with the rate of change in T2LV or the rate of brain atrophy. Patients with a greater increase in NfL from baseline to month 24, however, tended to have a higher T2LV at baseline and follow-up time points. As there was little effect of change in NfL on T2LV, the effect was similar across all visits: an increase of 1 extra doubling per year in NfL was associated with a 15.1 mL higher baseline T2LV and 15.9 mL higher month 25 T2LV. There was no evidence for an association between NfH and any MRI variables (Table 3; Figure 2).

NfL at month 24 was associated with concurrent T2LV and new/enlarging T2 lesions between baseline and month 12 and between months 12 and 25 when examined in separate models (Table 4). When these were combined together as

**Figure 2** Relationship Between Each of Baseline Serum NfL and NfH and Each of Whole Brain Atrophy Rate and Baseline T2 Lesion Volume



Points represent individual patient data. Whole brain atrophy rate is reported as yearly % change from baseline to month 25 and baseline T2 lesion volume in milliliters. (A) Baseline NfL and baseline to month 25 whole brain atrophy rate. (B) Baseline NfH and baseline to month 25 whole brain atrophy rate. (C) Baseline NfL and baseline T2 lesion volume. (D) Baseline NfH and baseline T2 lesion volume. NfH = neurofilament heavy; NfL = neurofilament light.

predictors in the same mutually adjusted model, both concurrent T2LV and month 12–25 new/enlarging T2 lesions remained independently associated with month 24 NfL, whereas the association with new and enlarging lesions between baseline and month 12 was lost. In the final model, month 24 NfL was increased by 5.7% for each new/enlarging T2 lesion between months 12 and 25 and by 0.89% for each milliliter increase in month 25 T2LV.

### Association of NfL and NfH With Clinical Variables

Higher baseline NfL was significantly associated with higher baseline EDSS score, but not with the rate of change in the EDSS score from baseline to 2 years (Table 5). Higher baseline NfL was also associated with worse baseline 9HPT performance and with a greater rate of worsening in the 25FW speed from baseline to 2 years. Baseline NfH was not materially associated with any clinical variables. Sensitivity analyses demonstrated that the results were not materially changed following the exclusion of 2 neurofilament outliers.

### Classification of Evidence

This study assessed the ability of serum NfL and NfH to act as biomarkers of a neuroprotective treatment response with high-dose simvastatin, compared with placebo, in patients with SPMS. It provides Class I evidence that these biomarkers, as quantified in this study, do not act as biomarkers of neuroprotection with simvastatin.

### Discussion

Our results demonstrate that despite simvastatin reducing the annualized whole brain atrophy rate by 43% per year, compared with placebo, we did not find evidence to support a simvastatin treatment effect on serum NfL or NfH. We also replicate previously observed findings, demonstrating that higher NfL is associated with a greater subsequent rate of whole brain atrophy and that recent inflammatory activity (new/enlarging T2 lesions), as well as T2LV, is associated with higher NfL.

The existing literature suggests that in MS, the degree of neuroaxonal injury reflected by serum NfL is predominantly

**Table 4** MRI Predictors of Month 24 Serum NfL

Predictor variable	Separate model for each T2 lesion variable			Combined, mutually adjusted, model		
	% Increase in 24 mo NfL per unit increase in predictor	95% CI	p Value	% increase in 24 mo NfL per unit increase in predictor	95% CI	p Value
Month 0–12 T2 new/enlarging lesions (count)	3.23	0.29 to 6.26	0.031	-0.15	-3.08 to 2.86	0.919
Month 12–25 T2 new/enlarging lesions (count)	6.76	3.45 to 10.18	<0.001	5.73	2.40 to 9.17	0.001
Month 25 T2 lesion volume (mL)	1.10	0.51 to 1.67	<0.001	0.89	0.29 to 1.49	0.004

Abbreviation: NfL = neurofilament light.

Results are presented from the 3 separate models, and then from the combined linear regression model, with month 24 NfL out the outcome and T2 lesion variables as predictors. Results are from covariate adjusted models as indicated in the Methods section. As previously, NfL was log<sub>2</sub> transformed. T2 new/enlarging lesions are reported as a count and T2 lesion volume in milliliters. Coefficients are expressed as % increase in 24 month NfL per unit increase in T2 lesion variables.

related to ongoing neuroinflammation. Such neuroinflammation may be detected by conventional MRI measures, such as recent T1 gadolinium-enhancing lesions or new T2 lesions, or by advanced MRI measures of chronic neuroinflammation, such as the identification of chronic active lesions via their paramagnetic rims.<sup>5,22</sup> Our previous findings suggested that simvastatin treatment was not systemically immunomodulating in this cohort, hence providing 1 possible rationale for the absence of a treatment effect on NfL.<sup>2</sup>

Many of the pathophysiologic mechanisms contributing to progressive MS ultimately converge on neuroaxonal damage, which may be reflected by increased NfL.<sup>23</sup> The dissociation between the previously observed benefits of simvastatin (on whole brain atrophy and disability measures) and the absence of a treatment effect on NfL, however, does also highlight the potential importance of additional processes, independent of neuroaxonal injury, in the pathophysiology of SPMS. Comorbidities, particularly cardiovascular, are prevalent within the MS population and are known to have an impact on future disability and brain atrophy.<sup>24–26</sup> MRI measures of brain atrophy have been validated against clinical treatment effects, long-term disability outcomes, and measures of neuroaxonal loss.<sup>27–30</sup> Brain atrophy is, however, not specific to neuroaxonal injury and may be influenced by volume changes in other CNS tissue compartments. The mechanism of action of simvastatin in progressive MS is the subject of an ongoing mechanistic vascular perfusion study (OPT-MS, NCT03896217), and the ultimate confirmation of the efficacy of simvastatin in SPMS awaits the results of the ongoing phase 3 MS-STAT2 trial (NCT03387670). Our data therefore suggest that caution is required when considering NfL as an outcome measure for treatments in progressive MS if the mechanism of action is not known to directly affect neuroaxonal injury, such as with simvastatin.

Although our results are not necessarily generalizable to other neuroprotective treatments in PMS, they are supported by data from ibudilast. The SPRINT-MS study demonstrated a significant 48% reduction in the rate of brain atrophy in PMS

with ibudilast compared with placebo.<sup>31</sup> There was, however, no significant difference between the treatment groups in either serum or CSF NfL.<sup>9</sup> Although ibudilast is likely to have pleiotropic effects (such as modulation of CNS innate immunity through inhibition of phosphodiesterases, macrophage migration inhibitory factor, and Toll-like receptor 4), like simvastatin, it is not thought to be systemically immunomodulatory.<sup>32</sup>

NfL has shown utility as a biomarker of treatment with fingolimod, siponimod, natalizumab, and ocrelizumab in PMS cohorts.<sup>6–8</sup> These treatments all share a predominantly immunomodulatory mechanism of action, and their ability to reduce NfL is therefore entirely in keeping with the known association between NfL, neuroaxonal injury, and markers of inflammatory activity in PMS.<sup>5</sup> Supporting this, subgroup analyses suggest that there may be a greater treatment effect of natalizumab, siponimod, and ocrelizumab on NfL in patients with recent inflammatory activity.<sup>6–8</sup>

The estimated treatment effects of simvastatin on NfL and NfH were small and not statistically significant, with the 95% CI sufficiently narrow to exclude an important treatment effect in either direction. Exploratory analyses did find that the month 24 serum NfL level increased by 6% (95% CI 2% to 9%) for each new/enlarging T2 lesion in the preceding year and by 0.9% (95% CI 0.3% to 1.5%) for each milliliter increase in concurrent T2LV, further supporting the known relationship between NfL and neuroinflammation. The key question of this study, however, was to determine the extent to which serum neurofilaments may act as biomarkers of a neuroprotective treatment that does not appear to have direct effects on neuroinflammation, using simvastatin as our example. Indeed, NfL has shown utility as a biomarker of noninflammatory neurodegeneration and neuroprotection in other neurologic conditions.<sup>33,34</sup> Our results, however, together with those from SPRINT-MS, suggest that either these treatments produce benefits on the rate of whole brain atrophy by mechanisms independent of neuroaxonal injury or that the degree of neuroprotection induced is insufficient to produce material



**Table 5** Relationship Between Baseline Serum NfL and Serum NfH and Clinical Variables

	Parameter estimate	95% CI	p Value
<b>EDSS score</b>			
<b>Relationship with baseline EDSS score</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	0.284	0.096 to 0.500	<0.01
Baseline NfH (per doubling)	-0.030	-0.109 to 0.058	>0.05
<b>Relationship with change in the EDSS score (units per year)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	0.026	-0.052 to 0.112	>0.05
Baseline NfH (per doubling)	0.002	-0.030 to 0.036	>0.05
<b>25FW</b>			
<b>Relationship with baseline 25FW (1/s)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	-0.159	-0.390 to 0.056	>0.05
Baseline NfH (per doubling)	0.055	-0.044 to 0.147	>0.05
<b>Relationship with change in 25FW (1/s per year)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	-0.183	-0.312 to -0.055	<0.01
Baseline NfH (per doubling)	0.028	-0.041 to 0.081	>0.05
<b>9HPT</b>			
<b>Relationship with baseline 9HPT (1,000/s)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	-3.600	-5.488 to -1.629	<0.01
Baseline NfH (per doubling)	-0.113	-1.267 to 1.011	>0.05
<b>Relationship with change in 9HPT (1,000/s per year)</b>			
<b>Predictor variable</b>			
Baseline NfL (per doubling)	-0.046	-0.904 to 0.858	>0.05
Baseline NfH (per doubling)	-0.168	-0.524 to 0.162	>0.05

Abbreviations: 9HPT = 9-hole peg test; 25FW = timed 25-foot walk; EDSS = Expanded Disability Status Scale; NfH = neurofilament heavy; NfL = neurofilament light.

The results of 6 separate models are presented, with EDSS, 25FW, or 9HPT as the dependent variables and NfL or NfH data as the predictor variables. EDSS score is reported as the score; both 9HPT and 25FW are reported as a speed (9HPT as  $1,000 \times s^{-1}$  and 25FW as  $s^{-1}$ ). In all analyses, neurofilament data were  $\log_2$  transformed. Results are from covariate-adjusted models as indicated in the Methods section. p Value bounds (</>0.05 and </>0.01) can only be inferred from the 95% and 99% bias-corrected and accelerated cluster bootstrap (10,000 replications) CIs.

changes in serum NfL. We speculate that the latter may be due to the association between neuroinflammation and NfL persisting independent of such neuroprotective treatment. Future work should focus on replicating this NfL analysis in samples from the ongoing phase 3 MS-STAT2 clinical trial, once the effects of simvastatin on clinical disability progression are confirmed, and also on developing novel CNS biomarkers capable of capturing neuroprotective treatment effects independent of neuroinflammation.

In 2 studies assessing the neuroprotective potential of sodium channel blockade (with phenytoin in acute optic neuritis and lamotrigine in SPMS), NfH has shown promise as a biomarker of neuroprotective treatment.<sup>11,12</sup> Our data, however, found no evidence of a simvastatin treatment effect on NfH or any associations of NfH with MRI or clinical measures of disease severity. Although previous studies have shown strong and consistent correlations between serum and CSF NfL,<sup>35</sup> inconsistent results have been found for correlation between

serum and CSF NfH.<sup>36-38</sup> This suggests that NfH instability between CSF and serum compartments may have limited its potential as a biomarker in our cohort. The Quanterix Simoa NF-Light Advantage assay has been widely used and validated against clinical and MRI outcomes. The Simoa pNF-heavy Discovery assay, however, has been less widely used. One study has used this assay to demonstrate modest associations between serum NfH and T2LV in a mixed MS cohort.<sup>38</sup> Although Simoa digital ELISA platforms tend to improve sensitivity and accuracy over traditional ELISA techniques, the limited data from this NfH assay therefore suggest that the NfH data should be interpreted with caution.

In conclusion, our results show that despite simvastatin treatment being associated with a significant reduction in whole brain atrophy and benefits in secondary outcomes, our results are most compatible with no important effect on serum NfL or NfH in SPMS. Although higher NfL is associated with greater disease severity and faster progression, our results, together with those from ibudilast, suggest that candidate nonimmunomodulatory neuroprotective treatments in PMS may act via mechanisms independent of the main determinants of serum neurofilament concentrations. While confirmation of the neuroprotective efficacy of simvastatin in SPMS is awaited from the ongoing phase 3 study, our results, together with those of others, suggest that the utility of serum neurofilaments as biomarkers of treatment response in progressive MS may be limited to interventions that are either known to suppress acute or chronic neuroinflammatory activity or to otherwise directly affect neuroaxonal injury.

## Study Funding

No targeted funding reported.

## Disclosure

T.E. Williams has received honorarium for educational talks from Novartis and Merck. K.P. Holdsworth, J.M. Nicholas, A. Eshaghi, T. Katsanouli, H. Wellington, and A. Heslegrave report no disclosures relevant to the manuscript. H. Zetterberg has served at scientific advisory boards for Denali, Roche Diagnostics, Wave, Samumed, Siemens Healthineers, Pinteon Therapeutics, Nervgen, AZTherapies, and CogRx; has given lectures in symposia sponsored by Cellectricon, Fujirebio, Alzecure, and Biogen; and is a cofounder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program. C. Frost reports no disclosures. J. Chataway has received support from the Efficacy and Mechanism Evaluation Programme and Health Technology Assessment Programme (NIHR); UK Multiple Sclerosis Society; the National Multiple Sclerosis Society; and the Rosetrees Trust. He is supported in part by the National Institute for Health Research, University College London Hospitals, Biomedical Research Centre, London, UK. He has been a local principal investigator for a trial in MS funded by the Canadian MS society; a local principal investigator for commercial trials funded by: Actelion, Biogen, Novartis and Roche; has received an investigator grant from Novartis; and has taken part in advisory boards/consultancy for Azadyne,

Biogen, Celgene, Janssen, MedDay, Merck, NervGen, Novartis and Roche. Go to [Neurology.org/NN](http://Neurology.org/NN) for full disclosures.

## Publication History

Received by *Neurology: Neuroimmunology & Neuroinflammation* June 16, 2021. Accepted in final form November 23, 2021.

## Appendix Authors

Name	Location	Contribution
<b>Thomas E. Williams, BA, MB BChir, MRCP</b>	Queen Square Multiple Sclerosis Centre, UCL Queen Square Institute of Neurology, United Kingdom	Acquisition of patient samples, neurofilament laboratory analysis, data analysis and interpretation, and drafting of the manuscript and revisions
<b>Katherine P. Holdsworth, MSc</b>	London School of Hygiene and Tropical Medicine, United Kingdom	Data analysis and interpretation and manuscript revisions
<b>Jennifer M. Nicholas, PhD</b>	London School of Hygiene and Tropical Medicine, United Kingdom	Data analysis and interpretation and manuscript revisions
<b>Arman Eshaghi, PhD</b>	Queen Square Multiple Sclerosis Centre, UCL Queen Square Institute of Neurology, United Kingdom	MRI analysis and manuscript revisions
<b>Theodora Katsanouli, MSc</b>	London School of Hygiene and Tropical Medicine, United Kingdom	Data analysis and interpretation and manuscript revisions
<b>Henrietta Wellington, PhD, FRCP</b>	UK Dementia Research Institute at UCL	Supervision of neurofilament laboratory analysis and manuscript revisions
<b>Amanda Heslegrave, PhD</b>	UK Dementia Research Institute at UCL	Supervision of neurofilament laboratory analysis and manuscript revisions
<b>Henrik Zetterberg, PhD</b>	UK Dementia Research Institute at UCL	Oversight of neurofilament laboratory analysis, data analysis and interpretation, and manuscript revisions
<b>Chris Frost, PhD</b>	London School of Hygiene and Tropical Medicine, United Kingdom	Supervision of data analysis and interpretation and manuscript revisions
<b>Jeremy Chataway, PhD, FRCP</b>	Queen Square Multiple Sclerosis Centre, UCL Queen Square Institute of Neurology, United Kingdom	Chief investigator of the MS-STAT clinical trial, study design and conception, data interpretation, and manuscript revisions

## References

- Lublin FD, Reingold SC, Cohen JA, et al. Defining the clinical course of multiple sclerosis. *Neurology*. 2014;83(3):278-286.
- Chataway J, Schuerer N, Alsanousi A, et al. Effect of high-dose simvastatin on brain atrophy and disability in secondary progressive multiple sclerosis (MS-STAT): a randomised, placebo-controlled, phase 2 trial. *Lancet*. 2014;383(9936):2213-2221.
- Chan D, Binks S, Nicholas JM, et al. Effect of high-dose simvastatin on cognitive, neuropsychiatric, and health-related quality-of-life measures in secondary progressive multiple sclerosis: secondary analyses from the MS-STAT randomised, placebo-controlled trial. *Lancet Neurol*. 2017;16(8):591-600.

4. Sormani MP, Haering DA, Kropshofer H, et al. Blood neurofilament light as a potential endpoint in Phase 2 studies in MS. *Ann Clin Transl Neurol.* 2019;6(6):1081-1089.
5. Williams T, Zetterberg H, Chataway J. Neurofilaments in progressive multiple sclerosis: a systematic review. *J Neurol.* 2021;268(9):3212-3222.
6. Bar-Or A, Thanei GA, Harp CT, et al. Blood neurofilament light levels are lowered to a healthy donor range in patients with RMS and PPMS following ocrelizumab treatment. Presented at the 35th Congress ofECTRIMS; 2019; Stockholm.
7. Kapoor R, Sellebjerg F, Hartung HP, et al. Natalizumab reduced serum levels of neurofilament light chain in secondary progressive multiple sclerosis patients from the phase 3 ASCEND study. Presented at the 34th Congress ofECTRIMS; 2018; Berlin.
8. Kuhle J, Kropshofer H, Haering D, et al. Neurofilament light levels in the blood of patients with secondary progressive MS are higher than in primary progressive MS and may predict brain atrophy in both MS subtypes. Presented at the 34th Congress ofECTRIMS; 2018; Berlin.
9. Fox R, Raska P, Barro C, et al. Neurofilament light chain in a phase 2 clinical trial of ibudilast in progressive multiple sclerosis. *Mult Scler.* 2021;27(13):2014-2022.
10. Kapoor R, Smith KE, Allegretta M, et al. Serum neurofilament light as a biomarker in progressive multiple sclerosis. *Neurology.* 2020;95(10):436-444.
11. Gnanapavan S, Grant D, Morant S, et al. Biomarker report from the phase II lamotrigine trial in secondary progressive MS—neurofilament as a surrogate of disease progression. *PLoS One.* 2013;8(8):e70019.
12. Raftopoulos R, Kuhle J, Grant D, et al. Neurofilament results for the phase II neuroprotection study of phenytoin in optic neuritis. *Eur J Neurol.* 2021;28(2):587-594.
13. Quanterix. Simoa NF-light Advantage Kit HD-1/HD-X data sheet. Epub 2017:1-2.
14. Quanterix. Simoa pNF-heavy Discovery Kit HD-1/HD-X data sheet. 2017:1-2.
15. Eshaghi A, Kievit RA, Prados F, et al. Applying causal models to explore the mechanism of action of simvastatin in progressive multiple sclerosis. *Proc Natl Acad Sci USA.* 2019;116(22):11020-11027.
16. Liu G, Lu K, Mogg R, Mallick M, Mehrotra D. Should baseline be a covariate or dependent variable in analyses of change from baseline in clinical trials? *Stat Med.* 2009;28(20):2509-2530.
17. European Medicines Agency. Statistical principles for clinical trials. *Eur Med Agency.* 1998;5(7):29.
18. Matthews J, Altman D, Campbell M, Royston P. Analysis of serial measurements in medical research. *Br Med J.* 1990;300(6725):230-235.
19. Frost C, Kenward MG, Fox NC. The analysis of repeated “direct” measures of change illustrated with an application in longitudinal imaging. *Stat Med.* 2004;23(21):3275-3286.
20. Barkhof F, Simon JH, Fazekas F, et al. MRI monitoring of immunomodulation in relapse-onset multiple sclerosis trials. *Nat Rev Neurol.* 2012;8(1):13-21.
21. World Medical Association. Declaration of Helsinki: ethical principles for medical research involving human subjects. 2013. Accessed January 2021. [wma.net/en/30publications/10policies/b3/17c.pdf](http://wma.net/en/30publications/10policies/b3/17c.pdf)
22. Maggi P, Kuhle J, Schädelin S, et al. Chronic white matter inflammation and serum neurofilament levels in multiple sclerosis. *Neurology.* 2021;97(6):e543–e553.
23. Ontaneda D, Thompson AJ, Fox RJ, Cohen JA. Progressive multiple sclerosis: prospects for disease therapy, repair, and restoration of function. *Lancet.* 2017;389(10076):1357-1366.
24. Palladino R, Marrie RA, Majeed A, Chataway J. Evaluating the risk of macrovascular events and mortality among people with multiple sclerosis in England. *JAMA Neurol.* 2020;77(7):820-828.
25. Marrie RA, Rudick R, Horwitz R, et al. Vascular comorbidity is associated with more rapid disability progression in multiple sclerosis. *Neurology.* 2010;74(13):1041-1047.
26. Jakimovski D, Gandhi S, Paunkoski I, et al. Hypertension and heart disease are associated with development of brain atrophy in multiple sclerosis: a 5-year longitudinal study. *Eur J Neurol.* 2019;26(1):87-e8.
27. Sormani MP, Arnold DL, De Stefano N. Treatment effect on brain atrophy correlates with treatment effect on disability in multiple sclerosis. *Ann Neurol.* 2014;75(1):43-49.
28. Sastre-Garriga J, Pareto D, Battaglini M, et al. MAGNIMS consensus recommendations on the use of brain and spinal cord atrophy measures in clinical practice. *Nat Rev Neurol.* 2020;16(3):171-182.
29. Cifelli A, Arridge M, Jezzard P, Esiri MM, Palace J, Matthews PM. Thalamic neurodegeneration in multiple sclerosis. *Ann Neurol.* 2002;52(5):650-653.
30. Popescu V, Klaver R, Voorn P, et al. What drives MRI-measured cortical atrophy in multiple sclerosis? *Mult Scler.* 2015;21(10):1280-1290.
31. Fox RJ, Coffey CS, Conwit R, et al. Phase 2 trial of ibudilast in progressive multiple sclerosis. *N Engl J Med.* 2018;379(9):846-855.
32. Barkhof F, Hulst HE, Drulovic J, Uitdehaag BMJ, Matsuda K, Landin R. Ibudilast in relapsing-remitting multiple sclerosis: a neuroprotectant? *Neurology.* 2010;74(13):1033-1040.
33. Bridel C, Van Wieringen WN, Zetterberg H, et al. Diagnostic value of cerebrospinal fluid neurofilament light protein in neurology: a systematic review and meta-analysis. *JAMA Neurol.* 2019;76(9):1035-1048.
34. Olsson B, Alberg L, Cullen NC, et al. NFL is a marker of treatment response in children with SMA treated with nusinersen. *J Neurol.* 2019;266(9):2129-2136.
35. Novakova L, Zetterberg H, Sundstrom P, et al. Monitoring disease activity in multiple sclerosis using serum neurofilament light protein. *Neurology.* 2017;89(22):2230-2237.
36. Vorobyeva A, Fominykh V, Zakharova M. Biochemical markers of neurodegeneration in multiple sclerosis. *Mult Scler.* 2013;19(11):369.
37. Eikelenboom MJ, Uitdehaag BMJ, Petzold A. Blood and CSF biomarker dynamics in multiple sclerosis: implications for data interpretation. *Mult Scler Int.* 2011;2011:1-6.
38. Verberk I, Koel-Simmelink M, Twaalfhoven H, et al. Ultrasensitive immunoassay allows measurement of serumneurofilament heavy in multiple sclerosis. *Mult Scler Relat Disord.* 2021;50:102840.

# Neurology<sup>®</sup> Neuroimmunology & Neuroinflammation

## Assessing Neurofilaments as Biomarkers of Neuroprotection in Progressive Multiple Sclerosis: From the MS-STAT Randomized Controlled Trial

Thomas E. Williams, Katherine P. Holdsworth, Jennifer M. Nicholas, et al.  
*Neurol Neuroimmunol Neuroinflamm* 2022;9;  
DOI 10.1212/NXI.0000000000001130

This information is current as of January 14, 2022

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://nn.neurology.org/content/9/2/e1130.full.html">http://nn.neurology.org/content/9/2/e1130.full.html</a>
<b>References</b>	This article cites 32 articles, 2 of which you can access for free at: <a href="http://nn.neurology.org/content/9/2/e1130.full.html##ref-list-1">http://nn.neurology.org/content/9/2/e1130.full.html##ref-list-1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>Class I</b> <a href="http://nn.neurology.org/cgi/collection/class_1">http://nn.neurology.org/cgi/collection/class_1</a> <b>Clinical trials Randomized controlled (CONSORT agreement)</b> <a href="http://nn.neurology.org/cgi/collection/clinical_trials_randomized_controlled_consort_agreement">http://nn.neurology.org/cgi/collection/clinical_trials_randomized_controlled_consort_agreement</a> <b>Multiple sclerosis</b> <a href="http://nn.neurology.org/cgi/collection/multiple_sclerosis">http://nn.neurology.org/cgi/collection/multiple_sclerosis</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://nn.neurology.org/misc/about.xhtml#permissions">http://nn.neurology.org/misc/about.xhtml#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://nn.neurology.org/misc/addir.xhtml#reprintsus">http://nn.neurology.org/misc/addir.xhtml#reprintsus</a>

*Neurol Neuroimmunol Neuroinflamm* is an official journal of the American Academy of Neurology. Published since April 2014, it is an open-access, online-only, continuous publication journal. Copyright Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology.. All rights reserved. Online ISSN: 2332-7812.

